



قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا  
عَلَّمْتَنَا إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ

صدق الله العظيم

سورة البقرة آية (٣٢)



---

---

# **Role of CT virtual endoscopy and multiplanar reformatting in the evaluation of laryngeal and hypopharyngeal carcinoma**

## **Thesis**

Submitted for Partial Fulfillment of MD Degree  
in **Radiodiagnosis**

**By**

**Heba Serag El-Din Apas**

M.B.,B.CH.,M.SC

Ain Shams University

**Under Supervision of**

**Prof. Dr. Sahar Naiem Mohammed**

Professor of Radiodiagnosis

Faculty of Medicine – Ain Shams University

**Prof. Dr. Lobna El-Fiky**

Professor of otorhinolaryngology

Faculty of Medicine – Ain Shams University

**Prof. Dr. Amany Emad El-Din Radi**

Professor of Radiodiagnosis

Faculty of Medicine – Ain Shams University

**Dr. Reem Hassan Basiouny**

Lecturer of Radiodiagnosis

Faculty of Medicine – Ain Shams University

**Faculty of Medicine**

**Ain Shams University**

**2012**

---

---

# Contents

---

	Page No.
List of tables .....	<i>i</i>
List of figures .....	<i>iv</i>
List of charts .....	<i>xx</i>
List of abbreviations .....	<i>xxii</i>
<b>Introduction .....</b>	<b>1</b>
<b>Aim of the Work .....</b>	<b>3</b>
<i>Review of literature</i>	
<b>1. Gross and CT anatomy of the larynx .....</b>	<b>4</b>
Gross and CT anatomy of the hypopharynx.....	22
Cervical lymph nodes.....	28
Normal CT virtual laryngoscopy anatomy.....	36
<b>2. Pathology of laryngeal and hypopharyngeal carcinoma .....</b>	<b>39</b>
<b>3. Physical Aspects of MDCT .....</b>	<b>63</b>
<b>4. Patients and Methods .....</b>	<b>84</b>
<b>5. Results .....</b>	<b>92</b>
<b>6. Illustrative Cases .....</b>	<b>119</b>
<b>7. Discussion.....</b>	<b>186</b>
<b>8. Summary and conclusion .....</b>	<b>195</b>
<b>9. Recommendations.....</b>	<b>200</b>
<b>10. References .....</b>	<b>201</b>
<b>Arabic Summary .....</b>	<b>—</b>

## List of Tables

<b>Table No.</b>	<b>Title</b>	<b>Page No.</b>
<b>1</b>	AJCC stage grouping for laryngeal cancer.	<b>57</b>
<b>2</b>	AJCC 2002 stage groupings for hypopharyngeal cancer.	<b>61</b>
<b>3</b>	Distribution of group I (laryngeal carcinoma) & group II (hypopharyngeal carcinoma) as regards age and gender	<b>93</b>
<b>4</b>	Distribution of group I (laryngeal carcinoma) & group II (hypopharyngeal carcinoma) as regards complaint	<b>94</b>
<b>5</b>	Distribution of group I (laryngeal carcinoma) & group II (hypopharyngeal carcinoma) as regards tumor staging	<b>95</b>
<b>6</b>	Distribution of group I (laryngeal carcinoma) & group II (hypopharyngeal carcinoma) as regards T score	<b>96</b>
<b>7</b>	Distribution of group I (laryngeal carcinoma) & group II (hypopharyngeal carcinoma) as regards lymph nodes staging	<b>97</b>
<b>8</b>	Distribution of patients in group I and II as regards tumoral extension	<b>98</b>
<b>9</b>	Comparison between endoscope and CTVL as regards mucosal surface irregularity. Fisher exact test. S= significant.	<b>100</b>
<b>10</b>	Comparison between direct endoscope and CTVL as regards primary subsite of the tumor. Chi-square test ( $X^2$ ).HS= highly significant.	<b>101</b>
<b>11</b>	Comparison between direct laryngoscopy and CTVL as regards anterior commissure involvement. Chi-square test ( $X^2$ ). S= significant.	<b>102</b>
<b>12</b>	Comparison between endoscope and CTVL as regards posterior commissure involvement. Chi-square test ( $X^2$ ).HS= highly significant.	<b>103</b>
<b>13</b>	Comparison between endoscope and CTVL as regards subglottic extension of the tumor. Chi-square test ( $X^2$ ).HS= highly significant	<b>104</b>

<b>14</b>	Comparison between direct endoscope and CTVL as regards vocal cords involvement. Chi-square test ( $X^2$ ).HS= highly significant.	<b>105</b>
<b>15</b>	Comparison between direct endoscope and CTVL as regards detection of the primary subsite of the tumor. Chi-square test ( $X^2$ ).HS= highly significant.	<b>107</b>
<b>16</b>	Comparison between direct endoscope and CTVL as regards involvement of the right pyriform sinus. Fisher exact test. S= significant.	<b>108</b>
<b>17</b>	Comparison between direct endoscope and CTVL as regards involvement of the left pyriform sinus. Fisher exact test. NS= non significant.	<b>109</b>
<b>18</b>	Comparison between direct endoscope and CTVL as regards extend of the tumor to posterior pharyngeal wall. Fisher exact test. NS= non significant.	<b>109</b>
<b>19</b>	Comparison between direct endoscope and CTVL as regards extent to post cricoid region. Fisher exact test. NS= non significant.	<b>110</b>
<b>20</b>	Comparison between direct endoscope and CTVL as regards extent of the tumor to vocal cords. Fisher exact test	<b>111</b>
<b>21</b>	Comparison between direct endoscope and CTVL as regards mucosal surface irregularity. Fisher exact test	<b>111</b>
<b>22</b>	Comparison between axial cuts and CT virtual laryngoscopy as regards subglottic extension of the tumor (group I ). Chi-square test ( $X^2$ ) . HS= highly significant.	<b>112</b>
<b>23</b>	Comparison between axial CT and MPR as regards LN staging in group I patients. Chi-square test ( $X^2$ ) . HS= highly significant.	<b>114</b>
<b>24</b>	Comparison between axial CT and MPR as regards lymph nodes enlargement and staging in group II patients. Fisher exact test. NS= non significant.	<b>115</b>
<b>25</b>	Comparison between axial cuts and coronal reformatted images as regards paraglottic space infiltration in group I& II. Fisher exact test. HS = highly significant, S= significant.	<b>116</b>

---

---

<b>26</b>	Comparison between axial CT cuts and sagittal CT cuts as regards preepiglottic space infiltration by the tumor in group I. Fisher exact test. HS= highly significant.	<b>117</b>
<b>27</b>	Comparison between axial CT and sagittal reconstruction as regards anterior commissure invasion in group I patients. Fisher exact test. HS= highly significant.	<b>117</b>
<b>28</b>	Comparison between axial CT and sagittal reformatting as regards detection of tumor extension to the posterior pharyngeal wall in group II patients. Fisher exact test. S= significant.	<b>118</b>

## List of Figures

Figure No.	Title	Page No.
<b>1.1</b>	(a) Anterolateral view of the laryngeal cartilages and ligaments. (b) Lateral view of the laryngeal cartilages and ligaments.	<b>5</b>
<b>1.2</b>	Posterior view of the cartilages and ligaments of the larynx.	<b>6</b>
<b>1.3</b>	Coronal section of larynx and upper part of trachea	<b>13</b>
<b>1.4</b>	Coronal reformatted image of CT of the larynx and upper part of trachea showing its three subdivisions lined by the black lines.	<b>13</b>
<b>1.5</b>	The hyoid bone (yellow arrow) and epiglottis (red arrow), the valleculae (asterix).	<b>15</b>
<b>1.6</b>	The hyoid bone (yellow arrow) and epiglottis (red arrow), the valleculae (asterix).	<b>15</b>
<b>1.7</b>	(a, b, c): supraglottic larynx: thyroid cartilage (red arrow), aryepiglottic fold (black arrow), pre epiglottic space (astrix), and para glottic space (orange arrow).	<b>15</b>
<b>1.8</b>	false vocal folds (arrows).	<b>15</b>
<b>1.9</b>	(a, b): true vocal folds (arrows).	<b>17</b>
<b>1.10</b>	(a,b,c): showing progressive change in shape of air column of subglottic larynx. Thyroid cartilage (red arrow), cricoids cartilage (blue arrow).	<b>19</b>
<b>1.11</b>	Hypopharyngeal anatomy.	<b>22</b>
<b>1.12</b>	Posterior view of the pharynx.	<b>23</b>
<b>1.13</b>	Subdivisions of the pharynx. Sagittal CT scan showing subdivisions of the pharynx (demarcated by dashed lines). The nasopharynx ( <b>NP</b> ) extends superiorly to the skull base and inferiorly to the level of the hard palate. The oropharynx ( <b>OP</b> ) extends from the inferior border of the nasopharynx to the level of the hyoid bone and the hypopharynx ( <b>HP</b> ) continues inferiorly to the caudal margin of the <b>cricoid</b> cartilage.	<b>23</b>
<b>1.14</b>	showing upper pyriform sinus (asterix) and posterior hypopharyngeal wall (arrow).	<b>25</b>

<b>1.15</b>	showing mid pyriform sinus bounded by aryepiglottic folds medially and posterior pharyngeal wall posteriorly.	<b>25</b>
<b>1.16</b>	showing lower pyriform sinus.	<b>25</b>
<b>1.17</b>	showing cricoarytenoid joint (white arrow) and posterior wall of post cricoid area (small black arrows).	<b>26</b>
<b>1.18</b>	showing attachment of postcricoid musculature to inferior horn of thyroid cartilage at the level of middle of cricoid cartilage (black arrows).	<b>27</b>
<b>1.19</b>	showing Inferior margin of cricoid cartilage. Note the change in shape of muscular wall with lack of intra mural fat planes.	<b>27</b>
<b>1.20</b>	Schematic Illustration of the Nodal Classification.	<b>28</b>
<b>1.21</b>	(a): Axial images show the inferior margin of the <b>Level I</b> lymph nodes that are formed by the hyoid bone. Its superior margin is formed by mylohyoid muscle (b): The posterior margin of the Level I nodes is formed by white line connecting the posterior margin of the submandibular glands. <b>Level IA</b> lymph nodes are located between the mylohyoid muscles (arrowheads = mylohyoid muscle, small arrow = location of Level IA nodes). The <b>Level IB</b> nodes are located lateral to the mylohyoid muscles (large white arrows).	<b>29</b>
<b>1.22</b>	<b>A.</b> The superior margin of the <b>Level II</b> lymph nodes is formed by the skull base jugular foramen (arrow), <b>B.</b> The lower margin is formed by the hyoid bone (arrow), <b>C.</b> The Level II nodes are located anterior to a transverse line connecting the posterior edge of the sternocleidomastoid muscles (long line) and posterior to a transverse line connecting the posterior edge of the submandibular glands (small line). <b>D.</b> <b>Level IIA</b> nodes are situated anterior, medial or lateral to the internal jugular vein (arrow) whereas <b>Level IIB</b> nodes are posterior to the internal jugular vein and have an identifiable fat plane between the lymph node and the vein (arrowheads).	<b>30</b>



<b>1.23</b>	<p><b>A.</b> The hyoid bone denotes the superior extent of the <b>Level III</b> nodes (arrow). <b>B.</b> The cricoid cartilage arch forms the lower border (arrow). <b>C.</b> Level III lymph nodes are situated anterior to a "line" connecting the posterior margins of the sternocleidomastoid muscle and lateral to the medial margin of either the common carotid artery or the internal carotid artery. (arrowhead = internal carotid arteries).</p>	<b>31</b>
<b>1.24</b>	<p><b>A.</b> The cricoid cartilage (arrow) forms the superior boundary of the <b>Level IV</b> nodes. <b>B.</b> The clavicle forms the inferior margin (arrow). <b>C.</b> These nodes are anterior and medial to an "oblique line" drawn connecting the posterior edge of the sternocleidomastoid muscle and the posterolateral edge of the anterior scalene muscle and lateral to the carotid artery (arrowheads).</p>	<b>32</b>
<b>1.25</b>	<p><b>A.</b> The skull base forms the superior margin of the <b>Level V</b> nodes.  <b>B.</b> This group of nodes extends inferior to the level of the clavicle (arrow).  <b>C. Level VA</b> nodes lie between the skull base and the level of the lower margin of the cricoid cartilage arch. Unlike Levels II and III, Level VA nodes are behind the posterior edge of the sternocleidomastoid muscle (arrowheads) and anterior to the trapezius muscle (arrows).  <b>D. Level VB</b> nodes are located between the lower margin of the cricoid cartilage and the clavicle. Unlike Level IV nodes, they are behind an oblique line connecting the posterior edge of the sternocleidomastoid muscle and the posterolateral edge of the anterior scalene muscle.</p>	<b>33</b>
<b>1.26</b>	<p><b>A.</b> The superior border is formed by the hyoid bone.  <b>B.</b> This group of nodes extends inferiorly to the top of the manubrium.  <b>C.</b> These nodes are located between the medial margins of the left and right common (arrowheads) or internal carotid arteries.</p>	<b>34</b>

<b>1.27</b>	These nodes are situated at or caudal to the clavicles (arrows) and lateral to the carotid arteries (arrowheads).	<b>35</b>
<b>1.28</b>	<b>Level VII</b> nodes lie caudal to the top of the manubrium (arrowhead), between the medial margins of the left and right common carotid arteries, and extend caudally to the level of the innominate vein. (Arrow=unopacified innominate vein).	<b>35</b>
<b>1.29</b>	showing virtual endoscopy from upside down at the level of epiglottis. Tip of epiglottis (arrow) and vallecula (asterix).	<b>36</b>
<b>1.30</b>	virtual scope from upside down behind the epiglottis showing the eminence for arytenoids posteriorly (arrows) with pyriform sinuses (asterix).	<b>36</b>
<b>1.31</b>	virtual scope from upside down at a lower level showing two false vocal folds (white arrow) and true vocal folds below them (black arrow).	<b>37</b>
<b>1.32</b>	virtual scope from upside down at a closer view to the true vocal cords showing the anterior commissure (arrow).	<b>37</b>
<b>1.33</b>	virtual scope from upside down at the subglottic level showing the tracheal rings.	<b>37</b>
<b>1.34</b>	virtual scope from downwards up showing the undersurface of true vocal cords (arrow).	<b>38</b>
<b>1.35</b>	virtual scope from downwards up with a closer view to the true vocal cords showing the anterior commissure (arrow).	<b>38</b>
<b>1.36</b>	virtual scope from downwards up above the vocal folds showing the undersurface of epiglottis (arrow).	<b>38</b>
<b>2.1</b>	Basaloid squamous cell carcinoma. A lobular (solid) and cord-like patterns. B, central comedotype. C, basaloid-appearing cells showing marked pleomorphism. D, the squamous cell component is focally seen in the form of keratinization.	<b>41</b>

<b>2.2</b>	Laryngeal papillary squamous cell carcinoma. <b>A</b> , filiform growth with finger-like projections and identifiable fibrovascular cores. <b>B</b> , broad-based bulbous to exophytic growth with rounded projections resembling a cauliflower-like growth pattern with fibrovascular cores. <b>C</b> , the squamous epithelium is cytologically malignant.	<b>42</b>
<b>2.3</b>	<b>A</b> , laryngeal spindle cell squamous carcinoma showing a differentiated squamous cell carcinoma ( <i>top</i> ) associated with a malignant (undifferentiated) spindle cell proliferation. <b>B</b> , the malignant cell component of spindle cell squamous carcinoma may include ( <i>left</i> ) spindle-shaped cells and ( <i>right</i> ) epithelioid cells.	<b>43</b>
<b>2.4</b>	Laryngeal verrucous carcinoma (VC). <b>A</b> , squamous cell proliferation with marked surface keratinization. <b>B</b> , VC is characterized by bland, uniform cells showing maturation, retention of polarity, no dysplasia nor nuclear pleomorphism, absence of increased nuclear-to-cytoplasmic ratio, and absence of dyskeratosis.	<b>44</b>
<b>2.5</b>	A well-differentiated neuroendocrine carcinoma ("carcinoid" tumor) of the larynx consists of sharply demarcated nests of uniform cells.	<b>46</b>
<b>2.6</b>	<b>1.</b> Primary renal cell carcinoma removed 7 years earlier. <b>2.</b> Metastatic renal cell carcinoma of the larynx with intact overlying squamous mucosa.	<b>47</b>
<b>3.1</b>	Diagram shows the difference between single-row detector and multiple-row detector CT designs. The multiple-row detector array shown is asymmetrical and represents that of one particular manufacturer.	<b>64</b>
<b>3.2</b>	Anatomic coverage. ( <b>a</b> , <b>b</b> ) The coverage for multisection CT ( <b>a</b> ) can be eight times longer than for single-section helical CT ( <b>b</b> ) at the same pitch and section thickness. ( <b>c</b> , <b>d</b> ) To scan the same volume in the same time with single-section helical CT, one must increase the pitch ( <b>c</b> ) or section thickness ( <b>d</b> ), thereby degrading image quality. ( <b>e</b> ) To achieve the same image quality with single-section helical CT, the scanning time would have to be lengthened eightfold.	<b>68</b>

<b>3.3</b>	Diagram illustrates how a sequence of axial images can be considered to form a vertical “stack.” By sampling a three-dimensional stack of CT numbers along the y-z plane, sagittal views can be generated. Similarly, sampling in the x-z plane generates coronal views. The CT reformatting process is thus fundamentally different (and simpler) than the CT reconstruction process (in which raw CT projection signal data are used to create an axial image), and these two terms ( <i>reformat</i> and <i>reconstruction</i> ) should not be used interchangeably.	<b>67</b>
<b>3.4</b>	Effects of AIP on an image of the liver. <b>(a)</b> Coronal reformatted image created with a default thickness of 1 pixel (approximately 0.8 mm). <b>(b)</b> Increasing the slab thickness to 4 mm by using AIP results in a smoother image with less noise and improved contrast resolution. The image quality is similar to that used in axial evaluation of the abdomen.	<b>71</b>
<b>3.5</b>	<b>(a)</b> Oblique maximum-intensity projection of a slab (cropped volume) of the hepatic and pancreatic regions clearly demonstrates the enhancing regional vasculature and orally administered gastrointestinal contrast material. The image clearly shows vascular stenosis (arrows) caused by a surrounding tumor, but the tumor margins are not well seen. <b>(b)</b> Oblique minimum-intensity projection (slab view) of the same patient emphasizes the hepatic biliary tree. The tumor margins (arrows) are delineated better with this technique. A slightly different slab view volume was used in this view.	<b>72</b>
<b>3.6</b>	<b>(a)</b> Surface-rendered view from a multiple-row detector CT series demonstrates an aneurysm (arrow) arising at the origin of the posterior inferior cerebellar artery (arrowhead) from the vertebral artery (circle). <b>(b)</b> On a view from the interior of the artery, the left vertebral artery ( <i>L Vert</i> ), posterior inferior cerebellar artery ( <i>PICA</i> ), and neck of the aneurysm are all easily recognized, and their relative locations are readily appreciated.	<b>74</b>

<b>3.7</b>	(a) Frontal volume-rendered image of a patient with horseshoe kidney who was scheduled to undergo surgery for a mass in the right posterior-inferior aspect of the fused kidney (arrowheads). The aorta is readily apparent (arrow). (b) Another view is somewhat closer and more superior than the view seen in a, and the kidney tissue has been rendered slightly more transparent. The unusual vasculature pattern in this case is readily apparent, with two upper pole arteries (open arrows) and two lower pole arteries (arrowheads) originating below the inferior mesenteric artery (solid arrow).	<b>76</b>
<b>3.8</b>	(a) Volume-rendered image for which the color map function was adjusted to help enhance the visibility of the tumors (arrows) on the cortex of the kidney. The kidney is shown as orange, the renal lesions are reddish, and the bones are white. Observing this case in motion (rotating right to left) helped the viewer appreciate the depth and interconnections of the anatomic structures. (b) An alternate frame of the movie loop from this case demonstrates the renal artery structure.	<b>77</b>
<b>3.9</b>	Orthographic volume rendering of the airways. Volume-rendered image of a patient with tracheal stenosis (arrow) includes the airway from the hypopharynx to just above the carina. The image is not distorted by proximity or angle of the viewpoint and provides an “external” view of anatomic relationships. Segmentation of the airways was achieved by assigning a spike in opacity at the interface between air and soft-tissue attenuation. Overlying lung tissue was removed with region-of-interest editing to avoid obscuring the trachea.	<b>78</b>

<b>3.10</b>	Perspective volume rendering of the airways. Axial (top right), coronal (bottom left), and sagittal (bottom right) chest CT scans show a left hilar mass, which is positioned between vascular structures. Virtual bronchoscopy (immersive rendering with a point of view within the tracheobronchial tree) (top left) was used to guide subsequent transbronchial biopsy, allowing six biopsy passes between central vascular structures without significant bleeding.	<b>80</b>
<b>3.11</b>	): Region-of-interest editing. (a) Three-dimensional volume-rendered image (posterior view) from chest CT performed in a trauma patient with a fracture of T10. A region including a portion of the left ribs is defined manually (green area). (b) The selected region is then removed from the image. (c) Rib removal allows visualization of the fracture on a lateral projection without interference from overlying ribs.	<b>82</b>
<b>3.12</b>	Use of an opacity threshold for segmentation, as shown on a full field of view 3D volume-rendered image of the chest and abdomen. (a) A low opacity threshold allows the skin to obscure the abdominal contents. A vertical row of shirt buttons is seen in the midline. (b–d) Progressively increasing the opacity threshold excludes first low-opacity soft tissues (skin, fat) (b) then high-opacity soft tissues (muscle, bowel wall) (c) while contrast-enhanced organs and vessels remain (d). (e) Eventually only the most opaque objects (bone, calcium, excreted contrast material) remain.	<b>83</b>
<b>6.1</b>	axial CT cuts are showing mass infiltrating both pyriform sinuses, post cricoid region (A) and posterior pharyngeal wall(B).	<b>119</b>
<b>6.2</b>	(A & B)sagittal reformatted CT images reveal no infiltration of the pre-epiglottic space. The lower limit of the mass is noted opposite D2 vertebral body. (C)Coronal CT image is showing clear paraglottic space bilaterally.	<b>120</b>
<b>6.3</b>	(A& B): coronal CT cuts are showing bilateral deep cervical lymph node enlargement.	<b>120</b>

<b>6.4</b>	Virtual laryngoscopy images reveals obliteration of both pyriform fosse (A) with encroachment on the posterior wall of the larynx (B) yet mucosal surface irregularity couldn't be evaluated.	<b>121</b>
<b>6.5</b>	(A) routine axial CT reveals mass at the left pyriform fossa and, (B) CT done with modified valsalva maneuver is showing the mass arising from its lateral wall .	<b>122</b>
<b>6.6</b>	axial CT images are showing the mass extending to the post cricoid region (A) and posterior pharyngeal wall (B).	<b>123</b>
<b>6.7</b>	(A) sagittal CT image reveals no infiltration of the pre-epiglottic space. Also lower limit of the mass is noted opposite D1 vertebral body. (B) Axial CT image is showing clear paraglottic space bilaterally.	<b>123</b>
<b>6.8</b>	(A & B) Coronal and axial CT images showing bilateral deep cervical lymph node more than 10 mm on the left.	<b>123</b>
<b>6.9</b>	Virtual laryngeoscopy images reveal obliteration of left pyriform fossa with clear right pyriform fossa (A) and encroachment on the posterior wall of the larynx (B) with mucosal surface irregularity noted.	<b>124</b>
<b>6.10</b>	(A & B): CT axial cuts are showing mass at the left pyriform fossa.	<b>125</b>
<b>6.11</b>	(A) coronal reformatted image reveals infiltration of the left paraglottic space. (B) Sagittal reformatted image reveals clear pre-epiglottic space with the lower limit of the mass is noted opposite D1 vertebral body.	<b>126</b>
<b>6.12</b>	Bilateral deep cervical lymph node enlargement is noted in the coronal image (A) at left side and in the sagittal image (B) at the right side.	<b>126</b>
<b>6.13</b>	Virtual laryngoscopy cranio-caudal image (A) is showing mucosal surface irregularity with free vocal cords noted in caudo-cranial image (B).	<b>126</b>
<b>6.14</b>	axial CT cuts showing the mass arising from left pyriform sinus (A) extending to the post cricoid region (B) and the posterior pharyngeal wall (C).	<b>129</b>
<b>6.15</b>	sagittal reformatted image reveals the lower limit of the mass opposite D1/D2.	<b>129</b>