

Analytical Study on Some Drugs Used for Treatment of Type 2 Diabetes Mellitus

Thesis presented by
Maya Shaaban Eissa Hussein

B.Sc. in Pharmaceutical Sciences
Faculty of Pharmacy - Ain Shams University
2006

Submitted for the partial fulfillment of
Master Degree in Pharmaceutical Sciences
(Pharmaceutical Analytical Chemistry)

Under supervision of:

Prof. Dr. Osama Ibrahim Abd El-Sattar

Professor of Pharmaceutical Analytical Chemistry
Chairman of the National Organization of Drug Control and Research

Associate Prof. Dr. Maha Farouk Abd El-Ghany

Associate Professor of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy - Ain Shams University

Dr. Omar Abd El-Aziz Ghonim

Lecturer of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy - Ain Shams University

Pharmaceutical Analytical Chemistry Department
Faculty of Pharmacy - Ain Shams University
2011

Approval Sheet

*Analytical Study on Some Drugs Used for
Treatment of Type 2 Diabetes Mellitus*

Name of the Candidate:

Maya Shaaban Eissa Hussein

Submitted to:

Pharmaceutical Analytical Chemistry Department, Faculty of
Pharmacy, Ain Shams University, 2011

Committee in Charge:

Prof. Dr. Osama Ibrahim Abd El-Sattar

.....
Professor of Pharmaceutical Analytical Chemistry
Chairman of the National Organization of Drug Control and Research

Prof. Dr. Hamed Mahmoud El-Fatattary

.....
Professor of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy, Tanta University

Prof. Dr. Sawsan Abd El-Moneim Abd El-Razik

.....
Professor of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy, Al-Azhar University (Girls)

Associate Prof. Dr. Maha Farouk Abd El-Ghany

.....
Associate Professor of Pharmaceutical Analytical Chemistry
Faculty of Pharmacy, Ain Shams University

Date: Monday, 10/ 01/ 2011

ACKNOWLEDGEMENT

*No words can ever express my sincere gratitude to **ALLAH** who guide, aid and bless me in everything and everywhere in my life.*

*I am deeply grateful to **Prof. Dr. Osama Ibrahim Abd El-Sattar**, Professor of Pharmaceutical Analytical Chemistry and Chairman of the National Organization of Drug Control and Research, for suggesting the point. His supervision, valuable advice, constructive comments and helpful suggestions are the things that I wish to express my sincere appreciation and my deepest thanks and gratitude. It's really an honour to be one of his students.*

*I am greatly indebted to **Associate Prof. Dr. Maha Farouk Abd El-Ghany**, Associate Professor of Pharmaceutical Analytical Chemistry, Faculty of Pharmacy, Ain Shams University, for her valuable supervision, beneficial discussion, great effort and continuous encouragement.*

*I also like to express my sincere thanks to **Dr. Omar Abd El-Aziz Ghonim**, Lecturer of Pharmaceutical Analytical Chemistry, Faculty of Pharmacy, Ain shams University, for his keen interest, kind guidance, constant advice and sincere help in all stages of the work,*

*I wish to express my cordial thanks and gratefulness to **my parents, my sister and my brother** for their continuous encouragement and praying and their effective support.*

*Finally, my hearty thanks to all **Staff of ACAP I pharmaceutical factory**, Badr city, for their keen help in carrying out my pratical work,*

Maya Shaaban

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

(نَرْفَعُ دَرَجَاتٍ مَن نَّشَاءُ)
وَفَوْقَ كُلِّ ذِي عِلْمٍ عَظِيمٌ

صدق الله العظيم

سورة يوسف آية 76

CONTENTS

PART I: Introduction

Section A

General Introduction

	Page
I.A.1.Diabetes Mellitus.....	1
I.A.2. Oral Hypoglycaemic Drugs.....	2
I.A.2.1. Drug Stability.....	3

Section B

Literature Review

I.B.1. Chemical structures and physicochemical properties.....	7
I.B.1.1. Repaglinide.....	7
I.B.1.2. Pioglitazone hydrochloride.....	7
I.B.1.3. Rosiglitazone maleate.....	8
I.B.2. Stability.....	8
I.B.3. Pharmacology.....	9
I.B.4. Structure Activity Relationship.....	11
I.B.5. Methods of Analysis.....	14
I.B.5.1. Repaglinide	14
I.B.5.1.1. Spectroscopic Methods	14
I.B.5.1.2. Electrochemical Methods.....	15
I.B.5.1.3 Chromatographic Methods.....	15
I.B.5.1.3.1. Thin Layer Chromatography.....	15
I.B.5.1.3.2. High Performance Liquid Chromatography.....	15
I.B.5.2.Pioglitazone hydrochloride.....	18
I.B.5.2.1. Spectroscopic Methods.....	18
I.B.5.2.2. Electrochemical Methods.....	19
I.B.5.2.3. Capillary Electrophoresis.....	20
I.B.5.2.4. Chromatographic Methods.....	20
I.B.5.2.4.1. Thin Layer Chromatography.....	20
I.B.5.2.4.2. High Performance Liquid Chromatography.....	20
I.B.5.3.Rosiglitazone maleate.....	25
I.B.5.3.1. Spectroscopic Methods.....	25
I.B.5.3.2. Electrochemical Methods.....	25
I.B.5.3.3. Capillary Electrophoresis.....	26
I.B.5.3.4. Chromatographic Methods	26
I.B.5.3.4.1. Thin Layer Chromatography.....	26
I.B.5.3.4.2. High Performance Liquid Chromatography.....	26

Part II: Chromatographic Methods

Introduction.....	31
-------------------	----

Section A

Chromatographic Stability-indicating Method for Determination of Repaglinide in Presence of its ‘Acid, Alkaline and Oxidative-Degradates’ and Their Structure Elucidation, Using Liquid chromatography-mass spectrometry Technique

II.A.1.Experimental.....	34
II.A.1.1. Instrumentation.....	34
II.A.1.2. Samples.....	35
II.A.1.3. Chemicals and Reagents.....	35
II.A.1.4. Repaglinide standard solutions.....	36
II.A.1.5. Preparation of the degraded solutions (100.00 µg.ml ⁻¹).....	36
II.A.1.6. Procedure.....	37
II.A.1.6.1. Chromatographic conditions.....	37
II.A.1.6.2. Conditions of liquid chromatography-mass spectrometry for structural elucidation.....	37
II.A.1.6.3. Method validation.....	38
II.A.1.6.3.1. Linearity.....	38
II.A.1.6.3.2. Accuracy.....	38
II.A.1.6.3.3. Precision.....	39
II.A.1.6.3.4. Specificity.....	39
II.A.1.6.3.5. Robustness.....	39
II.A.1.6.3.6. Ruggedness.....	40
II.A.1.6.4. Application to the pharmaceutical preparation.....	40
II.A.1.6.5. Validation by standard addition technique	40
II.A.2. Results and Discussion.....	42
II.A.2.1. Choice of mobile phase organic modifier.....	42
II.A.2.2. Importance of adding an ion-pairing agent, Triethylamine.....	43
II.A.2.3. Mobile phase composition.....	44
II.A.2.4. Results of method validation.....	45
II.A.2.5. Structural elucidation of repaglinide degradates using liquid chromatography-mass spectrometry technique.....	58
II.A.2.6. Proposed degradation pathways of repaglinide.....	58

Section B
Chromatographic Stability-indicating Method for Determination of
Pioglitazone Hydrochloride in Presence of its ‘Acid, Alkaline and
Oxidative-Degradates’ and Their Structure Elucidation, Using Liquid
chromatography-mass spectrometry Technique

II.B.1.Experimental.....	65
II.B.1.1. Instrumentation.....	65
II.B.1.2.Samples.....	65
II.B.1.3. Chemicals and Reagents.....	65
II.B.1.4. Pioglitazone hydrochloride standard solution.....	66
II.B.1.5. Preparation of the degraded solutions (100.00 µg.ml ⁻¹).....	66
II.B.1.6. Procedure.....	67
II.B.1.6.1. Chromatographic conditions.....	67
II.B.1.6.2. Conditions of liquid chromatography-mass spectrometry for structural elucidation.....	67
II.B.1.6.3. Method validation.....	68
II.B.1.6.3.1.Linearity.....	68
II.B.1.6.3.2.Accuracy.....	68
II.B.1.6.3.3.Precision.....	68
II.B.1.6.3.4.Specificity.....	69
II.B.1.6.3.5.Robustness.....	69
II.B.1.6.3.6.Ruggedness.....	69
II.B.1.6.4. Application to the pharmaceutical preparation.....	70
II.B.1.6.5. Validation by standard addition technique	70
II.B.2. Results and Discussion.....	71
II.B.2.1. Choice of mobile phase organic modifier.....	71
II.B.2.2. Importance of adding an ion-pairing agent, Triethylamine.....	72
II.B.2.3. Mobile phase composition.....	72
II.B.2.4. Results of method validation.....	73
II.B.2.5. Structural elucidation of pioglitazone hydrochloride degradates using liquid chromatography-mass spectrometry technique.....	86
II.B.2.6. Proposed degradation pathways of pioglitazone hydrochloride.....	86

Section C
**Chromatographic Stability-indicating Method for Determination of
Rosiglitazone Maleate in Presence of its ‘Acid, Alkaline and Oxidative-
Degradates’ and Their Structure Elucidation, Using Liquid
chromatography-mass spectrometry Technique**

II.C.1. Experimental.....	93
II.C.1.1 Instrumentation.....	93
II.C.1.2. Samples.....	93
II.C.1.3. Chemicals and Reagents.....	93
II.C.1.4. Rosiglitazone maleate standard solutions.....	94
II.C.1.5. Preparation of the degraded solutions (100.00 µg.ml ⁻¹).....	94
II.C.1.6.Procedure.....	95
II.C.1.6.1. Chromatographic conditions.....	95
II.C.1.6.2. Conditions of liquid chromatography-mass spectrometry for structural elucidation.....	95
II.C.1.6.3. Method validation.....	96
II.C.1.6.3.1. Linearity.....	96
II.C.1.6.3.2. Accuracy.....	96
II.C.1.6.3.3. Precision.....	96
II.C.1.6.3.4. Specificity.....	97
II.C.1.6.3.5. Robustness.....	97
II.C.1.6.3.6. Ruggedness.....	97
II.C.1.6.4. Application to the pharmaceutical preparation.....	98
II.C.1.6.5. Validation by standard addition technique	98
II.C.2. Results and Discussion.....	99
II.C.2.1. Choice of mobile phase organic modifier.....	99
II.C.2.2. Importance of adding an ion-pairing agent, Triethylamine.....	100
II.C.2.3. Mobile phase composition.....	100
II.C.2.4. Results of method validation.....	101
II.C.2.5.Structural elucidation of rosiglitazone maleate degradates using liquid chromatography-mass spectrometry technique.....	114
II.C.2.6. Proposed degradation pathways of rosiglitazone maleate.....	114

Part III: Spectrophotometric Stability-indicating Methods

Introduction.....	122
--------------------------	------------

Section A
Spectrophotometric Stability-indicating Methods for Determination of Repaglinide in Presence of its Acid, Alkaline and Oxidative-Degradates, Using Derivative and pH-induced Difference Techniques

III.A.1.Experimental.....	124
III.A.1.1. Instrumentation.....	124
III.A.1.2. Samples.....	124
III.A.1.3. Chemicals and Reagents.....	124
III.A.1.4. Repaglinide standard solutions.....	125
III.A.1.5. Preparation of the degraded solutions (100.00 $\mu\text{g.ml}^{-1}$).....	125
III.A.1.6. Procedure.....	125
III.A.1.6.1. Spectral characteristics of repaglinide and its ‘acid, alkaline and oxidative degradates’	125
III.A.1.6.1.a. Derivative spectrophotometry	125
III.A.1.6.1.b. pH-induced difference spectrophotometry.....	125
III.A.1.6.2. Method validation.....	126
III.A.1.6.2.1. Linearity.....	126
III.A.1.6.2.2. Accuracy.....	127
III.A.1.6.2.3. Precision.....	127
III.A.1.6.2.4. Specificity.....	128
III.A.1.6.2.5. Robustness of the pH-induced difference spectrophotometric method	128
III.A.1.6.2.6. Ruggedness.....	128
III.A.1.6.3. Application to the pharmaceutical preparation.....	128
III.A.1.6.4. Validation by standard addition technique	129
III.A.2. Results and Discussion.....	129

Section B
Spectrophotometric Stability-indicating Methods for Determination of Pioglitazone Hydrochloride in Presence of its Acid, Alkaline and Oxidative-Degradates, Using Derivative and pH-induced Difference Techniques

III.B.1. Experimental.....	151
III.B.1.1 Instrumentation.....	151
III.B.1.2. Samples.....	151
III.B.1.3. Chemicals and Reagents.....	151
III.B.1.4. Pioglitazone hydrochloride standard solutions.....	151
III.B.1.5. Preparation of the degraded solutions (100.00 $\mu\text{g.ml}^{-1}$).....	151
III.B.1.6. Procedure.....	152

III.B.1.6.1. Spectral characteristics of pioglitazone hydrochloride and its ‘acid, alkaline and oxidative-degradates’	152
III.B.1.6.1.a. Derivative spectrophotometry	152
III.B.1.6.1.b. pH-induced difference spectrophotometry	152
III.B.1.6.2. Method validation.....	153
III.B.1.6.2.1. Linearity.....	153
III.B.1.6.2.2. Accuracy.....	154
III.B.1.6.2.3. Precision.....	154
III.B.1.6.2.4. Specificity.....	155
III.B.1.6.2.5. Robustness of the pH-induced difference spectrophotometric method	155
III.B.1.6.2.6. Ruggedness.....	155
III.B.1.6.3. Application to the pharmaceutical preparation.....	155
III.B.1.6.4. Validation by standard addition technique	156
III.B.2. Results and Discussion.....	156

Section C

Spectrophotometric Stability-indicating Methods for Determination of Rosiglitazone Maleate in Presence of its Acid, Alkaline and Oxidative-Degradates, Using Derivative and pH-induced Difference Techniques

III.C.1. Experimental.....	178
III.C.1.1 Instrumentation.....	178
III.C.1.2. Samples.....	178
III.C.1.3. Chemicals and Reagents.....	178
III.C.1.4. Rosiglitazone maleate standard solutions.....	178
III.C.1.5. Preparation of the degraded solutions (100.00 $\mu\text{g.ml}^{-1}$).....	178
III.C.1.6. Procedure.....	179
III.C.1.6.1. Spectral characteristics of rosiglitazone maleate and its ‘acid, alkaline and oxidative-degradates’	179
III.C.1.6.1.a. Derivative spectrophotometry	179
III.C.1.6.1.b. pH-induced difference spectrophotometry.....	179
III.C.1.6.2. Method validation.....	180
III.C.1.6.2.1. Linearity.....	180
III.C.1.6.2.2. Accuracy.....	181
III.C.1.6.2.3. Precision.....	181
III.C.1.6.2.4. Specificity.....	181
III.C.1.6.2.5. Robustness of the pH-induced difference spectrophotometric method	182
III.C.1.6.2.6. Ruggedness.....	182

III.C.1.6.3. Application to the pharmaceutical preparation.....	182
III.C.1.6.4. Validation by standard addition technique	182
III.C.2. Results and Discussion.....	183

Part IV: Visible Spectrophotometric Methods

Introduction.....	205
--------------------------	------------

Section A

Spectrophotometric Methods for Determination of Repaglinide, Using Ion-pair and Charge-transfer Techniques

IV.A.1. Experimental.....	208
IV.A.1.1 Instrumentation.....	208
IV.A.1.2. Samples.....	208
IV.A.1.3. Chemicals and Reagents.....	208
IV.A.1.4. Repaglinide standard solutions.....	209
IV.A.1.5. Procedure.....	210
IV.A.1.5.1. Spectral characteristics of repaglinide produced complexes.....	210
IV.A.1.5.1.a. Repaglinide ion-pair complex with bromophenol blue.....	210
IV.A.1.5.1.b. Repaglinide charge-transfer complex with chloranilic acid.....	210
IV.A.1.5.2. Method validation.....	211
IV.A.1.5.2.1. Linearity.....	211
IV.A.1.5.2.2. Accuracy.....	212
IV.A.1.5.2.3. Precision.....	212
IV.A.1.5.2.4. Robustness.....	212
IV.A.1.5.2.5. Ruggedness.....	213
IV.A.1.5.3. Application to the pharmaceutical preparation.....	213
IV.A.1.5.4. Validation by standard addition technique	213
IV.A.2. Results and Discussion.....	214
IV.A.2.1. Method Development.....	214
IV.A.2.1.1. Determination of $\lambda_{\max}$ for the formed complexes.....	214
IV.A.2.1.2. Choice of solvent.....	214
IV.A.2.1.3. Reagent concentration.....	215
IV. A.2.1.4. Effect of phthalate-buffer (pH and volume) on the ion-pair complex formation.....	215
IV.A.2.1.5. Effect of time and temperature.....	216
IV.A.2.1.6. Stoichiometric relationship.....	216
IV.A.2.1.7.Suggested mechanism for ion-pair and charge-transfer	217

complex formation reaction.....	235
IV.A.2.2.Results of method validation.....	226

Section B

Spectrophotometric Method for Determination of Pioglitazone Hydrochloride, Using Ion-pair Technique

IV.B.1. Experimental.....	235
IV.B.1.1 Instrumentation.....	235
IV.B.1.2. Samples.....	235
IV.B.1.3. Chemicals and Reagents.....	235
IV.B.1.4. Pioglitazone hydrochloride standard solutions.....	235
IV.B.1.5. Procedures.....	235
IV.B.1.5.1. Spectral characteristics of pioglitazone ion-pair complex with bromophenol blue.....	235
IV.B.1.5.2. Method validation.....	236
IV.B.1.5.2.1. Linearity.....	236
IV.B.1.5.2.2. Accuracy.....	236
IV.B.1.5.2.3. Precision.....	237
IV.B.1.5.2.4. Robustness.....	237
IV.B.1.5.2.5. Ruggedness.....	237
IV.B.1.5.3. Application to the pharmaceutical preparation.....	237
IV.B.1.5.4. Validation by standard addition technique	238
IV.B.2. Results and discussion.....	238
IV.B.2.1. Method development.....	238
IV.B.2.1.1.Determination of $\lambda_{\max}$ for the formed ion-pair complex.....	239
IV. B.2.1.2. Choice of solvent.....	239
IV.B.2.1.3. Reagent concentration.....	239
IV. B.2.1.4. Effect of phthalate-buffer (pH and volume).....	239
IV.B.2.1.5. Effect of time and temperature.....	240
IV.B.2.1.6. Stoichiometric relationship.....	240
IV.B.2.1.7.Suggested mechanism for ion-pair and charge-transfer complex formation reaction.....	240
IV.B.2.2.Results of method validation.....	246

Section C

Spectrophotometric Methods for Determination of Rosiglitazone Maleate, Using Ion-pair and Charge-transfer Techniques

IV.C.1. Experimental.....	253
IV.C.1.1 Instrumentation.....	253

IV.C.1.2. Samples.....	253
IV.C.1.3. Chemicals and Reagents.....	253
IV.C.1.4. Rosiglitazone maleate standard solutions.....	253
IV.C.1.5. Procedures.....	254
IV.C.1.5.1. Spectral characteristics of rosiglitazone produced complexes.....	254
IV.C.1.5.1.a. Rosiglitazone ion-pair complex with bromophenol blue.....	254
IV.C.1.5.2.b. Rosiglitazone charge-transfer complex with chloranilic acid.....	254
IV.C.1.5.2. Method validation.....	255
IV.C.1.5.2.1. Linearity.....	255
IV.C.1.5.2.2. Accuracy.....	256
IV.C.1.5.2.3. Precision.....	256
IV.C.1.5.2.4. Robustness.....	256
IV.C.1.5.2.5. Ruggedness.....	257
IV.C.1.5.3. Application to the pharmaceutical preparation.....	257
IV.C.1.5.4. Validation by standard addition technique	257
IV.C.2. Results and Discussion.....	258
IV.C.2.1. Method Development.....	258
IV.C.2.1.1. Determination of $\lambda_{\max}$ for the formed complexes.....	258
IV.C.2.1.2. Choice of solvent.....	258
IV.C.2.1.3. Reagent concentration.....	259
IV. C.2.1.4. Effect of phthalate-buffer (pH and volume) on the ion-pair complex formation.....	259
IV.C.2.1.5. Effect of time and temperature.....	263
IV.C.2.1.6. Stoichiometric relationship.....	263
IV.C.2.1.7.Suggested mechanism for ion-pair and charge-transfer complex formation reaction.....	264
IV.C.2.2.Results of method validation.....	270

Part V General Discussion

V	General Discussion.....	279
---	-------------------------	-----

Part VI References

VI	References.....	283
----	-----------------	-----

Part VII Arabic Summary

LIST OF FIGURES

Figure		Page
[1]	The pathway for glucose transport and metabolism.	5
[2]	Common structural features found in RPG.	13
[3]	Common structural features found in thiazolidindiones PPAR- γ gonists; RGL and PGL	13
[4]	HPLC chromatogram of RPG (I); (40.00 $\mu\text{g.ml}^{-1}$).	48
[5]	HPLC chromatogram of RPG acid-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	49
[6]	HPLC chromatogram of RPG alkaline-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	49
[7]	HPLC chromatogram of RPG oxidative-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	49
[8]	Calibration curve correlating the peak area of the HPLC at 225.00 nm to the corresponding concentration of RPG (10.00 -50.00 $\mu\text{g.ml}^{-1}$).	50
[9]	HPLC chromatogram of mixture solution containing RPG (I); (40.00 $\mu\text{g.ml}^{-1}$) and its acid-degradates; (60.00 $\mu\text{g.ml}^{-1}$)	51
[10]	HPLC chromatogram of mixture solution containing RPG (I); (40.00 $\mu\text{g.ml}^{-1}$) and its alkaline-degradates; (60.00 $\mu\text{g.ml}^{-1}$)	51
[11]	HPLC chromatogram of mixture solution containing RPG (I); (40.00 $\mu\text{g.ml}^{-1}$) and its oxidative-degradates; (60.00 $\mu\text{g.ml}^{-1}$)	51
[12]	Mass spectrum of RPG	59
[13]	Mass spectrum of RPG acid-degradate 1	60
[14]	Mass spectrum of RPG acid-degradate 2	61
[15]	Mass spectrum of RPG alkaline-degradate	62
[16]	Mass spectrum of RPG oxidative-degradate	63
[17]	Chemical structure of RPG along with some possible hydrolysis products	64
[18]	HPLC chromatogram of PGL (I); (40.00 $\mu\text{g.ml}^{-1}$).	76
[19]	HPLC chromatogram of PGL acid-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	77

[20]	HPLC chromatogram of PGL alkaline-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	77
[21]	HPLC chromatogram of PGL oxidative-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	77
[22]	Calibration curve correlating the peak area of the HPLC at 220.00 nm to the corresponding concentration of PGL (10.00 -50.00 $\mu\text{g.ml}^{-1}$).	78
[23]	HPLC chromatogram of mixture solution containing PGL (I); (40.00 $\mu\text{g.ml}^{-1}$) and its acid-degradates; (60.00 $\mu\text{g.ml}^{-1}$)	79
[24]	HPLC chromatogram of mixture solution containing PGL (I); (40.00 $\mu\text{g.ml}^{-1}$) and its alkaline-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	79
[25]	HPLC chromatogram of mixture solution containing PGL (I); (40.00 $\mu\text{g.ml}^{-1}$) and its oxidative-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	79
[26]	Mass spectrum of PGL	87
[27]	Mass spectrum of PGL acid-degradate	88
[28]	Mass spectrum of PGL alkaline-degradate 1	89
[29]	Mass spectrum of PGL alkaline-degradate 2	90
[30]	Mass spectrum of PGL oxidative-degradate	91
[31]	HPLC chromatogram of RGL (I); (40.00 $\mu\text{g.ml}^{-1}$).	104
[32]	HPLC chromatogram of maleic acid; (20.00 $\mu\text{g.ml}^{-1}$).	104
[33]	HPLC chromatogram of RGL acid-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	105
[34]	HPLC chromatogram of RGL alkaline-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	105
[35]	HPLC chromatogram of RGL oxidative-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	105
[36]	Calibration curve correlating the peak area of the HPLC at 240.00 nm to the corresponding concentration of RGL (10.00 -60.00 $\mu\text{g.ml}^{-1}$).	106
[37]	HPLC chromatogram of mixture solution containing RGL (I); (40.00 $\mu\text{g.ml}^{-1}$) and its acid-degradates; (60.00 $\mu\text{g.ml}^{-1}$)	107
[38]	HPLC chromatogram of mixture solution containing RGL (I); (40.00 $\mu\text{g.ml}^{-1}$) and its alkaline-degradates; (60.00 $\mu\text{g.ml}^{-1}$).	107