

Phytochemical and Biological Studies of Some Artocarpus Species

Family Moraceae, Cultivated in EGYPT

Submitted in fulfillment of the requirements for the Master's degree in pharmaceutical sciences

(Pharmacognosy)

By

HaidySayedAbd El Atty

Quality control specialist

National Organization of Drug Control&Research

Supervised By

Prof. Dr. Abdel Nasser Badawi Singab

Professor of Pharmacognosy

Dean

Faculty of Pharmacy

Ain Shams University

Prof. Dr. Kamilia Fouly Taha

Professor of Pharmacognosy

Applied Research Center of Medicinal

Plants, National Organization of Drug

Control & Research

Pharmacognosy Department
Faculty of Pharmacy
Ain Shams University
(2012)

Acknowledgment

First of all, I thank Allah, the source of all knowledge, for helping me throughout this work. And gave me the power to continue.

My great appreciation and deepest gratitude are addressed to *prof. Abdel Nasser Badawi Singab*, Professor of pharmacognosy, Dean of faculty of pharmacy-Ain Shams University for suggesting the research point and supervision of this thesis, gave me valuable instructions, scientific guidance and continuous encouragement throughout the course of the practical work and helping me in biology work, NMR interpretation and writing the thesis.

My deepest thanks, and everlasting appreciation to *prof.Kamilia Fouly Taha*, Professor of pharmacognosy, Applied Research Center for Medicinal Plants in the National Organization for Drug Control and Research, for her supervision of this thesis, overwhelming support, valuable scientific opinions throughout the practical work and writing.

My deepest thanks to botanist specialist agricultural engineer\ Therese Labib Youssef in Orman garden herbarium for her help in bringing the plant samples and identified the plant

My everlasting appreciation and sincere indebtedness to **Assoc. prof/Hesham A.El-Beshbishy.** Assistant professor of Clinical Biochemistry and Molecular Biology, Faculty of Applied Medical Sciences, Taibah University, Al-Madinah Al-Munawarah, Saudi Arabia for carrying out the biological studies.

My deep gratitude to **Assoc. prof\ Ziad Moussa,**Chemistry departement Faculty of Science, Taibah University, Al-Madinah Al-Munawarah, Saudi Arabia for carrying out the NMR studies.

My deep gratitude to **prof \Sedky Bolous,** Professor of botany, Botany department, Desert research center for carrying out the botanical study

Appreciation is expressed to all my colleagues in the Department of Pharmacognosy, Faculty of Pharmacy, Ain Shams University, and my colleagues at the Applied Research Center for Medicinal Plants, NODCAR, specially *Dr. Gamal Ghazal*, *Dr. Gouda Talaat*, *Dr. Waleed Gamal El Din and Dr. ZeinabTalaat*.

Finally a special gratitude is acknowledged to my great father,my great mother, my sister, my husband and my friends for their persistent help, true sympathy, honest prayers, and encouragement throughout the course of thesis.

Haidy Sayed

Table of Contents

	Page
List of figures	i
List of tables	iv
List of abbreviations	vii
Introduction	1
Review of literature	4
A. Family Moraceae	4
B. Botanical description of family Moraceae	4
C.Botanical description of genus Artocarpus	5
D. Botanical description of <i>Artocarpus heterophyllus</i> Lam and <i>Artocarpus altilis</i> Fosberg	5
E. Ecology and distribution of <i>Artocarpus heterophyllus</i> Lam and <i>Artocarpus altilis</i> Fosberg	6
F. Folk medicinal uses	7
G. Economic uses	8
H. Biological activities of plants belonging to genus Artocarpus	9
I. Chemical constituents of the genus Artocarpus	14
Taxonomy	31
Material, methods and apparatus	34
I. Material	34
II .Methods	42
III. Apparatus	61

Chapter I: Botanical study	62
-Macromorphology of Artocarpus heterophyllus Lam	62
- Macromorphology of Artocarpus altilis Fosberg.	64
-Micromorphological study of the leaves of Artocarpus heterophyllus Lam	66
Chapter II: Biological studies	72
1. Determination of in vitro biological study of Artocarpus heterophyllus	
Lam and Artocarpus altilis Fosberg	72
1.1. Assay of antioxidant activity	72
2. determination of <i>in vivo</i> biological studies of the leaves of <i>Artocarpus</i>	78
heterophyllus Lam	
2.1. Determination of median lethal dose	78
2.2. Antihyperglycemic activity	78
2.3. Antihyperlipedimic activity	81
Chapter III: Phytochemical studies	86
1. Pharmacopeal constants of Artocarpus heterophyllus Lam and	86
Artocarpus altilis Fosberg	
2. Preliminary phytochemical investigation of the leaves of Artocarpus	87
heterophyllus Lam and Artocarpus altilis Fosberg.	
3.Quantitive determination of total polyphenols and flavonoids of	88
Artocarpus heterophyllus Lam.	
3.1. Determination of total polyphenols of 70% ethanolic extract of	88
A.heterophyllus Lam by colorimetric method calculated as gallic acid	

3.2. Determination of total flavonoids of 70% ethanolic extract of <i>A.heterophyllus</i> Lam by UV spectroscopy calculated as quercetin.	90
4.Determination of free and combined sugars by paper chromatography of defatted air dried leaves powder of <i>A.heterophyllus</i> Lam	92
4.1. Identification of free sugars	92
4.2. Identification of combined sugars	93
5.Phytochemical investigation of n-butanol fraction of the leaves of	94
Artocarpus heterophyllus Lam(jack fruit)	
Identification of compound $\underline{1}$	98
Identification of compound $\underline{2}$	102
Identification of compound $\underline{3}$	109
Identification of compound $\underline{4}$	117
Identification of compound $\underline{5}$	123
Chapter IV: Investigation of the lipoidal matter of dried leaves of A.	127
heterophyllus Lam	
1.1.Preparation of unsaponifiable fraction	127
1.2.Analysis of fatty acid content	128
Gas liquid chromatography analysis of fatty acid metyl ester	129
Gas chromatography/Mass spectroscopy of unsaponifiable matter	130
General summary	133
Conclusions and recommendations	142
References	144
Arabic summary	

List of figures

Figures	Page
1. a. Photographs of <i>Artocarpus heterophyllus</i> Lam tree in winter, leaves and fruits of <i>Artocarpus altilis</i> Fosberg, <i>Artocarpus</i>	32
heterophyllus Lam tree in summer.	
1.b.Photographs of leaves of <i>Artocarpus heterophyllus</i> Lam, mature leaf of <i>Artocarpus heterophyllus</i> , immature fruit on tree of <i>Artocarpus heterophyllus</i> Lam, mature small sized fruit of <i>Artocarpus heterophyllus</i> Lam.	33
2. Scheme for extraction of powdered air dried leaves of <i>Artocarpus heterophyllus</i> Lam.	44
3. Scheme for extraction of powdered air dried leaves of <i>Artocarpus atilis</i> Fosberg.	44
4. a. Photographs of jack fruit rind, male flowers, T.S. in fruit showing seeds of <i>A.heterophyllus</i> Lam.	63
4. b. Photographs of bread fruit rind, male flowers, T.S. in fruit showing seeds of <i>A. altilis</i> Fosberg.	65
5. Photograph of transverse section in midrib(x250).	68
6. Photograph of mesophyll (x250).	69
7. Photograph of epidermis, palisade and spongy tissues.	69
8. Photograph of wavy lower epidermis of young leaf (x250).	70
9. Photograph of upper epidermis had no stomata(x250).	70
10. Photograph of lower epidermis had anomocytic type stomata(x250)	71

i

11. DPPH free radical scavenging activity of different fractions	73
obtained from air dried powdered leaves of jack fruit (Artocarpus	
heterophyllus Lam).	
12. DPPH free radical scavenging activity of different fractions	73
obtained from air dried powdered leaves of bread fruit (Artocarpus	
altilis Fosberg).	
13. Comparison between DPPH free radical scavenging activity of	74
different fractions obtained from air dried powdered leaves of bread	
fruit (Artocarpus altilis Fosberg) and air dried powdered leaves of	
jack fruit (Artocrpus heterophyllus Lam).	
14. Ferrous chelating activity of different fractions obtained from air	75
dried powdered leaves of jack fruit (Artocarpus heterophyllus Lam).	
15. Standard calibration curve of gallic acid equivalent.	89
16. Standard calibration curve of quercetin.	91
17. Scheme for chromatographic fractionation of n-butanol extract	95
of the air dried powdered leaves of A.heterophyllus Lam.	
18. H-NMR (CD ₃ OD, 400 MHz) of compound 1:3,4, 5-trihydroxy-	100
benzoic acid (gallic acid).	
19. ¹³ C-NMR (CD ₃ OD, 100 MHz) of compound <u>1</u> :3,4, 5-trihydroxy-	101
benzoic acid (gallic acid).	101
	104
20. ¹ H-NMR spectrum of compound <u>2</u> (CD ₃ OD, 400 MHz):	106
quercetin-3- O - α -rhamnosyl (1 \rightarrow 6)- β -glucoside (rutin).	
21. 13 C-NMR spectrum of compound $\underline{2}$: quercetin-3- O - α -rhamnosyl	107
$(1 \rightarrow 6)$ - β -glucoside (rutin).	

22. HSQC-NMR spectrum of compound 2: quercetin-3-O-α-108 rhamnosyl (1 \rightarrow 6)- β -glucoside (rutin). 23. ¹H-NMR (CD₃OD, 400 MHz) of compound 3: quercetin -3-O-113 β -glucoside (isoquercitrin). 24. ¹³C-NMR (CD₃OD, 100 MHz) of compound <u>3</u>: quercetin -3-*O*-114 β -glucoside (isoquercitrin). 25. HSQC-NMR of compound 3: quercetin $-3-O-\beta$ -glucoside 115 (isoquercitrin). 26. ¹H-NMR spectrum of compound <u>4</u> (CD₃OD, 400 MHz):3, 4', 120 5,7-tetrahydroxyflavone (kaempferol). 27. ¹³C-NMR spectrum of compound 4 (CD₃OD, 100 MHz):3, 4', 5, 121 7-tetrahydroxyflavone (kaempferol). 28. HSQC-NMR spectrum of compound 4: 3, 4', 5, 7-122 tetrahydroxyflavone (kaempferol). 29. ¹H-NMR spectrum of compound 5: Apigenin 7-O- β-D-126 glucopyranoside. 30. GLC analysis of fatty acids methyl esters of the saponifiable 129

matter of air dried leaves of Artocarpus heterophyllus Lam.

List of Tables

Tables	Page
1. Geographic distribution, adaptation and propagation of both	6
Artocarpus heterophyllus Lam and Artocarpus altilis Fosberg.	
2. Different biological activities of different <i>Artocarpus</i> species.	9
3. Different phytochemical constituents isolated from different	14
species of genus Artocarpus.	
4. Results of extractives, yield, color and weight of leaves of	45
Artocarpus heterophyllus Lam.	
5. Results of extractives, yield, color and weight of leaves of	45
Artocarpus altilis Fosberg.	
6. Numerical values of the leaves of Artocarpus heterophyllus	67
Lam.	
7.Influence of oral intake of JFEE, JFBE (200mgKg ⁻¹ day ⁻¹) or	80
GLB (500 mg Kg ⁻¹ day ⁻¹) for 10 days to STZ-diabetic rats on	
serum protein content, fasting blood glucose FBG (mg%), lipid	
peroxides expressed as thiobarbituric acid reactive substance	
TBARS (nmol/ml), percentage of glycosylated hemoglobin A1C	
(%HbA1C) and insulin (U/ml).	
8.Influence of oral intake of JFEE, JFBE (200mgKg ⁻¹ day ⁻¹) or	82
GLB (500 mg Kg ⁻¹ day ⁻¹) for 10 days to STZ-diabetic rats on	
serum lipid profiles including total cholesterol TC (mg%), low	
density lipoprotein cholesterol LDL-C (mg%), high density	
lipoprotein cholesterol HDL-C (mg%), LDL/HDL ratio, very low	
density lipoprotein VLDL-C (mg%) and triglycerides (mg%).	

9. Pharmacopial constants of the leaves of <i>A. heterophyllus</i> Lam and <i>A. altilis</i> Fosberg.	86
10. Results of preliminary phytochemical screening for detection of different phytochemical constituents in both <i>Artocarpus heterophyllus</i> Lam and <i>Artocarpus altilis</i> Fosberg.	87
11. The absorbance measured for different concentrations of standard gallic acid solutions.	89
12. The absorbance measured for different concentrations of standard quercetin solutions.	90
13. Results of paper chromatography of free sugars of defatted air dried leaves powder of <i>A. heterophyllus</i> Lam.	92
14. Results of paper chromatography of hydrolysed combined sugars of defatted air dried leaves powder of <i>A.heterophyllus</i> Lam.	93
15. Results of column chromatographic analysis of F-2.	96
16. Physicochemical, chromatographic and spectral data of C1.	98
17. Physicochemical, chromatographic and spectral data of compound C2.	102
18. Physicochemical, chromatographic and spectral data of compound C3.	109
19. Physicochemical, chromatographic and spectral data of compound C4.	117
20. Physicochemical, chromatographic and spectral data of compound $C\underline{5}$.	123
21. GLC analysis of fatty acids methyl esters of the lipoidal matter of leaves of <i>Artocarpus heterophyllus</i> Lam.	129

22. GC\Ms of sterols and hydrocarbon of unsaponifiable fraction 131 of leaves of *Artocarpus heterophyllus* Lam.