



# **STUDIES ON THE DETECTION OF SOME INSECTICIDES IN ENTOMOLOGICAL SPECIMENS OF MEDICOLEGAL IMPORTANCE**

A Thesis

Submitted to The Department of Entomology, Faculty of Science,  
Ain-Shams University

In partial fulfillment of requirements for award of the Master  
Degree in Entomology

**By**

**Samah Mohammed Ahmed Hussein Ahmed**

B. Sc.

## **Supervisors**

**Prof. Dr. Reda Fadeel Ali Bakr**, Professor of Insect Toxicology,  
Entomology Department, Faculty of Science,  
Ain-Shams University

**Dr. Sanaa Abd El-Kader Mohamed El-Sawi**, Assistant  
Professor, Agricultural Research Center, Ministry of Agriculture

**Dr. Rawheia Hassan Ramadan**, Assistant Professor, Entomology  
Department, Faculty of Science, Benha University

**Entomology Department**

**Faculty of Science**

**Ain Shams University**

**2012**

## **Biography**

**Name:** Samah Mohammed Ahmed Hussien

**Date and place of birth:** 24 August 1984, Cairo.

**Date of Graduation:** 2005

**Degree awarded:** B.Sc. (Entomology/Chemistry), 2005,  
Faculty of Science, Ain Shams University.

**Date of registration for the M. Sc. Degree:** 11/6/2007

**Courses studied by the candidate in partial fulfillment of the requirements for Master Science Degree (M.Sc.) in Entomology:**

- 1) Population genetics and evolution.
- 2) Natural communities of insects.
- 3) Molecular Biology.
- 4) Insect Biochemistry.
- 5) Pollution of the environment.
- 6) Insects of Medical and Veterinary importance.
- 7) Insect Microbiology.
- 8) Parasitology and Immunology.
- 9) Insect Transmission of phytopathogens.
- 10) How to write and publish scientific reports.

**Statistical Course:** Bio-Statistics and Complete Science.

**Language Course:** English: M.Sc. Course.

## **Board of Supervision**

### **Prof. Dr. Reda Fadeel Ali Bakr**

Professor of Insect Toxicology  
Entomology Department  
Faculty of Science  
Ain-Shams University

### **Dr. Sanaa Abd El-Kader Mohamed El-Sawi**

Assistant Professor  
Agricultural Research Center  
Ministry of Agriculture

### **Dr. Rawheia Hassan Ramadan**

Assistant Professor  
Entomology Department  
Faculty of Science  
Benha University

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

جامعة عين شمس  
كلية العلوم

اسم الطالبة: سماح محمد أحمد حسين أحمد

عنوان الرسالة: دراسات علي كشف بعض المبيدات الحشرية في عينات  
حشرية ذات أهمية في الطب الشرعي

الدرجة العلمية: الماجستير

القسم التابع له: قسم علم الحشرات

الكلية التابع لها: كلية العلوم

الجامعة التابع لها: جامعة عين شمس

سنة التخرج: ٢٠٠٥

أَتَقَدِّمُ بِخَالِصِ آيَاتِ الشُّكْرِ وَالْعُرْفَانِ لِلَّهِ عَزَّ وَجَلَّ وَلِأَسَاتِذِي الْكَرَامِ:

أ.د. رضا فضيل علي بكر / أستاذ علم المبيدات – قسم علم الحشرات –  
كلية العلوم – جامعة عين شمس

د. سناء عبد القادر محمد الصاوي / أستاذ مساعد – المعمل المركزي  
لتحليل متبقيات المبيدات – مركز البحوث الزراعية

د. روية حسن رمضان / أستاذ مساعد – قسم علم الحشرات –  
كلية العلوم – جامعة بنها

جامعة عين شمس

كلية العلوم

رسالة ماجستير

اسم الطالبة: سماح محمد أحمد حسين أحمد

عنوان الرسالة: دراسات علي كشف بعض المبيدات الحشرية في عينات  
حشرية ذات أهمية في الطب الشرعي

الدرجة العلمية: الماجستير

لجنة الإشراف:

أستاذ علم المبيدات – كلية العلوم –  
جامعة عين شمس

أ. د. رضا فضيل علي بكر

أستاذ مساعد بالمعمل المركزي  
لتحليل متبقيات المبيدات – مركز  
البحوث الزراعية

د. سناء عبد القادر محمد الصاوي

أستاذ مساعد – كلية العلوم –  
جامعة بنها

د. روية حسن رمضان

لجنة الممتحنين:

١- أ. د. /

٢- أ. د. /

٣- أ. د. /

تاريخ البحث:

تاريخ المناقشة: / /

الدراسات العليا:

سنة المنح:

أجيزت الرسالة بتاريخ: / /

ختم الإجازة:



## LIST OF FIGURES

<b>No.</b>	<b>FIGURE NAME</b>	<b>Page No.</b>
<b>1</b>	Concentration of malathion (mg/kg) in tissues of rabbits administered different dosages of malathion. R0 = Control, R1 = 513 mg/kg, R2 = 1026 mg/kg	<b>46</b>
<b>2</b>	Correlation coefficient between administered dosage and malathion concentration in different tissues	<b>48</b>
<b>3</b>	Concentration of malathion (mg/kg) in pupae and larvae fed on (rabbits, liver tissues, muscle).	<b>50</b>
<b>4</b>	Correlation coefficient between administered dosage and malathion concentration in the third feeding instar larvae of <i>C. megacephala</i> (reared on control and treated rabbit carcasses, liver, muscle tissues) and pupae.	<b>52</b>
<b>5</b>	Mean body length (mm), over time (hrs), for <i>C. megacephala</i> larvae (from 2 <sup>nd</sup> instar to pupariation) reared on rabbit carcasses administered different dosages of malathion. R0 = control, R1 = 510 mg/kg, R2 = 1026 mg/kg	<b>65</b>
<b>6</b>	Mean body length (mm), over time (hrs), for <i>C. megacephala</i> larvae (from 2 <sup>nd</sup> instar to pupariation) fed on liver tissue from control and treated rabbit carcasses	<b>69</b>

<b>7</b>	Mean body length (mm), over time (hrs), for <i>C. megacephala</i> larvae (from 2 <sup>nd</sup> instar to pupariation) fed on muscle tissues from control and treated rabbit carcasses	<b>73</b>
<b>8</b>	Summary of body length over time for <i>C. megacephala</i> larvae reared on liver, muscle tissues and control & treated rabbit carcasses	<b>77</b>
<b>9</b>	Comparison of duration of the egg, larval and pupal stages to the total duration of <i>C. megacephala</i> reared on control and treated rabbit carcasses	<b>85</b>
<b>10</b>	Comparison of duration of the egg, larval and pupal stages to the total durations of <i>C. megacephala</i> reared on liver tissues from control and treated rabbit carcasses	<b>87</b>
<b>11</b>	Comparison of duration of the egg, larval and pupal stages to the total durations of <i>C. megacephala</i> reared on muscle tissues from control and treated rabbit carcasses	<b>89</b>
<b>12</b>	Summary of different mean durations of the developmental stages of <i>C. megacephala</i> at different temperature regimes (25.7, 28.4, 30.6 °C)	<b>101</b>
<b>13</b>	Electron micrograph of a dark columnar cell (dense compressed cells) in middle mid-gut showing nucleus (arrow). Magnification X 2,000	<b>106</b>
<b>14</b>	Electron micrograph of a light columnar cell in middle mid-gut (less electron dense cytoplasm) showing nucleus (Nu) surrounded by numerous vesicles of	<b>106</b>

	rough endoplasmic reticulum (rer-v) and chromatin patches of varying densities (head arrow) lysosomes (arrow). Magnification X 1,500	
<b>15</b>	Electron micrograph of a columnar cell in middle mid-gut showing nucleus (Nu) surrounded by numerous vesicles of rough endoplasmic reticulum (rer-v), lysosomes (arrow) and clumped (eu & hetero) chromatin (big arrow). Magnification X 1,000	<b>107</b>
<b>16</b>	Electron micrograph of a dark columnar cell in middle mid-gut showing microvilli (Mv), small dense secretory granules (arrow), small vesicles of rough endoplasmic reticulum (head arrows) and elongated & oval mitochondria (Mt). Magnification X 2,000	<b>107</b>
<b>17</b>	Electron micrograph of a columnar cell in middle midgut showing rarified mitochondria (Mt), rarified areas of cytoplasm (big white arrow), ribosomes (head arrow), rough endoplasmic membrane (REM) and Nuclear membrane (N.M). Magnification X 5,000	<b>108</b>
<b>18</b>	Electron micrograph of a columnar cell in middle mid-gut showing rarified mitochondria (Mt) and homogenously scattered lysosomes (Ly) between mitochondria. Magnification X 2,000	<b>108</b>
<b>19.a</b>	Electron micrograph of a dark columnar cell in middle midgut showing irregular nucleus and lacked clumping of chromatin. Magnification X 2,000	<b>109</b>
<b>19.b</b>	Electron micrograph of a dark columnar cell in middle midgut showing irregular nucleus. Magnification X 2,000	<b>109</b>

<b>20</b>	Electron micrograph of a columnar cell in middle midgut showing significant engulfed granules in curved mitochondria. Magnification X 4,000	<b>110</b>
<b>21</b>	Electron micrograph of a columnar cell in middle midgut showing accumulation of fat vacuoles at the periphery in between mitochondria and absence of lysosomes. Magnification X 1,200	<b>110</b>
<b>22</b>	Electron micrograph of a columnar cell in middle midgut showing no change in M.V (arrow) & fat vacuoles (F.v) in between mitochondria. Magnification X 4,000.	<b>111</b>
<b>23.a</b>	Electron micrograph of a columnar cell in middle midgut showing nucleus (Nu) abundant of fat vacuoles (small black arrows) and rarified areas of cytoplasm (white arrows) & coagulated cytoplasmic granules & organells (head arrow). Magnification X 2,000	<b>112</b>
<b>23.b</b>	Electron micrograph of a columnar cell in middle midgut showing abundant of fat vacuoles (black arrows). Magnification X 2,000	<b>112</b>
<b>24</b>	Electron micrograph of a columnar cell in middle midgut showing abundant of fat vacuoles (long white arrows) microvilli shrinkage (small black arrow). Magnification X 4,000	<b>113</b>
<b>25</b>	Electron micrograph of a columnar cell in middle midgut showing coagulated and shrinkage microvilli. Magnification X 2,000	<b>113</b>

## LIST OF TABLES

<b>No.</b>	<b>TABLE NAME</b>	<b>Page No.</b>
<b>1</b>	Recovery (%) of malathion from rabbit tissues and insects (larvae & pupae)	<b>40</b>
<b>2</b>	Concentration of malathion (mg/kg) in tissues of rabbits administered different dosages of malathion. R0 = Control, R1 = 513mg/kg, R2 = 1026 mg/kg	<b>45</b>
<b>3</b>	Correlation between administered dosage and malathion concentration in different tissues	<b>47</b>
<b>4</b>	Concentration of malathion (mg/kg) in pupae and larvae fed on (rabbits, liver tissues, muscles)	<b>49</b>
<b>5</b>	Correlation between administered dosage and malathion concentration in the third feeding instar larvae of <i>C. megacephala</i> (reared on control and treated rabbit carcasses, liver, muscle tissues) and pupae.	<b>51</b>
<b>6</b>	Correlation between concentration in (the third larval instars and pupae), (liver and muscles)	<b>53</b>
<b>7</b>	Correlation between concentration in (larvae and pupae) and Kidney	<b>54</b>
<b>8</b>	Correlation between concentration in (larvae and pupae) and Lung	<b>55</b>
<b>9</b>	Correlation between concentration in (larvae and pupae) and Fat	<b>56</b>

<b>10</b>	Correlation between concentration in (larvae and pupae) and Heart	<b>57</b>
<b>11</b>	Total body length of <i>C. megacephala</i> larvae (from 2 <sup>nd</sup> instar to Pupariation) reared on rabbit carcasses administered different dosages of malathion. R0 = Control, R1 = 513mg/kg, R2 = 1026 mg/kg	<b>64</b>
<b>12</b>	Significance table for rate of development of mean larval length of <i>C. megacephala</i> reared on rabbit carcasses administered different dosages of malathion. R0 = control, R1 = 510 mg/kg, R2 = 1026 mg/kg	<b>66</b>
<b>13</b>	The time of maximum larval length and duration of larval and pupal stages for colonies of <i>C. megacephala</i> reared on rabbit carcasses administered different dosages of malathion R0 = control, R1 = 510 mg/kg, R2 = 1026 mg/kg (mean $\pm$ SE)	<b>67</b>
<b>14</b>	Total body length of <i>C. megacephala</i> larvae (from 2 <sup>nd</sup> instar to pupariation) reared on liver tissues of rabbits administered different dosages of malathion	<b>68</b>
<b>15</b>	Significance table for rate of development of mean larval length of <i>C. megacephala</i> reared on liver tissues of rabbits treated with different dosages of malathion	<b>70</b>

<b>16</b>	The time of maximum larval length and duration of larval and pupal stages for colonies of <i>C. megacephala</i> reared on liver tissues of rabbits treated with malathion (mean $\pm$ SE)	<b>71</b>
<b>17</b>	Total body length of <i>C.megacephala</i> larvae (from 2 <sup>nd</sup> instar to Pupariation) reared on muscle tissues of rabbits administered different dosages of malathion	<b>72</b>
<b>18</b>	Significance table for rate of development of mean larval length of <i>C. megacephala</i> reared on muscle tissues of rabbits treated with different dosages of malathion	<b>74</b>
<b>19</b>	The time of maximum larval length and duration of larval and pupal stages for colonies of <i>C. megacephala</i> reared on muscle tissues of rabbits treated with malathion (mean $\pm$ SE)	<b>75</b>
<b>20</b>	ANOVA results for the differences in larval length and development rate between all diets	<b>76</b>
<b>21</b>	Summary of best fit of body length over time for larvae of <i>C. megacephala</i> reared on rabbit carcasses treated with malathion	<b>79</b>
<b>22</b>	Summary of best fit of body length over time for larvae of <i>C.megacephala</i> reared on liver from rabbits carcasses treated with malathion	<b>80</b>