



PRODUCTION AND EFFICIENCY TESTING OF INORGANIC NANOPHOTOCATALYSTS FOR WATER TREATMENT

Thesis submitted

By

Mona Karam Farhan

B.Sc. (Chemistry) 2007

To

CHEMISTRY DEPARTMENT

FACULTY OF SCIENCE

AIN-SHAMS UNIVERSITY

For

THE DEGREE

M.Sc. IN CHEMISTRY

(2011)

Thesis Advisors

Prof. Dr. M. S. A. Abdel-Mottaleb

Professor of Inorganic and Photochemistry

Faculty of Science, Ain Shams University

Prof. Dr. Mohamed Abo- Aly

Professor of Inorganic and Photochemistry

Faculty of Science, Ain Shams University

Dr. Essam Bakir

Lecturer of inorganic and Photochemistry

Faculty of Science, Ain Shams University

Approval Sheet

Name of candidate: Mona Karam Farhan

Degree: M.Sc. Degree in Chemistry

**Thesis Title: PRODUCTION AND EFFICIENCY TESTING OF
INORGANIC NANOCATALYST FOR WATER TREATMENT**

This Thesis has **been approved by:**

1- Prof. Dr. M. S. A. Abdel-Mottaleb

2- Prof. Dr. Mohamed Abo- Aly

3- Dr. Essam Bakir

Approval

Chairman of Chemistry Department

Prof. Dr. Maged Shafik Antonious Nakhla

PRODUCTION AND EFFICIENCY TESTING OF INORGANIC
NANOCATALYST FOR WATER TREATMENT

By

Mona Karam Farhan
B.Sc. (Chemistry) 2007
Under the supervision of:

1- Prof. Dr. M. S. A. Abdel-Mottaleb

Chemistry Department, Faculty of Science, Ain Shams University, Abbasia, Cairo,
Egypt

2- Prof. Dr. M. Abu-Ali

Chemistry Department, Faculty of Science, Ain Shams University, Abbasia, Cairo,
Egypt

3- Dr. Essam Bakir

Chemistry Department, Faculty of Science, Ain Shams University, Abbasia, Cairo,
Egypt



كلية العلوم
قسم الكيمياء

إنتاج واختبار كفاءة حفازات ضوئية نانوية غير عضوية لمعالجة المياه

رسالة مقدمة من

منى كرم فرحان

بكالوريوس العلوم (كيمياء)

إلى

كلية العلوم – جامعة عين شمس

للحصول على درجة الماجستير في العلوم (كيمياء)

تحت إشراف

أ.د. محمد صبري عبد المطلب

أستاذ الكيمياء كلية العلوم جامعة عين شمس

أ.د. محمد محمود أبو علي

أستاذ الكيمياء غير العضوية

د. عصام محمد عدلي مصطفى

مدرس الكيمياء غير العضوية



كلية العلوم
قسم الكيمياء

رسالة الماجستير

اسم الطالب: منى كرم فرحان.
عنوان الرسالة: إنتاج واختبار كفاءة حفازات ضوئية نانوية غير عضوية لمعالجة المياه

اسم الدرجة: ماجستير في العلوم .

لجنة الاشراف :

- ١ أ.د. محمد صبري عبد المطلب
أستاذ الكيمياء غير العضوية والضوئية
- ٢ أ.د. محمد محمود أبو علي: أستاذ الكيمياء غير العضوية
- ٣ عصام محمد عدلي مصطفى بكير: مدرس كيمياء غير العضوية والضوئية.

لجنة تحكيم الرسالة:

١. أ.د. محمد صبري عبد المطلب
أستاذ الكيمياء غير العضوية والضوئية
- ٢ أ.د. مجدي حسني السيد :
أستاذ الكيمياء المتفرغ مركز بحوث الصحراء
- ٢ بدر عواد:
٣ أستاذ الكيمياء كلية العلوم جامعة الأزهر.

Umm Al-Qura

ختم الإجازة

أجيزت الرسالة

موافقة مجلس الكلية

بتاريخ / /
موافقة مجلس الجامعة

/

/ /



جامعة عين شمس
كلية العلوم

اسم الطالب: منى كرم فرحان رجب

تاريخ الميلاد: ١٩٨٦/٧/١٣

اسم الدرجة: ماجستير في العلوم

القسم التابع له: (الكيمياء غير العضوية – الكيمياء الضوئية)

سنة التخرج: ٢٠٠٧

سنة المنح:



جامعة عين شمس
كلية العلوم

شكر

أتقدم بأسمى آيات الشكر والعرفان لله عز وجل ولأساتذتي الأفاضل الذين قاموا بالاشراف
على هذا العمل وهم:
أ.د. محمد صبري عبد المطلب.
أ.د. محمد محمود أبو علي.
د. عصام محمد عدلي

كذلك أشكر أعضاء الفريق العلمي للكيمياء الضوئية والكيمياء الشمسية

CONTENTS

	Aim of the work	1
	Introduction	4
1	Nanotechnology	4
1.1.1	What is nanotechnology	4
1.1.2	Nanocatalysts	5
1.1.3	Nanotechnology and the Environment	6
1.2	Water Pollution	7
1.3	Treatment of Textile Wastewater	9
1.4	Advanced Oxidation Process	12
1.4.1	Homogeneous Photocatalysis (AOPs)	15
1.4.1.1	<i>The O₃/UV Process</i>	15
1.4.1.2	<i>The H₂O₂/UV Process</i>	16
1.4.1.3	<i>UV Fenton 's Processes</i>	17
1.4.2	Heterogeneous Photocatalysis	19
1.4.2.1	Photocatalytic Process	21
1.4.2.2.	Photocatalytic semiconductor	22
1.4.3	Titanium dioxide	25
1.4.3.1	Historical background of TiO ₂ .	25
1.4.3.2	Titanium dioxide structure	28
1.4.3.3	Titanium Degussa	29
1.4.3.4	Titanium dioxide Photocatalysis	30
1.4.3.5	Enhancement of TiO ₂ Photocatalyst	34
1.4.3.5.1	Composite Semiconductors	37
1.4.3.6	Practical application of TiO ₂ photocatalyst	40
1.5	Literature Review	43
	CHAPTER 2 Experimental Techniques and Methods	
2.1	Materials	50
2.2	Instruments	52
2.2.1	UV-VIS Measurements	52
2.2.2	XRD Measurements	52
2.2.3	TEM Measurements	52

2.2.4	Photoillumination Setup	53
2.2.5	Centrifuge	53
2.2.6	pH meter	53
2.3	Preparations and General Procedures	54
2.3.1	Photodegradation of Dyes	54
2.3.2	Decolorization Analysis	54
2.4	Methods of Data Analysis	55
2.4.1	The kinetic Rate Laws	55
2.4.2	Adsorption and Photodegradation Kinetics:	57
2.4.5	XRD analysis	60
2.5	Data Analysis and Handling	61
2.5.1	Statistical and Data Handling Package:	61

CHAPTER 3 Results and Discussion

3.1	Characterization	62
3.1.1	Crystal properties and morphology	62
3.1.1.1	XRD	62
3.1.1.2	TEM	68
3.1.2	Optical properties	72
3.2	Photocatalytic Efficiency test using Remazol Red RB-133 Dye as a model compound.	73
3.2.1	Absorption spectra of Remazol Red RB-133 Dye	73
3.2.2	Photostability of Remazol Red Dye	73
3.3	Studies of different factors affecting the photodegradation of RRRB-133 dye using TiO₂	75
3.3.1	Effect of Operating parameters: effect of different concentrations of Remazol Red Dye RRRB-133	75
3.3.2	Effect of Operating parameters: different TiO ₂ concentration	81
3.3.3.	Influence of pH on photocatalytic degradation of dyes in waste water	87

3.4	Enhancement of Photocatalytic activity of TiO₂	95
3.4.1	Influence of catalyst composition process	101
3.4.1.1	<i>Irradiation Under UV-visible light</i>	101
3.4.1.2	<i>Irradiation under UV</i>	107
3.4.1.3	<i>Irradiation under visible</i>	115
3.4.2	Influence of Oxidizing agents addition	115
	Summary and conclusions	124
	References	131

List of Figures

1.1	Schematic representation of the “band gap model.”	21
1.2	Energies for various semiconductors in aqueous electrolytes at pH = 1.	21
1.3	Bulk structure of rutile and anatase	28
1.4	Schematic diagram of photocatalytic process initiated by photon acting on the semiconductor.	31
1.5	Steps in heterogeneous photocatalytic reaction.	33
1.6	Excitation steps using dye molecule sensitizer	36
1.7	Photoexcitation in composite semiconductor Photocatalyst	37
1.8	Major areas of activity in titanium dioxide photocatalysis	40
1.9	Thin-film-fixed-bed reactor	41
1.10	Super-hydrophilicity	42
2.1	Schematic diagram of Remazol Red RRB-133	50
3.1	XRD pattern of a,b,c,d,e where [a: pure TiO ₂ nanocrystal, b: TiO ₂ - CdS nanocomposite, c: TiO ₂ - ZnO nanocomposite, d: TiO ₂ - SnS nanocomposite, e: TiO ₂ - Fe ₃ O ₄ nanocomposite, f: TiO ₂ - ZnS nanocomposite	63
3.2	XRD pattern of a,b,c,d,e where [a: pure TiO ₂ nanocrystal, and TiO ₂ - CdS nanocomposite, b: pure TiO ₂ nanocrystal, and pure ZnS nanocatalyst and TiO ₂ - ZnS nanocomposite, c: pure TiO ₂ nanocrystal and pure Fe ₃ O ₄ nanocatalyst and TiO ₂ - Fe ₃ O ₄ nanocomposite, d: pure TiO ₂ nanocrystal, and TiO ₂ - ZnO nanocomposite, e: pure TiO ₂ nanocatalyst and TiO ₂ - SnS nanocomposite	66
3.3	TEM image of a,b,c,d,e where [a: pure TiO ₂ nanocrystal, b: CdS nanocatalyst, c: TiO ₂ - CdS nanocomposite, d: ZnS nanocatalyst, Fe ₃ O ₄ CdS nanocatalyst	69
3.4	Normalized absorbance spectrum of TiO ₂ degauss and TiO ₂ doped with 2% wt of nano CdS. (solid line TiO ₂ and dash line CdS-TiO ₂).	72
3.5	The absorption spectrum of RRRB-133 in aqueous solution	74

3.6	Absorption spectrum of RRRB-133 (1×10^{-5} mole.l ⁻¹) under irradiation of UV-VIS light	74
3.7	Change in absorbance of 1g.l ⁻¹ loaded TiO ₂ on photodegradation of aqueous solution of RRRB-133 dye, a: (1×10^{-5} mole.l ⁻¹) b: (3×10^{-5} mole.l ⁻¹) at different time with wavelength ($\lambda_{\text{analytical}} = 524\text{nm}$)	79
3.8	First order kinetic plots of loaded TiO ₂ on the degradation of RRRB-133 dye ($C_0 = 1 \times 10^{-5}$ mole.l ⁻¹), aqueous medium, ($\lambda_{\text{analytical}} = 524\text{nm}$)	80
3.9	Representation of decolorization of RRRB-133 at different initial concentration [TiO ₂]=1g.l ⁻¹ in aqueous solution, ($\lambda_{\text{analytical}} = 524\text{nm}$)	81
3.10	Change in absorbance of (a:0.2g.l ⁻¹ , b: 0.6g.l ⁻¹ : c: 1g.l ⁻¹) loaded TiO ₂ on photodegradation of aqueous solution of (1×10^{-5} mole.l ⁻¹) RRRB-133 dye at different time with wavelength, ($\lambda_{\text{analytical}} = 524\text{nm}$)	82
3.11	First order kinetic plots of loaded TiO ₂ on the degradation of RRRB-133 dye ($C_0 = 1 \times 10^{-5}$ mole.l ⁻¹), aqueous medium. ($\lambda_{\text{analytical}} = 524\text{nm}$)	84
3.12	Plot of (k) versus [TiO ₂] for RRRB-133 dye(using 1×10^{-5} mole.l ⁻¹) [dye] and Degussa TiO ₂), ($\lambda_{\text{analytical}} = 524\text{nm}$)	85
3.13	Representation of decolorization of RRRB-133 at different concentrations [TiO ₂] = [0.2g.l ⁻¹ , 0.6g.l ⁻¹ , 1g.l ⁻¹] in aqueous solution, ($\lambda_{\text{analytical}} = 524\text{nm}$)	85
3.14	First order kinetic plots of loaded TiO ₂ on the degradation of RRRB-133 dye at different pH values ($C_0 = 1 \times 10^{-5}$ mole.l ⁻¹), TiO ₂ , ($\lambda_{\text{analytical}} = 524\text{nm}$)	93
3.15	Plot of (k) versus value of pH for RRRB-133 dye(using 1×10^{-5} mole.l ⁻¹ [dye] and Degussa TiO ₂), ($\lambda_{\text{analytical}} = 524\text{nm}$)	94
3.16	Representation of decolorization of RRRB-133 loaded with [TiO ₂] = [1g.l ⁻¹] at different pH in aqueous solution, ($\lambda_{\text{analytical}} = 524\text{nm}$)	94
3.17	(a) Energy diagram illustrating the coupling of two SC in which vectorial electron transfer occurs from the light-activated SC to the nonactivated TiO ₂ . (b) Diagram valence and conductance band position for various semiconductors, and useful, relevant redox couples depicting the coupling of SC in which vectorial movement of electron and holes is possible	98

3.18	Valence and Conductance band position for various semiconductors, and useful, relevant redox couples.	99
3.19	Change in absorbance of (a: TiO_2 (1g.l^{-1}) doped with CdS (0.2g.l^{-1}) b: TiO_2 (1g.l^{-1}) alone c: TiO_2 (1g.l^{-1}) doped with ZnO (0.2g.l^{-1})) on photodegradation of aqueous solution of RRRB-133 dye (3×10^{-5} mole.l ⁻¹), ($\lambda_{\text{analytical}} = 524\text{nm}$) at different time, under UV – visible radiation,	104
3.20	Initial rate First order kinetic plots of nano-composite TiO_2 catalyst on the degradation of RRRB-133 dye ($C_0 = 1 \times 10^{-5}$ mole.l ⁻¹), under irradiation of UV	105
3.21	Representation of initial kinetic rate constants of decolorization of RRRB-133 dye ($C_0 = 3 \times 10^{-5}$ mole.l ⁻¹), ($\lambda_{\text{analytical}} = 524\text{nm}$) using different nano-composite TiO_2 catalyst under UV- Visible light irradiation (16 min)	106
3.22	Change in absorbance of (a: TiO_2 (1g.l^{-1}) doped with CdS (0.2g.l^{-1}) b: TiO_2 (1g.l^{-1}) alone c: TiO_2 (1g.l^{-1}) doped with ZnO (0.2g.l^{-1})) on photodegradation of aqueous solution of RRRB-133 dye (3×10^{-5} mole.l ⁻¹), ($\lambda_{\text{analytical}} = 524\text{nm}$) at different time, under UV radiation,	108
3.23	Initial rate First order kinetic plots of nano-composite TiO_2 catalyst on the degradation of RRRB-133 dye ($C_0 = 1 \times 10^{-5}$ mole.l ⁻¹), under UV irradiation light ($\lambda_{\text{analytical}} = 524\text{nm}$) (2.5min)	109
3.24	Representation of initial kinetic rate constants of decolorization of RRRB-133 dye ($C_0 = 3 \times 10^{-5}$ mole.l ⁻¹), ($\lambda_{\text{analytical}} = 524\text{nm}$) using nano-composite TiO_2 catalyst under UV irradiation at time (2.5min)	110
3.25	Representation of percentage of degradation of RRRB-133 dye ($C_0 = 3 \times 10^{-5}$ mole.l ⁻¹), ($\lambda_{\text{analytical}} = 524\text{nm}$) using different nano-composites TiO_2 catalyst under irradiation of UV at time (2.5 min)	111
3.26	Change in absorbance of (a: TiO_2 (1g.l^{-1}) alone, b: TiO_2 (1g.l^{-1}) doped with CdS (0.2g.l^{-1}) c: TiO_2 (1g.l^{-1}) doped with ZnO (0.2g.l^{-1})) on photodegradation of aqueous solution of RRRB-133 dye (3×10^{-5} mole.l ⁻¹), ($\lambda_{\text{analytical}} = 524\text{nm}$) at different time, under Visible irradiation.	112

- 3.27 Initial rate First order kinetic plots of nano-composites TiO₂ catalyst on the degradation of RRRB-133 dye ($C_0 = 1 \times 10^{-5}$ mole.l⁻¹), under Visible light, ($\lambda_{\text{analytical}} = 524\text{nm}$) 114
- 3.28 Representation of initial kinetic rate constants of decolorization of RRRB-133 dye ($C_0 = 3 \times 10^{-5}$ mole.l⁻¹), using different nano-composites TiO₂ catalyst under visible irradiation (2.5min), ($\lambda_{\text{analytical}} = 524\text{nm}$) 115
- 3.29 First order kinetic plots of decolorization of RRRB-133 with nano-composites TiO₂-SnS catalyst at different concentrations of H₂O₂, under irradiation of UV- visible light. [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, SnS (0.02g.l⁻¹) H₂O₂ (2mM, 4mM, 6mM) ($\lambda_{\text{analytical}} = 524\text{nm}$) 118
- 3.30 First order kinetic plots of decolorization of RRRB-133 with nano-composite TiO₂-ZnS catalyst at different concentration of H₂O₂, under UV- visible light irradiation [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, ZnS(0.02g.l⁻¹) H₂O₂ (2mM, 4mM, 6mM), ($\lambda_{\text{analytical}} = 524\text{nm}$) 119
- 3.31 Effect of hydrogen peroxide concentrations on photocatalytic rate constant of RRRB-133 Dye= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, SnS (0.02g.l⁻¹), H₂O₂ (2mM, 2mM, 4mM, 6mM) ($\lambda_{\text{analytical}} = 524\text{nm}$) 119
- 3.32 Effect of hydrogen peroxide concentrations on photocatalytic rate constant of RRRB-133 [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, ZnS(0.02g.l⁻¹), H₂O₂ (2mM, 4mM, 6mM), ($\lambda_{\text{analytical}} = 524\text{nm}$) 120
- 3.33 Effect of hydrogen peroxide concentrations on photocatalytic rate constant of RRRB-133 [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, CdS(0.02g.l⁻¹), H₂O₂ (0mM, 4mM, 6mM), ($\lambda_{\text{analytical}} = 524\text{nm}$) 120
- 3.34 Kinetic rate constant of RRRB-133 with nano-composite TiO₂-SnS catalyst at different concentrations of H₂O₂, under irradiation of UV- visible light. RRRB-133 [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, SnS(0.02g.l⁻¹), H₂O₂ (2mM, 4mM, 6mM) ($\lambda_{\text{analytical}} = 524\text{nm}$) 121
- 3.35 Represent the kinetic rate constant of RRRB-133 with nano-composites TiO₂-ZnS catalyst at different concentration of H₂O₂, under irradiation of UV-visible light. RRRB-133 [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, ZnS(0.02g.l⁻¹), H₂O₂ (2mM, 4mM, 6mM), ($\lambda_{\text{analytical}} = 524\text{nm}$) 121

- 3.36 The kinetic rate constant of RRRB-133 with nano-composites TiO₂-CdS catalyst at different concentrations of H₂O₂, under irradiation of UV- visible light. RRRB-133 [Dye](3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, CdS(0.02g.l⁻¹), H₂O₂ (0mM, 4mM , 6mM) ($\lambda_{\text{analytical}} = 524\text{nm}$) 122
- 3.37 First order initial kinetic plots of decolorization of RRRB-133 with nano-composites TiO₂-SnS, TiO₂-ZnS, TiO₂-CdS, H₂O₂, under irradiation of UV- visible light. [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]= 1g.l⁻¹, SnS(0.02g.l⁻¹), ZnS (0.02g.l⁻¹), CdS(0.02g.l⁻¹) ,H₂O₂ (4mM) ($\lambda_{\text{analytical}} = 524\text{nm}$) 122
- 3.38 First order initial kinetic plots of decolorization of RRRB-133 with nano-composites TiO₂-SnS, TiO₂-ZnS, TiO₂-CdS, H₂O₂, under irradiation of UV- Visible light. [Dye]= (3×10^{-5} mole.l⁻¹), [TiO₂]=1g.l⁻¹, SnS(0.02g.l⁻¹), ZnS (0.02g.l⁻¹), CdS(0.02g.l⁻¹) , (4mM)H₂O₂ ($\lambda_{\text{analytical}} = 524\text{nm}$) 123