

Utility of isatin in synthesis of heterocycles of anticipated biological activity

A Thesis Submitted for the Degree of M. Sc

In

Organic Chemistry

Presented

By

Rabaa Mohamed Kamel Mohamed

Chemistry Department Faculty of Science Ain Shams University

Supervised

By

Prof. Dr. Ahmed Said Ahmed YoussefProf. of Organic Chemistry

Dr.Samir Abd Elaziz Emara Dr.Magdy Mohamed Hemdan

Ass. Prof. of Organic Chemistry

Ass. Prof. of Organic Chemistry



Utility of isatin in synthesis of heterocycles of anticipated biological activity

Thesis Advisors

Thesis Approved

Prof. Dr. Ahmed said Ahmed Youssef
Prof. of Organic Chemistry

Dr. Samir Abd Elaziz EmaraAss. Prof. of Organic Chemistry

Dr. Magdy Mohamed Hemdan Ass.Prof. of Organic Chemistry

Head of Chemistry Department Prof. Dr. Hamed Ahmed Derbala

•••••

Approval of Chemistry Department Council //2014
Approval of Faculty Council //2014
Approval of University Council //2014

DEDICATION

To my distinguished parents:

I do appreciate my God for giving me
such wonderfully parents for their continuous
support, encouragement, and enlighting my
life.

ACKNOWLEDGEMENT

The author wishes to refer her deep appreciation and gratitude

To

Prof. Dr. Ahmed Said Ahmed Youssef, Professor of Organic Chemistry, Chemistry Department, Faculty of Science, Ain Shams University.

Dr. Samir Abd Elaziz Emara, Ass. Professor of Organic Chemistry, Chemistry Department, faculty of Science, Ain Shams University.

J.

Dr. Magdy Mohamed Hemdan, Ass. Professor of Organic Chemistry, Chemistry Department, Faculty of Science, Ain Shams University.

For

Suggesting the subject, interpreting the results, their valuable scientific guidance, help, and encouragement during the work of this thesis.

QUALIFICATION

Name: Rabaa Mohamed Kamel Mohamed

Scientific Degree: B. Sc.

Department: Micro- Chemistry

Faculty: Faculty of Science

University: Ain Shams University

B. Sc. Graduation Year: 2009

M. Sc. Graduation Year: 2014

Contents

SUMMARY	1
INTRODUCTION	1
Synthesis of Isatins	
The syndmeyer methodology	5
Use of nitroacetanilides	
The stole procedure	
The martinet isatin synthesis	8
Metalation of anilide derivatives	9
Reactivity of isatin and its derivatives towords electronic	<u>rophiles</u>
N-alkylation	11
N-methylene amino derivatives	16
N-acylation and N-sulfonylation	18
Application of isatins in organic synthesis	21
Reduction of the heterocyclic ring	24
Oxidation of the heterocyclic ring	28
Nucleophilic attack at positions C-2 or C-3	30
Miscellaneous reactions	42
Conclusions	48
Aim of the work	49
DISCUSSION & RESULTS	
Reaction of isatin and its derivatives with different	
nucleophiles	
Reaction of isatins with hydrazine hydrate & its derivat	ives51
Reaction of 1b with hydroxylamine hydrochloride	
Reaction of compounds 1a,b with o-phenylene diamine	
Reactions of isatin with hydrazides	
Reactions of isatin with carbanions	
Benzoylation and acylation of compounds 1a,b	
Benzoylation of compound 1a	87
Acetylation of compounds 1a,b	
,	

Reactions of compounds 15a,b with hydrazines	.101
<u>Utilities of hydrazones 2a,7 in the synthesis of some</u>	
<u>heterocycles</u>	
Action of carbon nucleophile on hydrazone 2a	113
Reaction of ethyl acetoacetate with hydrazone 7	123
Reaction of hydrazone 2a with benzoyl/lauroyl	
isothiocyanate	127
Action of acetic anhydride on hydrazone 7	133
EXPERIMENTAL	143
REFERENCES	162
ARABIC SUMMARY	1

List of tables

Table	Title	Page
No.		
1	IR spectral data for compounds 2a-d	53
2	¹ H-NMR spectral data for compounds 2a-d	54
3	Ions & relative intensities of 2a	56
4	Ions & relative intensities of 2b	58
5	Ions & relative intensities of 2c	60
6	Ions & relative intensities of 2d	62
7	IR spectral data for compound 3	66
8	¹ H-NMR spectral data for compound 3	67
9	Ions & relative intensities of 3	68
10	IR spectral data for compounds 4,5	72
11	¹ H-NMR spectral data for compounds 4,5	73
12	Ions & relative intensities of 4	74
13	IR spectral data for compounds 7,9	79
14	¹ H-NMR spectral data for compounds 7,9	80
15	Ions & relative intensities of 7	81
16	IR spectral data for compounds 11,12	83
17	¹ H-NMR spectral data for compounds 11,12	84
18	Ions & relative intensities of 11	85
19	Ions & relative intensities of 12	86
20	IR spectral data for compounds 13,14	88
21	¹ H-NMR spectral data for compounds 13,14	89
22	Ions & relative intensities of 13	90
23	Ions & relative intensities of 14	91
24	IR spectral data for compounds 15a,b&16	95
25	¹ H-NMR spectral data for compounds 15a,b&16	96
26	Ions & relative intensities of 15a	97
27	Ions & relative intensities of 16	99

28	IR spectral data for compounds 18,19	104
29	¹ H-NMR spectral data for compounds 18,19	105
30	Ions & relative intensities of 18	105
31	IR spectral data for compound 20	100
32		110
33	H-NMR spectral data for compound 20 Ions & relative intensities of 20	111
34	IR spectral data for compounds 23a,b	114
35	¹ H-NMR spectral data for compounds 23a,b	115
36	Ions & relative intensities of 23a	116
37	Ions & relative intensities of 23b	118
38	IR spectral data for compounds 25-27	121
39	¹ H-NMR spectral data for compounds 25-27	122
40	IR spectral data for compound 28	124
41	¹ H-NMR spectral data for compound 28	125
42	IR spectral data for compounds 30a,b&31	128
43	¹ H-NMR spectral data for compounds 30a,b&31	130
44	Ions & relative intensities of 30a	131
45	Ions & relative intensities of 31	132
46	IR spectral data for compounds 32,33	136
47	¹ H-NMR spectral data for compound 32	136
48	¹ H-NMR spectral data for compound 33	137
49	Ions & relative intensities of 32	138
50	Ions & relative intensities of 33	139
51	IR spectral data for compound 34	141
52	¹ H-NMR spectral data for compound 34	142

Summary

Isatin and some of its derivatives are used in this investigation on the hope of getting heterocyclic compounds bearing indolinone moiety to enhance their biological activities. Thus, 2 - (2- oxindolin -3 - ylidene) malononitrile (1a) or (E,Z)ethyl 2-cyano-2-(2-oxindolin-3-vlidene)acetate (**1b**) or isatin-β-thiosemicarbazone (**1c**) undergo reactions with prototype hydrazine hydrate itself and some of its simple congeners and gave hydrazone indoline-2-one derivatives bearing moiety (2a-d). of (E,Z) ethyl 2-cyano-2-(2-oxindolin-3-Refluxing ylidene)acetate (1b) with hydroxylamine hydrochloride in ethanol and a catalytic amount of ammonium acetate produced the oxime derivative 3. Treatment of 1a, b with o-phenylenediamine in acetic acid yielded 5H-indolo[2,3blguinoxaline(4) in one-pot reaction. On the other hand the reaction of **1a** with o-phenylenediamine in ethanol produced di-hydroquinoxaline derivative 5 in good yield. The reaction of hydrazide derivative 6 and isatin 1 in ethanol afforded the hydrazone derivative 7 in a good vield. Refluxing of isatin1 and imidazolone derivative 8 in n-butanol produced compound **9** in a good yield. The reaction of isatin **1** with 2-benzamidoacetic acid in acetic anhydride not afforded the expected oxazolone derivative **10**, instead it produced the open adduct **11**. On the other hand the reaction of **1** with p-nitrobenzyl cyanide in the presence of a catalytic amount of triethyl amine gave a mixture of diastereo isomers of **12**.

We aimed to undergo benzovlation or acylation to compounds 1a and 1b hoping to prevent the removal of dicyano methylene of orcyanocarboethoxy methylene moieties of compounds 1a 1b by the action of hydrazines. Heating of **1a** with benzoyl chloride in dry acetone in the presence of anhydrous potassium carbonate as a base yielded an adduct which was proved to be a mixture of two compounds, the 2-(1-benzoyl-2-oxindolin-3-ylidene) malononitrile **(13)** and ethyl 2-(2benzamidophenyl)-3,3-dicyanoacrylate (14). Compound 14 is formed by ring opening of 13 during its crystallization with ethanol. Heating of compounds 1a and 1b in acetic anhydride furnished compounds 15a,b as a mixture of N-acetyl and O-acetyl derivatives in addition

to ethyl 2-amino-3-cyanoquinoline-4-carboxylate (16) in case of the reaction with 1a. Treating an ethanolic solution of 15a with hydrazine hydrate yielded the cinnoline derivative 18 in addition to a minor amount of the hydrazone derivative 2a. Similar treatment of compound 15b with hydrazine hydrate gave the N-acetyl hydrazone 19 and small amounts of the hydrazone derivative 2a. The reactions of 15a and 15b with phenyl hydrazine furnished (Z)-1-acetyl-3-(2-phenyl hydrazono) indolin-2-one (20), beside compound 2b in minor amount.

The authors utilized hydrazones 2 and 7 in the synthesis of some heterocycles. Thus, refluxing the derivative with hydrazone 2a acetyl acetone, ethyl acetoacetate, diethyl malonate or malononitrile in pyridine afforded the spiro derivatives 23a, 23b, 25 and 27, in addition to the fused system 26 in case of its reaction with diethyl malonate. Treating of hydrazone7 with ethyl acetoacetate in dioxane with a catalytic amount of pipredine; gave the dihydropyrrole derivative 28. Refluxing of hydrazone 2a with benzovl isothiocyanate and/or lauroylisothiocyanate in dry acetonitrile furnished the thiosemicarbazide derivatives **30a,b** in good yields. Compound **30a** was heated with hydrazine hydrate in the presence of a catalytic amount of pipredine, to give the 1,2,4-triazole derivative **31.**Compound **7**was heated in acetic anhydride on the hope of getting a heterocyclic system bearing indoline moiety. Instead, it furnished a mixture of oxazolone derivative **32** and hydrazone derivative **33**.

Refluxing of compound **2d** in acetic anhydride gave the N-acetyl derivative **34**.

The chemistry of isatins

1. Introduction

Isatin (1H-indole-2,3-dione) (1) is a synthetically versatile substrate, where it can be used for the synthesis of a large variety of heterocyclic compounds, such as indoles, quinolines, and as a raw material for drug synthesis. Isatin has also been found in mammalian tissues, and its function as a modulator of biochemical processes has been the subject of several discussions. The advances in the use of isatin for organic synthesis are reported in this thesis.

Isatin (1H-indole-2,3-dione) was first obtained by Erdman and Laurent in 1841 as a product from the oxidation of indigo by nitric and chromic acids.

In nature, isatin is found in plants of the genus *Isatis*, in *Calanthe discolor* LINDL.² and in

CouroupitaguianensisAubl,³ and has also been found as a component of the secretion from the parotid gland of *Bufo*frogs,⁴ and in humans as it is a metabolic derivative of adrenaline.⁵⁻⁷ Substituted isatins are also found in plants, for example the melosatin alkaloids (methoxy phenyl pentyl isatins) obtained from the Caribbean tumorigenic plant *Melochiatomentosa*⁸⁻¹⁰ as well as from fungi: 6-(3'-methylbuten-2'-yl)isatin was isolated from *Streptomyces albus* ¹¹ and 5-(3'-methylbuten-2'-yl)isatin from *Chaetomium globosum*. ¹²Isatin has also been found to be a component of coal tar. ¹³

Recently, 6-methoxy-1-methylisatin was isolated from the plant *Boronellakoniambiensislies*. ¹⁴ Isatin-3-oximes are useful antiallergic agents with potent and prolonged action. Derivatives of these compounds are also potassium channel openers for treatment of respiratory diseases, convulsions, renal disorders, urinary incontinence and diarrhea. ^{15,16}

It has been reported that several compounds containing an isatin moiety possess antibacterial, antifungal, anticonvulsant, anti-inflammatory, and anti-