

# **Ultrasound in Pediatric Anesthesia**

*An Essay*

Submitted for partial fulfillment of Master Degree in  
Anesthesia

*By*

**Sherif Mahrous Sayed Ali**  
(M.B.B.CH.)

*Under Supervision Of*

**Prof. Dr. Seif Al Islam Abd Elaziz**

Professor of Anesthesia, Intensive Care and Pain Management  
Faculty of Medicine Ain Shams University

**Prof. Dr. Sherif Farouk Ibrahim**

Professor of Anesthesia, Intensive Care and Pain Management  
Faculty of Medicine Ain Shams University

**Dr. Hanaa Mohamed Abd Allah ElGendy**

Lecturer of Anesthesia, Intensive Care and Pain Management  
Faculty of Medicine Ain Shams University

*Faculty of medicine  
Ain Shams University  
2014*



## *Acknowledgement*

*First of all, all gratitude is due to **God** almighty for blessing this work, until it has reached its end, as a part of his generous help, throughout my life.*

*Really I can hardly find the words to express my gratitude to **Prof. Dr. Seif Al Islam Abd Elaziz** Professor of Anesthesia, Intensive Care and Pain Management, Faculty of Medicine, Ain Shams University, for his supervision, continuous help, encouragement throughout this work and tremendous effort he has done in the meticulous revision of the whole work. It is a great honor to work under her guidance and supervision.*

*I am greatly honored to express my sincere appreciation and indebted to **Prof. Dr. Sherif Farouk Ibrahim**, Professor of Anesthesia, Intensive Care and Pain Management, Faculty of Medicine, Ain Shams University, for his guidance, valuable advice, continuous encouragement, moral support, constructive direction and sincere supervision of this work,*

*I would like also to express my sincere appreciation and gratitude to **Dr. Hanaa Mohamed Abd Allah El Gendy** Lecturer of Anesthesia, Intensive Care and Pain Management, Faculty of Medicine, Ain Shams University, for her continuous directions supportive guidance, valuable facilities, unlimited time and effort offered to me and support throughout the whole work,*

*Last but not least, I dedicate this work to my family, Mother, Father (God mercy on him), my wife and my kids.*



---

## Contents

	<i>Page</i>
List of Figures	I
List of Tables	VII
List of Abbreviations	VIII
Introduction	1
Chapter 1 : Anatomical and Physiological Differences between Adults and Pediatrics.	3
Chapter 2 : Physical Principles of Ultrasound.	31
Chapter 3 : Safety and Efficacy of Ultrasound in Pediatric Anesthesia.	60
♦ Safety and Efficacy of Ultrasound Guided Regional Anesthesia in Pediatrics.	62
♦ Safety and Efficacy of Ultrasound Guided Vascular Access in Pediatrics.	98
Summary	130
References	132
Arabic Summary	150

---

## List of Figures

Figure N0.	Title	Page N0.
Figure (1)	A, adult larynx, B, infant larynx, A, anterior; P, posterior.	4
Figure (2)	The percentage of type I muscle fibers composition of the diaphragm and intercostal muscles in children and adults.	6
Figure (3)	Comparison of Glomerular Filtration Rate in different ages of children and adults.	13
Figure (4)	Changes in total serum protein and albumin values with maturation.	17
Figure (5)	Distance from the skin to the epidural and subarachnoid spaces at different intervertebral levels and through the sacral hiatus.	23
Figure (6)	Ultrasound imaging of the caudal part of the spine in a 3-week-old infant. A: Long-axis view. B: Short-axis view (transverse echotomography).	24
Figure (7)	Position of the distal end of the spinal cord in relation to the vertebral column and meninges according to the age of development. A: 24 weeks B: newborn C: 8 years old child D: adult.	25
Figure (8)	Ultrasound imaging of the inguinal area to locate the femoral nerve at different ages. The femoral nerve is clearly visible in the 16-year-old adolescent but difficult to identify in the 3-year-old child.	28
Figure (9)	Specular and scattered echo waves.	36
Figure (10)	Doppler ultrasound. A: Color Doppler. B: Power Doppler. C: Pulsed Doppler.	38
Figure (11)	Transverse sonogram of the forearm demonstrating reduced gain (A), excess gain (B), and optimal gain (C).	40
Figure (12)	Schematic diagram illustrating the different types of ultrasound transducers. Note how the ultrasound beam is emitted from each of these transducers.	42
Figure (13)	Axis of scan. A: Transverse scan. B: Longitudinal scan. SCM: sternocleidomastoid muscle, THY: thyroid, CA: carotid artery, IJV: internal jugular vein.	46
Figure (14)	Schematic diagram to demonstrate the relative echogenicity of various tissues	47
Figure (15)	The plane of ultrasound imaging. Note that the ultrasound beam is only 1 mm thick and for a needle to be visible during an ultrasound-guided intervention it must lie within this plane of imaging.	48
Figure (16)	Axis of intervention. Out-of-plane (A) and in-plane (B) techniques	48

Figure (17)	Anisotropy. Note how a small change in angle of the ultrasound beam from the neutral position (A) has affected the visibility of the median nerve in the forearm (white arrow, B).	50
Figure (18)	Ultrasound appearance of peripheral nerves. A: Transverse sonogram of the sciatic nerve in the thigh. B: Transverse sonogram of the brachial plexus in the interscalene groove. C: Transverse sonogram of the median nerve in the forearm. D: Longitudinal sonogram of the sciatic nerve in the thigh.	51
Figure (19)	Longitudinal sonogram of the intercostal space demonstrating the ultrasound appearances of bone and pleura.	52
Figure (20)	Tissue harmonic imaging (THI). Sagittal sonogram of the infraclavicular fossa. A: Conventional scan. B: Conventional scan with THI.	53
Figure (21)	Compound imaging. Transverse sonogram of the axilla. A: Conventional scan. B: Conventional scan with compound imaging.	54
Figure (22)	Panoramic imaging. Transverse panoramic scan of the forearm. FDS: flexor digitorum superficialis; FCU: flexor carpi ulnaris; FPL: flexor pollicis longus; FDP: flexor digitorum profundus.	55
Figure (23)	Hypoechoic nerve structures. C5–7 nerve roots at the level of the posterior interscalene groove. Yellow arrow indicates the C5 root.	65
Figure (24)	Hyperechoic nerve structure. Musculocutaneous nerve between the short head of the biceps and the coracobrachialis muscles.	65
Figure (25)	Ultrasonographic visualization of neuraxial structures at the thoracic level in a 1000 g neonate. White arrow: posterior dura mater; gray arrow: flavum ligament; oblique green arrow: epidural catheter.	66
Figure (26)	Sonographic appearance of the brachial plexus (white arrows) in the supraclavicular fossa. SCA: subclavian artery; SA: scalenus anterior; SM: scalenus medius.	69
Figure (27)	Sonoanatomy at the supraclavicular block location using a linear hockey stick probe placed in the coronal oblique plane.	70
Figure (28)	Sagittal sonogram of the left infraclavicular fossa. AA: axillary artery; AV: axillary vein.	71
Figure (29)	Sagittal sonogram of the infraclavicular fossa after local anesthetic injection “doughnut sign.” AA: axillary artery; LA: local anesthetic; Pec: pectoralis.	72
Figure (30)	Schematic diagram showing the positions of the cords of the brachial plexus in infraclavicular block and the sites at which the local anesthetic is injected: (1) posterior cord, (2) lateral cord, and (3) medial cord.	73

Figure (31)	Sonoanatomy at the interscalene groove using a linear hockey stick probe.	75
Figure (32)	Sonoanatomy of the axillary block location using a linear hockey stick probe placed transversely to the humerus.	76
Figure (33)	Transverse sonogram of the left axilla. AA: axillary artery; AV: axillary vein; CRB: coracobrachialis muscle; MN: median nerve; RN: radial nerve; MCN: musculocutaneous nerve.	77
Figure (34)	Schematic diagram showing the positions of the three main nerves (median, radial, and ulnar) of the brachial plexus in relation to the axillary artery, as identified on transverse ultrasonography, at the axilla.	77
Figure (35)	Transverse sonogram of the cubital fossa. BA: brachial artery; BCR: brachioradialis.	78
Figure (36)	Transverse sonogram of the forearm. FDP: flexor digitorum profundus; FDS: flexor digitorum superficialis; RA: radial artery; UA: ulnar artery.	78
Figure (37)	Transverse sonogram of the arm, just above the elbow, showing the radial nerve. BCR: brachioradialis; ECRL: extensor carpi radialis longus.	79
Figure (38)	Sonoanatomy at the lumbar plexus block location using a linear probe placed transversely at L4/ 5.	80
Figure (39)	Longitudinal sonogram of the lumbar paravertebral region showing the lumbar plexus. TP: transverse process; ESM: erector spinae muscle.	80
Figure (40)	Transverse sonogram of the inguinal region showing the femoral nerve and its relations. FA: femoral artery; FV: femoral vein.	83
Figure (41)	Transverse section through the gluteal region at the level of the quadratus femoris muscle showing the subgluteal space and its contents.	84
Figure (42)	Sonoanatomy at the subgluteal location using a linear probe. Shows sciatic nerve lies at the midpoint between ischial tuberosity and greater trochanter of femur.	84
Figure (43)	Sonoanatomy in the popliteal fossa just distal to bifurcation of the sciatic nerve using a linear hockey stick probe.	85
Figure (44)	Transverse sonogram between the greater trochanter and the ischial tuberosity. GM: gluteus maximus; QF: quadratus femoris.	85
Figure (45)	Transverse sonogram of the sciatic nerve at the subgluteal space after distribution of local anesthetic injection around the sciatic nerve. GM: gluteus maximus; LA: local anesthetic; QF: quadratus femoris.	85
Figure (46)	Transverse sonogram of the sciatic nerve at the apex of the popliteal fossa. AM: adductor magnus; BF: biceps femoris.	87



Figure (47)	Transverse sonogram of the thigh showing the adductor canal. AL: adductor longus; FA: femoral artery; FV: femoral vein; VM: vastus medialis.	88
Figure (48)	Rectus sheath sonography.	89
Figure (49)	Transverse sonogram of the inguinal region. ASIS: anterior superior iliac spine; IO: internal oblique; TA: transverse abdominis.	91
Figure (50)	Ilioinguinal iliohypogastric nerve block. The in-plane needle insertion technique. ASIS: anterior superior iliac spine; IO: internal oblique; TA: transverse abdominis.	92
Figure (51)	Longitudinal paramedian sonogram of the thoracic spine in a neonate. CSF: cerebrospinal fluid.	92
Figure (52)	Transverse sonogram of the thoracic spine in a neonate. CSF: cerebrospinal fluid.	93
Figure (53)	Longitudinal paramedian sonogram of the thoracolumbar spine in a neonate showing the termination of the spinal cord. CSF: cerebrospinal fluid.	93
Figure (54)	Sonoanatomy of the spinal column and canal in a transverse axis at the L3/ L4 level using a linear probe.	95
Figure (55)	Sonoanatomy of the spinal column and canal in a paramedian longitudinal view at the midthoracic level using a linear probe.	95
Figure (56)	Transverse sonogram of the sacrum.	97
Figure (57)	Probe position for short-axis ultrasound picture of the internal jugular vein (IJV) and common carotid artery (CCA). <b>B:</b> Short-axis ultrasound picture of the IJV and the CCA.	101
Figure (58)	<b>A:</b> The probe position for long-axis ultrasound picture of the IJV. <b>B:</b> Long-axis ultrasound picture of the IJV.	102
Figure (59)	Short-axis view of the IJV and the CCA with IJV compressed by probe pressure.	102
Figure (60)	Short-axis view of the IJV and the CC with the IJV distended with a Valsalva maneuver.	102
Figure (61)	Short-axis view of the IJV (blue) and the CCA (red) with color Doppler.	103
Figure (62)	<b>A:</b> The ultrasound probe is positioned for a short-axis view of the IJV and the needle is out of plane. <b>B:</b> In the ultrasound picture of the short-axis view of the IJV, the needle appears as a small dot within the vein.	103
Figure (63)	<b>A:</b> The probe is positioned for a long-axis view of the IJV and the needle is in plane. <b>B:</b> In the ultrasound picture of the long-axis view of IJV, the needle shaft can be seen and the tip is entering the vein.	105
Figure (64)	Subclavian ultrasound-guided approach one: probe above and introducer below the clavicle in a 2.2-kg neonate. Cl: clavicle; SN: sternal notch.	106

Figure (65)	Subclavian ultrasound-guided approach one: probe above and introducer below the clavicle in a 9-y-old child. Cl: clavicle; SN: sternal notch.	107
Figure (66)	Needle entry point: Subclavian ultrasound-guided approach one: 3-y-old child: probe above and introducer below the clavicle.	107
Figure (67)	Subclavian ultrasound-guided approach one: ultrasound view. CA: carotid artery; CL: clavicle; FR: first rib; IJV: internal jugular vein; SCV: subclavian vein.	107
Figure (68)	Subclavian ultrasound-guided approach one: ultrasound view with the needle in the lumen of the subclavian vein (SCV). CA: carotid artery; CL: clavicle; FR: first rib; IJV: internal jugular vein.	108
Figure (69)	Subclavian ultrasound-guided approach two: probe above and introducer above the clavicle. Needle entry point.	108
Figure (70)	Subclavian ultrasound-guided approach two ultrasound view: probe above and introducer above the clavicle.	109
Figure (71)	Subclavian ultrasound-guided approach two ultrasound view: needle in situ.	109
Figure (72)	Subclavian ultrasound-guided approach two: skin exit site for nontunneled catheter.	109
Figure (73)	Subclavian ultrasound-guided approach two: skin exit site for tunneled catheter. Note that it is easier to dress and maintain with a small child.	110
Figure (74)	Probe position for ultrasound-guided lateral subclavian-axillary vein access. C: coracoid process.	112
Figure (75)	Needle entry point for ultrasound-guided lateral subclavian-axillary vein.	112
Figure (76)	Ultrasound image: ultrasound-guided lateral subclavian-axillary vein access.	112
Figure (77)	Catheter skin exit site, ultrasound-guided lateral subclavian-axillary vein access. Note that the site is much easier to dress to prevent dislodgment than neck positions in the small child.	113
Figure (78)	Ultrasound-guided lateral subclavian-axillary vein access: approach two: in-plane. Note that this is difficult because of the smaller surface area to fit the probe in the smaller child or infant.	113
Figure (79)	child with probe and needle in position for a short-axis view of the IJV with out of-plane needle approach.	117
Figure (80)	Ultrasound short-axis view of the neck. CA: carotid artery; IJV: internal jugular vein.	117
Figure (81)	A child with the ultrasound probe in position for a long-axis view of the IJV with the needle entering in an out-of-plane approach.	118

Figure (82)	Ultrasound-guided approach to the femoral vein in a child with ultrasound probe position and needle position for femoral vein cannulation.	118
Figure (83)	Ultrasound view of the femoral region.	119
Figure (84)	Ultrasound picture of arm veins: basilic vein (BAv), brachial vein (BRv), and brachial artery (BRa). The basilic vein is situated alone, whereas the brachial veins are often paired on either side of the artery.	123
Figure (85)	The same patient as in Figure 85 with color Doppler to delineate the artery. On the ultrasound, the artery walls are pulsatile.	123
Figure (86)	Short-axis ultrasound view of the basilic vein. With color Doppler and a distal arm squeeze, the vein fills briefly with a flash of color.	123
Figure (87)	Cannula insertion technique. Short-axis out-of-plane approach. It is sometimes easier with this view to have a second person holding the probe.	124
Figure (88)	Ultrasound view short-axis with the needle entering the vein in an out-of-plane technique. The needle tip is first seen compressing the vein wall, then appears as a dot in the vein.	125
Figure (89)	Long-axis out-of-plane ultrasound-guided approach. Place the probe to obtain a short-axis view and centralize it over the vein.	125
Figure (90)	Rotate the probe to the long-axis view.	126
Figure (91)	The ultrasound image of the long-axis view.	126
Figure (92)	Needle entry in the long-axis view. The needle is seen compressing the anterior vein wall.	126
Figure (93)	Find and mark technique for radial artery cannulation.	128
Figure (94)	Real time 2-dimensional ultrasound radial artery cannulation.	128

## List of Tables

<b>Table No.</b>	<b>Title</b>	<b>Page N0.</b>
<b>Table (1)</b>	The major differences in static lung volumes and other variables between infants and adults.	8
<b>Table (2)</b>	Normal range of resting heart rate and blood pressure in children.	11
<b>Table (3)</b>	Main Anatomic and Physiologic Factors in the Pediatric Period That Can Influence the Selection or Performance of a Regional Block Procedure. AAG: $\alpha_1$ -acid glycoprotein; Cmax: peak plasma concentration; HAS: human serum albumin; Tmax: time to reach Cmax.	28-30
<b>Table (4)</b>	Propagation velocity of sound in body tissues.	33
<b>Table (5)</b>	Scanning Routine.	58
<b>Table (6)</b>	Results of neuraxial imaging according to age groups.	66
<b>Table (7)</b>	Common peripheral and neuraxial nerve blocks in children, IP: in-plane; OOP: out-of plane.	66-67
<b>Table (8)</b>	Comparison of Dynamic Ultrasound, Static Ultrasound, and Landmark Approaches to the Internal Jugular vein Cannulation From Randomized, Controlled Trials. CA: carotid artery; DUS: dynamic ultrasound; LM: landmark; SUS: static ultrasound.	115

## List of Abbreviations

2,3-DPG	2,3-Diphosphoglycerate
2D	Two-dimensional
3D	Three-dimensional
A	Articular process
AA	Axillary artery
AAG	$\alpha_1$ -acid glycoprotein
ABL	Allowable blood loss
AL	Adductor longus
AM	Adductor magnus
ASAS	Anterior spinal artery syndrome
ASIS	Anterior superior iliac spine
AV	Axillary vein
BA	Brachial artery
BAv	Basilic vein
BBB	Blood Brain Barrier
BCR	Brachioradialis
BF	Biceps femoris
BRa	Brachial artery
BRv	Brachial vein
C	Corocoid process
C5	Cervical vertebra number five
CA	Carotid artery
CBF	Cerebral blood flow
CC	Closing capacity
CCA	Common carotid artery
CDF	Color doppler flow
CL	Clavicle
cm	Centimeter
Cmax	Peak plasma concentration
CRB	Coracobrachialis muscle
CSF	Cerebrospinal fluid
CT	Computerized tomography
CVCs	Central venous catheters
dB	Decibel
DGC	Depth-gain compensation
DUS	Dynamic ultrasound

EBV	Estimate of Blood Volume
ECF	Extracellular fluid
ECG	Electrocardiogram
ECRL	Extensor carpi radialis longus
ESM	Erector spinae muscle
f	Frequency
FA	Femoral artery
FCU	Flexor carpi ulnaris
FDP	Flexor digitorum profundus
FDS	Flexor digitorum superficialis
FPL	Flexor pollicis longus
FR	First rib
FRC	Functional residual capacity
FV	Femoral vein
Gen	General
GFR	Glomerular Filtration Rate
GM	Gluteus maximus
HbF	Fetal hemoglobin
HR	Heart rate
hr	Hour
HSA	Human serum albumin
Hz	Hertz
IJV	Internal jugular vein
IO	Internal oblique
IP	In-plane
IV	Intravenous
IVC	Inferior vena cava
Kcal	Kilocalory
Kcl	Potassium chloride
kg	Kilogram
kHz	Kilohertz
L	Liter
L1	Lumber vertebra number one
LA	Local anesthetic
LM	Landmark
MCN	Musculocutaneous nerve
MHz	Megahertz
min	Minute
mL	Milliliter
mm	Millimeter

mm Hg	Millimeters of mercury
MN	Median nerve
mol	Mole
mos	Months
N	Number
NaCl	Sodium chloride
NICE	The National Institute for Clinical Excellence
OOP	Out-of plane
PA	Popliteal artery
Pec	Pectoralis
Pen	Penetration
PICCS	Peripherally inserted central catheters
PRAN	The Pediatric Regional Anesthesia Network
PZT	Lead Zirconium Titanate
QF	Quadratus femoris
RA	Radial artery
RN	Radial nerve
S	Spinous process
S1	Sacral vertebra number one
SA	Scalenus anterior
SCA	Subclavian artery
SCM	Sternocleidomastoid muscle
SCV	Subclavian vein
sec	Second
SM	Scalenus medius
SN	Sternal notch
SONAR	Sound Navigation and Ranging
SUS	Static ultrasound
SVC	Superior vena cava
T	Thoracic vertebra
TA	Transverse abdominis
TAP	Transverses abdominis plane
TGC	Time gain compensation
THI	Tissue harmonic imaging
THY	Thyroid
TLC	Total lung capacity
Tmax	Time to reach peak plasma concentration
TP	Transverse process
UA	Ulnar artery
US	Ultrasound