



Role of MR Enterography in the assessment of inflammatory bowel diseases

A Thesis

Submitted to the Faculty of Medicine, Ain Shams University
In Partial Fulfilment of the Requirements for the MD Degree in
Radiology

By

Maha Ahmed Sayed Abdel Karim

M.B.,B.Ch., M.Sc.

Faculty of Medicine, Ain Shams University

Under supervision of

Prof. Dr. Aida Mohamed El Shibiny

Professor of Radiology

Faculty of Medicine - Ain Shams University

Prof. Dr. Mohammad Amin Sakr

Professor of Tropical Medicine

Faculty of Medicine - Ain Shams University

Prof. Dr. Mohamed Shaker Ghazy

Assistant Professor of Radiology

Faculty of Medicine - Ain Shams University

Dr. Rasha Salah El Din Hussein

Lecturer of Radiology

Faculty of Medicine - Ain Shams University

Faculty of Medicine

Ain Shams University

2018



Acknowledgement

*First, I wish to express my deep thanks, sincere gratitude to **ALLAH**, who always helps me, care for me and grunted me the ability to accomplish this thesis.*

*I would like to express my deepest gratitude, thanks and gratefulness to **Prof. Dr. Aida Mohamed El Shibiny**, Professor of Radiology, Faculty of Medicine, Ain Shams University, for her enthusiastic support, continuous encouragement valuable scientific advices, and great help through out of the accomplishment of this work,*

*I am very grateful to **Prof. Dr. Mohammad Amin Sakr**, Professor of Tropical Medicine, Faculty of Medicine, Ain Shams University, for his kind supervision, support, indispensable suggestion, and great help through out of course of my thesis.*

*My sincere thanks to **Prof. Dr. Mohamed Shaker Ghazy**, Assistant Professor of Radiology, Faculty of Medicine, Ain Shams University, for his kind and meticulous supervision, support, help, valuable supervision all through the work,*

*Many thanks, and sincere gratitude to **Dr. Rasha Salah El Din Hussein**, Lecturer of Radiology, Faculty of Medicine, Ain Shams University for her kind supervision encouragement and meticulous revision of this work,*

Words can never express my sincere thanks to my family for their generous support and continuous encouragement.

Maha Abdel Karim

LIST OF CONTENTS

List of Abbreviations.....	II
List of Tables	III
List of Figures	V
Introduction and Aim of the work.....	1
Review of literature.....	5
Anatomy of the small bowel.....	5
Pathophysiology of IBD	18
Manifestations of IBD	32
Imaging of IBD.....	42
Technique of MR Enterography.....	68
Patients and Methods.....	80
Results.....	89
Illustrative Cases	110
Discussion.....	156
Summary & Conclusions	170
References	173
Arabic Summary	-

LIST OF ABBREVIATIONS

IBD	Inflammatory Bowel Diseases.
CD	Crohn's Disease
UC	Ulcerative Colitis.
MRE	Magnetic Resonance Enterography.
CTE	Computed Tomography Enterography.
DWI	Diffusion Weighted Image
ADC	Apparent Diffusion Coefficient.
HASTE	Half-Fourier Acquisition Single-Shot Turbo Spin Echo.
SSFSE	Single Shot Fast Spin Echo.
SSFP	Steady State Free Precession Sequence.
CDAI	Crohn's Disease Activity Index.
CDEIS	Crohn's Disease Endoscopic Score of Severity.
CDAS	Crohn's Disease Activity Score.
AIS	Acute Inflammation Score.

LIST OF TABLES

Table no.	Title	Page no.
1	Differences between jejunum and ileum.....	8
2	Macroscopic features used for the diagnosis of IBD	27
3	Microscopic features used for the diagnosis of IBD.....	28
4	Truelove and Witts' severity index for ulcerative colitis	36
5	Manifestations of CD & UC.....	37
6	The CDAI.....	39
7	MRE characteristics of the four subtypes of Crohn's disease	50
8	Characteristics of the study population	89
9	Distribution of bowel wall thickening.....	91
10	Enhancement patterns and status of lymphadenopathy	92
11	Demographic data of IBD cases.....	94
12	Distribution of small and large bowel affection in CD and UC	96
13	Maximum wall thickness range and its mean in all IBD cases.....	97
14	Difference in mean of maximum wall thickness between active and non active IBD cases	97

Table no.	Title	Page no.
15	Mean ADC value in diseased and normal bowel loops	98
16	Cut –off ADC value in Crohn’s disease & Ulcerative colitis	98
17	Percentage of high T2 signal, enhancement, lymphadenopathy and engorged vasa recta in IBD cases.....	99
18	Percentage of clinical activity in IBD cases.....	100
19	Percentage of MR activity in IBD cases	101
20	Correlation between MRE results and clinical results in activity assessment of CD	102
21	Correlation between MRE results and clinical results in activity assessment of UC	103
22	Diagnostic accuracy of MR activity in prediction of clinical activity results.....	104
23 & 24	Correlation between DWI restriction and disease activity	105
25	Correlation between lymphadenopathy, engorged vasa recta and MR disease activity	107
26	Percentage of complications in IBD	108
27	Complications of CD and their percentage	109

LIST OF FIGURES

Fig. no.	Title	Page no.
1	Small and large intestine in situ	5
2	Relations of duodenum.....	6
3	Relations of duodenum.....	9
4	Greater omentum and mesentery of small intestine	11
5	Coronal MR enterography coronal image of the jejunum.....	15
6	Coronal MR enterography image of the normal distal ileum with fewer folds visualized in comparison to that of the jejunum and duodenum.....	15
7	Coronal T2WI showing adequate bowel distention and excellent wall conspicuity of both jejunal and ileal loops.	16
8	Coronal T1WI post contrast with fat saturation obtained during the portal phase	16
9	Multiple small mesenteric lymph nodes and vascular branches exhibit low signal intensity on true FISP images.....	17
10	Small intestine: Layers of wall of small intestine	18
11	Histology of the small intestine.....	19
12	An example of restricted diffusion within the wall of an acutely inflamed distal ileum in a patient with acute Crohn's disease	51
13	Wall thickening and mesenteric changes	51
14	Active inflammatory CD in a 15-year-old girl with intermittent right lower quadrant pain of increasing severity and elevated inflammatory biomarker levels ..	52
15	Axial post contrast T1 weighted images of a 34-year-old man with acute Crohn's disease.....	53
16	Coronal T1WI post contrast	53

Fig. no.	Title	Page no.
17	(A) Coronal T2WI, (B) Axial T2WI both showing mesenteric hypertrophy	54
18	Axial contrast enhanced T1 weighted image demonstrate extensive inflammation and enhancement of small bowel and mesentery in the pelvis in a patient with acute Crohn's disease	54
19	(A) Coronal T2WI, (B) Coronal T1WI post contrast with fat saturation showing star sign.....	55
20	(A) True fast imaging with steady-state precession, (B) pre contrast and (C) post contrast T1 weighted images of a 43-year-old patient with chronic Crohn's disease.....	55
21	Mucosal healing on serial MRE in a patient with Crohn's disease affecting the transverse colon	56
22	Magnetic resonance enterography in a 24-year-old male with ulcerative colitis	57
23	Ultrasound of the pelvis in a 33 years old female with right iliac fossa pain	61
24	Selected intravenous and oral contrast enhanced computed tomography images of three different patients with acute Crohn's disease	63
25	Coronal oral and intravenous contrast enhanced computed tomography showing comb sign.....	64
26	Axial CT contrast enhanced image of a 29-year-old patient with fistulating Crohn's disease	64
27	CT of 32-year-old female patient with ulcerative colitis showing pneumoperitoneum	65
28	Axial contrast enhanced computed tomography in a patient with chronic Crohn's disease	65
29	Coronal intravenous contrast enhanced CT showing lead pipe appearance in patient with UC.....	66

Fig. no.	Title	Page no.
30	Oral contrast enhanced positron emission tomography computed tomography images in a patient with symptomatic Crohn's disease.....	67
31	High quality images of the small intestine acquired by true FISP sequence	72
32	Coronal true FISP	73
33	Coronal half-Fourier acquisition single-shot turbo spin-echo (HASTE) image	74
34	35-year-old man with suspected Crohn's disease.....	75
35	Mesenteric lymphadenopathy in a patient with Crohn's disease	75
36	(A) Coronal T1post contrast showing the normal diffuse subtle homogenous bowel enhancement. (B&C) axial T1WI post contrast showing (B) Pathological accentuated enhancement of the terminal ileum, and (C) Engorged vasa recta known as comb sign	78
37	Male to female ratio	90
38	Final diagnosis of the included patients	90
39	Distribution of bowel wall thickening.....	91
40	Enhancement patterns	93
41	Lymphadenopathy percentage.....	93
42	Mean age of IBD cases.....	95
43	Sex predilection in IBD cases	95
44	Percentage of MR activity in IBD cases	101
45	Correlation between MRE results and clinical results in activity assessment of CD	102
46	Correlation between MRE results and clinical results in activity assessment of UC	103
47	Correlation between DWI restriction and disease activity	106

Fig. no.	Title	Page no.
48	Correlation between lymphadenopathy, engorged vasa recta and MR disease activity	107
49	Percentage of complications in IBD	108
50	(a), (b) Coronal T2WI single shot fast spin echo. (c) Coronal T2 SSFP showing thickened terminal ileum ..	111
51	Sagittal T2WI showing the terminal ileum mural thickening till the ileocecal junction	112
52	(a, c)Axial T2WI SPIR, (b ,d)Axial inverted diffusion	113
53	Axial inverted diffusion showing multiple restricted pericolic lymphadenopathy	113
54	Axial post contrast T1WI fat saturated showing the terminal ileum accentuated mural enhancement with the characteristic trilaminar appearance composed of an inner enhancing ring produced by the hyperemic mucosa.....	114
55	Coronal T1WI post contrast showing the trilaminar enhancement pattern of the affected ileal loop	115
56	(a), (b), and (c), Coronal T2WI single shot fast spin echo showing a long segment of diffuse mural thickening seen implicating the terminal ileum, (d) coronal T2WI SSFP showing same finding	118
57	(a) axial T2WI showing the diffuse mural thickening with mural high signal intensity, (b) inverted DWI showing corresponding restricted diffusion	119
58	Axial post contrast T1WI with fat saturation showing (a) appreciable contrast enhancement of the terminal ileum mucosa, (b) post contrast engorged vasa recta giving "the comb sign" impressive of active inflammatory process	119
59	Post contrast coronal T1WI showing stratified mural enhancement of the thick walled ileal segment giving the characteristic target sign appearance.....	120

Fig. no.	Title	Page no.
60	(a) Axial T2WI still showing the diffuse mural thickening of the terminal ileum with newly developed mesenteric localized fluid collection with air – fluid level and thickened wall, (b) Axial DWI showing restricted diffusion within center of the collection, (c) corresponding lowered ADC, (d–f) Axial T1WI post contrast with fat saturation showing: (d) layered pattern of enhancement of the terminal ileum, (e) the localized collection shows the air fluid level with marginal enhancement, (f) marked engorgement of the surrounding mesenteric vessels (comb sign) is seen (arrow), picture impressive of active Crohn's disease with abscess formation.....	121
61	(a), (c) & (e) Axial T2WI. (b),(d)& (f) Axial T2 SPIR, showing the long segment of distal ileal loop located behind the lower abdominopelvic wall showing mural thickening of maximum 6.7 mm with submucosal edema appreciated high signal intensity on SPIR images	124
62	Axial inverted DWI showing restricted diffusion of the distal ileal loops, associated restricted mesenteric lymphadenopathy are also noted	125
63	Axial T1WI post contrast with fat saturation showing appreciable post contrast enhancement of the distal ileal loop	125
64	(a) Axial T2WI, (b) Axial T2 SPIR showing diffuse mural thickening of the caecum with maximum wall thickening 8.9 mm, with mural high signal intensity best appreciated on SPIR image impressive of submucosal edema.....	127
65	(a) Axial T2WI, (b) Axial T2 SPIR showing diffuse mural thickening of the ascending colon, with mural high signal intensity best appreciated on SPIR image impressive of submucosal edema.....	127