### Effect of Two Lining Procedures on the Marginal Integrity of Compound Posterior Direct Resin Composite Restoration

#### Thesis

Submitted to the Faculty of Oral and Dental Medicine, Cairo University for Partial fulfillment of the Requirements for the Master Degree in Operative Dentistry

By

Nehal Hamdi Mohammed Ali B.D.S. (October 6<sup>th</sup> University) 2007

Faculty of Oral and Dental Medicine
Cairo University
(2016)

### **Supervisors**

#### Dr. Mohsen Hussein Abi-Elhassan

Professor of Operative Dentistry

Faculty of Dentistry

Cairo University

#### **Dr. Mohamed Fouad Haridy**

Assoc. Professor of Operative Dentistry

Faculty of Dentistry

Cairo University

# بسم الله الرحين الرحيم



سورة طه الاية رقم (١١٤) صدق الله العظيم

This research is dedicated to my great father for his advice and encouragement.

My mother, as always she has supported me throughout all that I have done. Thank you for your endless prayers and much needed love throughout my years.

My sisters, their love, and support have lightened up my spirit to finish this study.

My beautiful wonderful and helpful daughter Nancy for making me always happy.

Most of all thanks to Allah who continues to make the impossible possible. Thank Allah for making me who I am today and for giving me the abilities that I have.

I would like to express my deepest thanks and gratitude to **Prof. Dr. Mohsen Abi El-Hassan**, Professor of Operative Dentistry, Faculty of Dentistry, Cairo University, for his assistance, guidance, encouragement, kindness and supervision during all the steps of the present work.

I must thank **Dr. Mohamed Fouad Haridy**, Assoc. Prof. of Operative Dentistry, Faculty of Dentistry, Cairo University, for his great efforts and valuable advice to introduce this useful study.

Finally, I would like to thank everybody helped me by anyway and supported me during this study.

### List of contents

Title	Page
LIST OF TABLE	Ι
LIST OF FIGURES	II
INTRODUCTION	1
REVIEW OF LITERATURE	4
AIM OF THE STUDY	72
MATERIALS &METHODS	73
RESULTS	88
DISCUSSION	104
SUMMERY &CONCLUSIONS	115
REFRENCES	121
ARABIC SUMMARY	1

## List of tables

<b>Table (1):</b> Materials composition, manufacturers and lot	
numbers	74
Table (2): Variables of the study	75
Table (3): Interaction of variables for microleakge	
study	<b>76</b>
Table (4): Statically analysis for the effect of lining	
materials on leakage score percentages	89
Table (5): The mean, standard deviation (SD) values of	
leakageof all groups	92
Table (6): The mean, standard deviation (SD) values of	
leakage regardless of testing period.	94
Table (7): The mean, standard deviation (SD) values of	
leakage regardless of liner type	96

# **List of Figures**

Figure (1): Nano hybrid resin composite (Teric N
Ceram)
Figure (2): Nano-filled, fluoride releasing single
component adhesive (Exite F Viva Pen)
Figure (3): Ortho-Phosphoric acid (N-Etch)
Figure (4): Self adhesive flowable composite (Fusio
Liquid Dentin)
Figure (5): Light cured resin modified glass ionomer
(Riva Light Cure)
Figure (6): Tooth before cavity preparation
Figure (7): Tooth after cavity preparation
Figure (8): Digital Caliper for measuring the cavity
dimensions
Figure (9): LED curing unit.
Figure (10): Computer controlled material testing
machine
Figure (11): Tooth surfaces covered with two coats of
fingernail polish
Figure (12): The specimens immersed in 5% red
fuschin.
<b>Figure (13):</b> The samples stored in incubator at 37°C
Figure (14): Teeth after removal from stain immersion

Figure (15): Sectioning of teeth for microleakage	
assessment	
Figure (16): SPI module sputter coating machine	
Figure (17): The specimens coated with gold sputter	
machine	
Figure (18): Analysis for microleakage under	
scanning electron microscope	
Figure (19): Bar chart representing leakage after 24	
hours groups	;
Figure (20): Bar chart representing leakage	
percentage of after three months groups	
Figure (21): leakage scores for resin composite	
restoration with different lining protocols	
Figure (22): Bar chart representing Leakage	
regardless of testing periods	
Figure (23): Bar chart representing Leakage scores	
regardless of Liner type	
Figure (24): A photomicrograph showing grade zero	
gingival microleakage of restored upper premolar	
tooth under the stereomicroscope (after 24h control	
group)	
Figure (25): A photomicrograph showing grade zero	
gingival microleakage of restored upper premolar	
tooth under the stereomicroscope (after 3 months	
control group)	

Figure (26): A photomicrograph showing grade zero	
gingival microleakage of restored upper premolar	
tooth under the stereomicroscope (after 24h self-	
adhesive flowable composite group	98
Figure (27): A photomicrograph showing grade zero	
gingival microleakage of restored upper premolar	
tooth under the stereomicroscope (after 3 months self	
adhesive flowable composite group	98
Figure (28): A photomicrograph showing grade one	
gingival microleakage of restored upper premolar	
tooth under the stereomicroscope (after 24h resin	
modified glass ionomer group)	99
Figure (29): A photomicrograph showing grade one	
gingival microleakage of restored upper premolar	
tooth under the stereomicroscope (after 3 months resin	
modified glass ionomer group)	99
Figure (30): Scanning electromicrograph of	
resin/dentin interface in control group – 24h (×1500)	101
Figure (31): Scanning electromicrograph of	
resin/dentin interface in control group -3 months	
(×1500)	101
Figure (32): Scanning electromicrograph of	
resin/dentin interface in self adhesive flowable	
composite group – 24h (×1500)	102

	Figure	(33):	Scanning	electromicrog	raph of	
	resin/den	itin int	erface in	self-adhesive	flowable	
composite group – 3 months interface (×1500) 10						102
	Figure	(34):	Scanning	electromicrog	raph of	
resin/dentin interface in resin modified glass ionomer						
group –24h (× 1500)				103		
	Figure	(35):	Scanning	electromicrog	raph of	
resin/dentin interface in resin reinforce glass ionomer						
	group – 3	3 months	s (× 1500)			103

المشرفون درمصن حسين أبي الحسن أبي الحسن أبي الحسن أبي الحسن أستاذ ورئيس قسم العلاج التحفظي كلية طب الفم و الاسنان جامعة القاهرة عميد كلية طب الأسنان – جامعة الفيوم

د/محمد فؤاد هريدي أستاذ مساعد العلاج التحفظي كلية طب الفم و الاسنان جامعة القاهرة

# تاثير طريقتين للتبطين على التماسك الحافي لترميمات الكومبوزيت الراتنجية المركبة المباشرة الخلفية

رسالة مقدمة الى
كلية طب القم و الاسنان جامعة القاهرة
كجزء من متطلبات الحصول على درجة الماجستير
(فرع العلاج التحفظي)

الطبيبة/ نهال حمدي محمد علي بكالوريوس طب و جراحة الفم و الاسنان جامعة ٦ اكتوبر

جامعة القاهرة ٢٠١٦

## لجنة الحكم والمناقشة

### أ.د. مصطفى عبد الحميد مصطفى

أستاذ بقسم العلاج التحفظي جامعة القاهرة.

### أ.د. محمد على اليسقى

أستاذ بقسم العلاج التحفظي جامعة الأزهر - عميد جامعة سيناء.

### أ.د. محسن حسين ابي الحسن

أستاذ ورئيس قسم العلاج التحفظي جامعة القاهرة.

### اً.م.محمد فؤاد هريدي

أستاذ مساعد بقسم العلاج التحفظي جامعة القاهرة ومعار للجامعة البريطانية.

#### <u>**Judgement Committee**</u>

#### Dr. Moustafa Abdullhamid Moustafa

Professor of Operative Dentistry
Faculty of Dentistry, Cairo University.

### Dr. Mohammed Ali Elyasaki

Professor of Operative Dentistry
Faculty of Dentistry, Al-Azhar University.

#### Dr. Mohsen Hussein Abi-Elhassan

Professor of Operative Dentistry Faculty of Dentistry, Cairo University.

### **Dr. Mohammed Fouad Haridy**

Assc. Professor of Operative Dentistry Faculty of Dentistry, Cairo University.