



Ain Shams University
Faculty of Medicine

ROLE OF ULTRASONOGRAPHY VERSUS MRI IN DIAGNOSIS OF ANKLE LESIONS

Essay

*Submitted for Partial Fulfillment of Master Degree
in Radio Diagnosis*

Presented By

Dalia Nazeeh Azer

*M.B., B.Ch,
Faculty of Medicine-Ain Shams University*

Under Supervision of

Prof. Dr. Marwa Ibrahim Fahmy

*Professor of Radiodiagnosis
Faculty of medicine - Ain shams university*

Dr. Yasser Ibrahim Abdel-Khalek

*Lecturer of Radiodiagnosis
Faculty of medicine - Ain shams university*

*Faculty of medicine
Ain shams university*

2016



Aknowledgement

*First and foremost, I feel indebted to **God**, the most merciful, who gave me the power to accomplish this work,*

*I wish to express my gratitude to my supervisor, **Prof. Dr. Marwa Ibrahim Fahmy**, Professor of Radiodiagnosis, Faculty of Medicine, Ain Shams University. Her unique supervision, meticulous advice and the long time and tremendous effort she offered were invaluable in accomplishing this work,*

*I also express my deepest thanks to **Dr. Yasser Ibrahim Abdel-khalek**, Lecturer of Radiodiagnosis, Faculty of Medicine, Ain Shams University for his patience, friendly attitude and for being generous with time and effort.*

Finally, Many thanks and appreciation to all those who generously supported me by all means, the staff of Radiodiagnosis Department, Ain Shams University.



List of Contents

<i>Title</i>	<i>Page No.</i>
List of Tables	I
List of Figures.....	II
List of Abbreviations	XIX
Introduction	1
Aim of the Work.....	3
Review of Literature	
- Chapter (1): Anatomy of the ankle.....	4
- Chapter (2): Pathological lesions of the ankle	40
- Chapter (3): MRI technique of ankle joint examination.....	64
- Chapter (4): US technique of ankle joint examination ...	72
- Chapter(5): Ultrasonographic and MRI manifestations of ankle lesions	100
- Chapter (6): Summary and Conclusion	177
References	190
Arabic Summary	-

List of Tables

<i>Table No.</i>	<i>Title</i>	<i>Page No.</i>
------------------	--------------	-----------------

Table (1): The signal intensity of various tissues 29

Table (2): Ankle MR imaging protocols at 1.5 and 3 pt.... 71

List of figure

<i>Figure No.</i>	<i>Title</i>	<i>Page No.</i>
Fig (1):	T1-weighted axial MR image showing the anterior compartment.....	5
Fig (2):	Anterior axial sonographic image shows the three tendons of the anterior compartment.....	7
Fig (3):	Anterior longitudinal sonographic image along the course of the tibialis anterior tendon.....	7
Fig (4):	Anatomic diagram showing the superior and inferior extensor retinacula	8
Fig (5):	Anterior axial sonographic image obtained at the level of the tibiofibular syndesmosis.....	9
Fig (6):	T1-weighted axial MR image.(a)The medial compartment of the ankle. Coronal MR image (b) clearly depicts the superficial and deep fasciculi of the medial collateral ligament and, superficially, the flexor retinaculum.....	10
Fig (7):	Sonographic image of the medial retromalleolar area obtained in the axial-oblique plane.	11
Fig (8):	Diagram showing the flexor retinaculum, which extends from the tibial malleolus to the medial aspect of the calcaneus	12
Fig (9):	Sonographic image of the medial submalleolar region	15
Fig (10):	Axial proton density (PD)-fat suppression (FS) MR image: The lateral compartment of the ankle	16
Fig (11):	Axial sonographic image of the retromalleolar region.....	18
Fig (12):	Anatomic diagram showing the superior and inferior peroneal retinacula.	19
Fig (13):	Axial PD FS MR image: the examination reveals an accessory peroneus quartus tendon.....	20
Fig (14):	Anterior perimalleolar sonogram.	21
Fig (15):	T1-weighted axial MR image (a) depicting the calcaneofibular .(b) shows the calcaneofibular ligament running beneath the peroneal tendons.....	22
Fig (16):	Coronal PD FS MR image: the posterior tibiotalar ligament	23
Fig (17):	T1W axial (a) and T1W sagittal (b) MR images. In the posterior compartment of the ankle	25
Fig (18):	Axial sonographic images of a normal Achilles tendon.....	25

List Of Figures

Fig (19):	Sagittal sonographic image of the Achilles tendon (arrows) from its origin at the confluence of the soleus (deep) and gastrocnemius (superficial) muscles to its insertion on the posterior surface of the calcaneus, distal to the posterosuperior tuberosity	26
Fig (20):	Sagittal sonographic image obtained at the level of the posterior calcaneal tuberosity.	28
Fig (21):	(a) The complex insertion of the tibialis posterior tendon . (b)The magic artifact is almost completely diminished when the same volunteer is scanned in supine position.	31
Fig (22):	Skeletal muscle: cross section	32
Fig (23):	Longitudinal tendon sonogram.....	34
Fig (24):	Transverse oblique ultrasound of the distal tibiofibular joint in the longitudinal axis of the anterior talofibular ligament.	36
Fig (25):	Extremity coils used for ankle and foot imaging.....	66
Fig (26):	Image planes for foot and ankle MRI. (A) foot positioned for axial images of the ankles.(B) Image planes obliqued to obtain true coronal through the ankle mortise.. (C) Oblique sagittal image planes to improve anatomic display of the osseous structures in the ankle. (D) Conventional sagittal image plane to improve evaluation of the Achilles tendon.	67
Fig (27):	Anterior tendons group: US appearance. Transverse US images obtained over the distal tibia (a) and talar dome (b).....	74
Fig (28):	Anterior tendons group: US appearance. Transverse US images obtained over the talar dome (a) and sinus tarsi (b).	76
Fig (29):	Anterior inferior tibiofibular ligament: normal US appearance. TIB: tibia, FIB: fibula (A) Transverse sonogram shows anterior tibiofibular ligament (arrows). (B) Corresponding transducer position.	78
Fig (30):	Lateral tendon group: US appearance. Transverse US images obtained over the peroneal tendons at the supra malleolar and malleolar location.	81
Fig (31):	Peroneus quadratus. US image obtained over the retro malleolar groove demonstrates the peroneus longus and peroneus brevis tendons together with an additional element, the peroneus quadratus, consisting of a discrete tendon surrounded by a hypoechoic muscle.	82

List Of Figures

Fig (32):	Lateral tendon group: US appearance. Coronal oblique US images obtained over the peroneal tendons at the proximal (a) and distal (b) inframalleolar region.	83
Fig (33):	Lateral tendon group: US appearance. Sagittal oblique US images obtained over the peroneus brevis at the supramalleolar (a) and inframalleolar (b) region.....	83
Fig (34):	Lateral tendon group: US appearance. Sagittal oblique US images obtained over the peroneus longus (white arrowheads) at the supramalleolar (a) and inframalleolar (b) region.....	84
Fig (35):	Anterior talofibular ligament: normal US appearance. Transverse oblique US images of the ligament (arrowheads) obtained with the transducer parallel (a) and oblique (b) to it...	86
Fig (36):	Calcaneofibular ligament: normal US appearance.	87
Fig (37):	Medial tendon group. Schematic sagittal drawing of the medial tendons showing: the TP , FDL, and FHL tendons and Achilles tendon	88
Fig (38):	Medial tendon group: US appearance. Transverse US images obtained over the posteromedial tendons at the supramalleolar location, (a) anterior and (b) posterior images.	90
Fig (39):	Medial tendon group: US appearance. Sagittal oblique US images obtained over the proximal (a) tibialis posterior and (b) flexor digitorum longus tendon.	91
Fig (40):	Posteromedial tendon group: US appearance. Longitudinal US images obtained over the distal (a) tibialis posterior and (b) flexor digitorum longus.	92
Fig (41):	Normal flexor hallucis longus tendon. Sagittal US image of the posterior ankle obtained with the transducer placed over the Achilles tendon.....	93
Fig (42):	Medial ligament group: normal US appearance. Coronal oblique US images of the deep and superficial deltoid ligament.....	94
Fig (43):	Posterior tendon group: US appearance. Transverse and sagittal US images obtained over the Achilles tendon..	97
Fig (44):	(A) Photograph shows the site for measuring AT (B) Transverse U/S scan shows a normal AT.....	98
Fig (45):	Photograph shows the site for longitudinal scanning of the distal Achilles tendon and retrocalcaneal bursa.	99

List Of Figures

Fig (46):	US aspect of tendinopathies. Left: longitudinal US. Right: axial US	103
Fig (47):	Stenosing tenosynovitis of the flexor hallucis longus tendon in a ballet dancer who had recent onset of pain during plantar flexion of the foot. a: Longitudinal and, b: transverse US images of the posteromedial ankle obtained at the level of the talar tubercles..	107
Fig (48):	Transverse US shows marked enlargement and internal hypoechogenicity of the peroneus brevis tendon with surrounding tenosynovitis	108
Fig (49):	Tibialis posterior tendinosis and tenosynovitis. A: Short-axis sonographic image shows a normal echogenic tibialis posterior tendon and an adjacent normal flexor digitorum longus tendon.	109
Fig (50):	Post-traumatic peritendinitis and partial rupture of peroneal tendons. Axial proton-density MR image shows a rim of low signal intensity surrounding partially ruptured peroneus longus and brevis tendon.....	112
Fig (51):	Longitudinal sonography image of tibialis posterior tendinosis shows a thickened tendon and an area of interstitial degeneration	113
Fig (52):	Power Doppler image showing Achilles insertional tendinosis with neovascularization at the area of tendinosis	113
Fig (53):	Transverse scan of Achilles tendinosis. The tendon is thickened and inhomogeneous with a small focus of calcification	114
Fig (54):	High-grade tendinopathy in a patient with continuous pain in the Achilles tendon	115
Fig (55):	A) Acute Achilles tendinosis. Sagittal T2-weighted MR image.B) Chronic tendinosis of the Achilles tendon. Sagittal TWI.	116
Fig (56):	Complete rupture of the Achilles tendon. Long-axis extended field-of-view 12-5 MHz US image demonstrates the separation of the torn tendon ends. , because of an ill-defined hypoechoic hematoma	119
Fig (57):	Complete acute rupture of the Achilles tendon. Long-axis US image demonstrates a gap between the proximal and distal ends of the ruptured Achilles tendon.....	119

List Of Figures

- Fig (58):** (A) Transverse US shows a longitudinal split in the peroneus brevis tendon creating two separate portions of the brevis at the level of the lateral malleolus. (B) Transverse US from a different patient show a longitudinal split in the peroneus brevis tendon at the level of the lateral malleolus 121
- Fig (59):** Short-axis sonographic image of the tibialis posterior tendon shows tendinosis and a linear hypoechoic interstitial split tear. The tendon is surrounded by a markedly thickened tendon sheath 122
- Fig (60):** Partial-thickness (type I) tear of the tibialis posterior tendon. (a) Long and (b) short-axis US images at the level of the medial malleolus 123
- Fig (61):** Longitudinal split of the tibialis posterior tendon. A) Transverse US shows the tibialis posterior tendon with a hypoechoic region extending to the deep surface of the tendon. B) Longitudinal US of the tibialis posterior tendon showing a corresponding linear region of hypoechogenicity 124
- Fig (62):** Complete tear of the tibialis posterior tendon, longitudinal sonogram obtained over the supramalleolar portion of the TP tendon showing a complete tear and the swollen hypoechoic and irregular proximal tendon 124
- Fig (63):** Sagittal proton density A) demonstrates grade I tear of Achilles tendon and T2 weighted image B) demonstrate high signal intensity with thickening and retraction commonly seen with complete tear of the Achilles tendon. 125
- Fig (64):** Sagittal T2 weighted image demonstrating a tear and retraction of the FHL 126
- Fig (65):** Sagittal (A) and axial B) T2-weighted image demonstrate focal thickening of the retinacula with increased signal intensity resulting from partial tear in the anterior tibial tendon 126
- Fig (66):** Traumatic complete rupture (type III) of the Achilles tendon, (A): Sagittal T1WI shows a gap represent complete tendon rupture, (B): Corresponding axial T2WI demonstrates fluid or edema at the gap. 127

List Of Figures

- Fig (67):** TP tendon tear (type 3) Axial T2WI at the level of the talus (A) and distally near the navicular (B) show fluid around the tendon with no definite tendon proximally in A and thickening in B distally. Sagittal gradient echo images (C, D) show the isolated FDL with high signal intensity in region of TP tendon.127
- Fig (68):** Peroneal tendon instability, transverse US images over the retro-malleolar groove obtained (a) at rest and (b) during forced eversion and dorsiflexion of the foot..... 128
- Fig (69):** 20-year-old woman with retro-fibular intra-sheath subluxation of left peroneal tendon. Axial sonogram obtained with affected ankle in neutral position shows peroneus longus and peroneus brevis tendons in normal relationship posterior to distal fibula. 130
- Fig (70):** Traumatic dislocation of the right posterior tibial tendon. MR image of both ankles shows the right posterior tibial tendon located anteromedial to the medial malleolus. 131
- Fig (71):** Sonogram of an ankle with an old ATFL sprain shows irregularities of the tip of the lateral malleolus due to a small healed avulsion. 134
- Fig (72):** US scan of lateral compartment of ankle. The ATFL is continuous but thickened with a hypoechoic line underscoring its superficial part for a 1st degree 135
- Fig (73):** US scan of lateral compartment of ankle. The ATFL is completely torn with abundant effusion 135
- Fig (74):** Sonogram of an ankle with an old sprain of the AITFL shows bulging of the ligament and capsule..... 136
- Fig (75):** A) Sonogram of an ankle with an old sprain shows hypoechogenicity, enlargement, and loss of tension of the CFL. B) Sonogram of the same patient's contralateral ankle shows the normal CFL 137
- Fig (76):** Sonogram of an ankle with a 3-month-old sprain of the CFL shows the deep location of the fibular tendons owing to the lack of tension in the enlarged, hypoechoic ligament 138
- Fig (77):** Longitudinal US scan shows hypoechoic thickening in the expected location of the ATFL with an avulsion fracture of the talar insertion. 138

List Of Figures

Fig (78):	Partial-thickness tear of the DL. Coronal 12-5 MHz US image over the medial ankle reveals hypoechoic thickening of the cranial portion of the tibiocalcaneal ligament consistent with a partial tear. Discrepant size between the injured (arrows) and the intact (arrowheads) portion of the ligament is noticed. C: calcaneus	140
Fig (79):	Partial-thickness tear of the DL in a patient presenting with chronic pain and swelling over the medial ankle. Coronal 12-5 MHz US image over the medial ankle reveals a hypoechoic cleft involving the deep portion of the deltoid ligament suggesting a partial tear	140
Fig (80):	Axial proton density MR image showing partial tear (arrow) of posterior inferior tibiofibular ligament.	142
Fig (81):	Oblique axial T2-weighted image of partial tear of calcaneofibular ligament shows fluid signal in ligament and mildly irregular contour.....	143
Fig (82):	Coronal proton density fat suppression MRIs in two cases of distal fibular fracture . a) showing bright fluid signal transects the posterior tibiotalar ligament representing complete tear (arrow) ,B) shows heterogeneous posterior tibiotalar ligament	146
Fig (83):	Tibialis posterior tenosynovitis (a) Axial ultrasound showing a minor widening of the tendon with a hypoechoic rim representing tenosynovitis.(b) Longitudinal section shows a widened tendon with fluid in the tendon sheath and synovial hypertrophy . (c) Power Doppler shows hypervascularity in the tendon and surrounding synovium.	148
Fig (84):	A 50-year-old woman with synovitis from rheumatoid arthritis. Sagittal sonogram shows ankle joint distention with predominantly hypoechoic synovium.....	148
Fig (85):	Rheumatoid arthritis, Coronal T1-weighted (a) and sagittal STIR (b) MR images show fluid and debris filling the sinus tarsi as well as subchondral edema and cysts	150
Fig (86):	A 64 yrs man with gout. Longitudinal sonogram of Tibialis posterior tendon shows distortion of distal tendon (T) and adjacent hypoechoic tenosynovitis	151
Fig (87):	A 64 yrs man with gout. Longitudinal sonogram of Tibialis posterior tendon shows distortion of distal tendon (T) and adjacent hypoechoic tenosynovitis	152

List Of Figures

- Fig (88):** Gouty arthritis of ankle (A) and (B) coronal T1-weighted images through the medial and posterior ankle joint show low signal intensity soft tissue foci associated with erosive changes. Axial T1-weighted fat suppressed, enhanced images of the ankle joint (C) and sinus tarsi (D) show marked enhancement of the tophi through the ankle with synovial enhancement of the extensor and peroneal tendon sheaths. ...154
- Fig (89):** Partial rupture of the ATFL , Long-axis 12-5 MHz US image over the ATFL in a patient following an inversion injury of the ankle.155
- Fig (90):** Partial rupture of the CFL in a patient with a torn ATFL. (a) Long-axis and (b) short-axis US scans show a markedly swollen and hypoechoic CFL , just deep to the peroneal tendon.156
- Fig (91):** Complete rupture of the ATFL. Long-axis US images obtained over the ATFL in two different patients only a few hours following an inversion injury of the ankle.....156
- Fig (92):** Sonographic anterior drawer test. A) Photograph demonstrates how to perform a stress test during scanning of the ATFL.157
- Fig (93):** Anterolateral impingement. A) Sagittal T1-weighted image shows low intensity soft tissue thickening around the lateral malleolus. B: Sagittal gradient echo image shows soft tissue enhancement and C) sagittal post-gadolinium image shows anterior tibial osteophyte.159
- Fig (94):** Anterior impingement syndrome. A) Longitudinal 12-5 MHz US image with b) T2*-weighted MR imaging correlation demonstrates a prominent spur at the distal end of the anterior tibia.....160
- Fig (95):** Stenosing tenosynovitis of the flexor hallucis longus tendon in a ballet dancer during plantar flexion of the foot. a, b) Sagittal and .c) transverse 12-5 MHz US images of the posteromedial ankle163
- Fig (96):** Partial-thickness tear of the deltoid ligament. Coronal US image over the medial ankle shows hypoechoic thickening of the cranial portion of the tibiocalcaneal ligament consistent with a partial tear.165

Fig (97):	Partial-thickness tear of the deltoid ligament . a) Coronal 12-5 MHz US image over the medial ankle with b) arthro-MR imaging.....	166
Fig (98):	Tarsal tunnel syndrome in a patient with tenosynovitis of the flexor hallucis longus tendon. a) Transverse 12-5 MHz US image over the posteromedial ankle shows displacement and compression of the tibial nerve by the distended sheath of the flexor hallucis longus tendon . b) diagram illustrates the relationship of the distended sheath of the flexor hallucis longus with the structures in the tarsal tunnel, including the tibial nerve, tibialis posterior and flexor digitorum longus tendons	167
Fig (99):	Tarsal tunnel syndrome in a patient with a tibiotalar ganglion cyst. a) Transverse 12-5 MHz US image of the posteromedial ankle with b) Schematic drawing illustrates the relationship of the ganglion cyst with the structures contained in the tarsal tunnel, including the tibial nerve, tibialis posterior, flexor digitorum and flexor hallucis longus tendons	168
Fig (100):	A schwannoma is identified within the tarsal tunnel on T1 weighted sagittal and fat-suppressed T2 weighted axial images.....	169
Fig (101):	This patient with symptoms of posterior tibial neuropathy was found to have varicosities (arrows) within the tarsal tunnel, well seen on this fat-suppressed T2-weighted axial image	169
Fig (102):	Sinus tarsi syndrome in a patient with RA. Sagittal T1-weighted MR image shows obliteration of fat by an area of fluid-like signal intensity in the subtalar joint.....	171
Fig (103):	Os trigonum syndrome A) Axial T2 image demonstrates edema and fluid around FHL. B) Sagittal PD image demonstrates fluid and marginal tibiotalar edema due to impingement	172
Fig (104):	Tarsal tunnel syndrome secondary to ganglion cyst. Axial T2-weighted MR image reveals a ganglion cyst with bright signal of its fluid content, interposed between the flexor digitorum longus (d) and flexor hallucis longus tendons and abutting the adjacent neurovascular bundle	175
Fig (105):	54-year-old woman with schwannoma. Longitudinal sonogram of the posterior tibial nerve shows solid and relatively homogeneous mass adjacent to talus and continuous with the nerve	176

List of Abbreviations

Abb.	Full term
AITFL	Anterior inferior tibio-fibular ligament.
CFL.....	Calcaneofibular ligament.
DWI.....	Diffusion weighted image.
EDL	Extensor digitorum longus.
EHL	Extensor hallucis longus.
FC	Calcaneofibular ligament.
FDL.....	Flexor digitorum longus.
FHL	Flexor hallucis longus
FTA.....	Anterior talofibular ligament.
FTP	Posterior talofibular ligament.
KFS.....	Kager fat space
LM.....	Lateral malleolus.
ML	Medial malleolus.
MRI	Magnetic resonance imaging.
PB	Peroneus brevis.
PQ.....	Peroneus quartus.
PL	Peroneus longus.
PT	Peroneus tertius.
PTFL	Posterior talofibular ligament.

List Of Abbreviations

<i>Abb.</i>	<i>Full term</i>
RA.....	Rheumatoid arthritis.
TA.....	Tibialis anterior.
TP	Tibialis posterior.
TTS.....	Tarsal tunnel syndrome.
T1WI	T1 weighted image.
T2WI	T2 weighted image.
US	Ultrasound