

بسم الله الرحمن الرحيم



-Call 6000





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم





جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة يعيدا عن الغيار













بالرسالة صفحات لم ترد بالأصل



An evaluation of two active lingual arches used for resolving moderate lower anterior crowding in mixed dentition

Thesis

Submitted in partial fulfillment for the requirements of Master Degree in Pedodontics

By Ahmed Ezat Abdel Azem Dawood B.D.S.(1998)

> Faculty of Dentistry Tanta University 2005

> > B 11197

المالة مرالت مرالت مرابية

Supervisors

Dr. Medhat Mohamed El Sakhawy

Professor and Head of Orthodontic Department
Faculty of Dentistry
Tanta University

Dr. Azza Mahmoud Tag El Din

Ass. Prof. of pedodontics

Pedodontics & Preventive Dentistry Department

Faculty of Dentistry

Tanta University

Dr. Talat Mohamed Beltagy

Lecture of Pedodontics

Pedodontics & Preventive Dentistry Department

Faculty of Dentistry

Tanta University

DEDICATION

Dedicated to

my parents &

my wife and

my daughter Basant.

ACKNOWLEDGEMENT

I would like to express my deepest thanks and appreciation to professor Dr. **Medhat Mohamed El Sakhawy** Professor and Head of Orthodontic Department, Faculty of Dentistry, Tanta University for his high appreciated advice and encouragement.

I would like also to express my sincere gratitude to **Dr. Azza Mahmoud Tag El Din** Ass. Prof. of pedodontics & Preventive Dentistry, Faculty of Dentistry, Tanta University for her help and continuous guidance throughout the course of this research.

I am also grateful to **Dr. Talat Mohamed Beltagy** Lecture of Pedodontics & Preventive Dentistry,
Faculty of Dentistry, Tanta University for his help throughout
the course of this research.

Contents

	Page
LIST OF FIGURES	
LIST OF TABLES	
I. INTRODUCTION	1
II. REVIEW OF LITERATURE	3
- Etiology of crowding	4
- Criteria of crowding	8
- Development of the mixed dentition	9
- Arch length analysis	11
- Dental arch development	17
- Study of crowding	24
- Management of crowding	27
III. AIM OF THE WORK	34
IV. MATERIALS AND METHODS	35
V. RESULTS	47
VI. DISCUSSION	77
/II. SUMMARY AND CONCLUSIONS	87
III. REFERENCES	91
IX. PROTOCOL	
X. ARABIC SUMMARY	
• •	

LIST OF FIGURES

Figures	Title	Page
Fig.(1):	Friel's spring lingual arch design appliance.	45
Fig.(2):	Finger's spring lingual arch design appliance.	45
Fig.(3):	Pedodontic diagnostic chart.	46
Fig.(4):	Children tolerance in the treated subgroup (A)	49
	Friel's spring & treated subgroup (B) Finger's	ļ
	spring during study periods.	
Fig.(5):	Percentage of appliance breakage in the treated	49
	subgroup (A) Friel's spring & treated subgroup	
7.	(B) Finger's spring during study periods.	
Fig.(6):	Percentage presence of tongue trauma in the	50
	treated subgroup (A) Friel's spring & treated	
	subgroup (B) Finger's spring during study	
Fic (7).	periods.	
Fig.(7):	Percentage of gingivitis in the treated subgroup	50
	(A) Friel's spring & treated subgroup (B)	
Fig.(8):	Finger's spring during study periods.	50
11g.(8).	A case before and after treatment in treated	52
Fig.(9):	subgroup (A) Friel's spring. Cephalometria, tracing of malar & includes the control of the cont	C 4
1 16.(2).	Cephalometric tracing of molar & incisor position in treated subgroup (A) Friel's spring.	54
Fig.(10):	A case before and after treatment in treated	56
1 18.(10).	subgroup (B) Finger's spring.	30
Fig.(11):	Cephalometric tracing of molar & incisor	58
	position in treated subgroup (B) Finger's spring.	
Fig.(12):		60
	anterior crowding.	
Fig.(13):	Cephalometric tracing of molar & incisor	62
	position in control group (C).	
Fig. (14):	The changes in intercanine width during the	73
	study period.	

Fig. (15):	The changes in intermolar width in the studied	73
	groups during study periods	
Fig. (16):	Arch depth changes in the studied groups during	73
·-	study periods.	
Fig. (17):	Mean leeway space in the studied groups during	74
	study periods.	
Fig. (18):	Mean arch length diameter in the studied groups	74
	during study periods.	
Fig. (19):	Mean SNA value in the studied groups before	75
	and after treatment.	
Fig. (20):	Mean SNB value in the studied groups before	75
	and after treatment.	
Fig. (21):	Mean molar position value in the studied groups	75
	before and after treatment.	
Fig. (22):	Mean incisor position in studied groups before	76
	and after treatment.	
Fig. (23):	Mean IIA value in studied groups before and	76
	after treatment.	

LIST OF DIAGRAMS

Diagram	Title	Page
Diagram(1):	Anterior crowding in mixed dentition.	37
Diagram(2):	Measuring points in the study models.	38

LIST OF TABLES

Table	Title	Page
Table(1):	Clinical assessment score.	44
Table(2):	Clinical assessment for treated groups during study periods.	47
Table(3):	Mean & standard deviation of linear measurements in treated subgroup (A) Friel's spring during study period.	51
Table(4):	Mean & standard deviation of angular measurements in treated subgroup (A) Friel's spring during study period.	53
Table(5):	Mean & standard deviation of linear measurements in treated subgroup (B) Finger's spring during study period.	55
Table(6):	Mean & standard deviation of angular measurements in treated subgroup (B) Finger's spring during study period.	57
Table(7):	Mean & standard deviation of linear measurements in control group (C) during study period.	59
Table(8):	Mean & standard deviation of angular measurements in control group (C).	61
Table(9):	Comparisons of mean and standard deviation in the linear measurements between the studied groups during study periods.	65
Table(10):	Spearman's correlation coefficient of linear measurements between the studied groups.	66
Table(11):	Comparisons of mean and standard deviation in angular measurements from cephalometric films	67