



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

*The Prevalence of Cigarette Smoking
among Egyptian Physicians
(In Cairo)*

Thesis

**Submitted for Partial Fulfillment of Master Degree
in Public Health**

By

**Nouran Hamdy Seif Elnasr
M.B., B.Ch, 1996**


Supervised By

**Dr/Ibrahim Soliman Hegazy
Professor of Public Health
Faculty of Medicine- Cairo University**

**Dr/ Omaila Kamel ElSalamony
Assistant professor of Public Health
Faculty of Medicine- Cairo University**

2001

B N C N.





بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

وَكَانَ فَضْلُ اللَّهِ عَلَيْكَ عَظِيمًا

سورة النساء الآية 113

صَدَقَ اللَّهُ الْعَظِيمُ

This work wouldn't have been carried out without Allah's great mercy and well.



100-100
(100)

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

100-100

Abstract

Smoking habits were studied among 750 physicians working in different healthcare facilities in four health districts in Cairo Governorate, in the year 2000, representing about 3.5% of the total number of physicians registered in Cairo Physicians Syndicate.

Smoking physicians were 170 (165 males, 5 females), with an overall prevalence of cigarette smoking 22.7% (32.5% of males, 2.1% of females). The most commonly associated health problems were cardiovascular diseases 10.6%, chronic respiratory diseases 12.4% & chronic renal problems, 1.2% with a statistically significant higher prevalence of renal problems.

Tendency for quitting was observed in about 68.2%, most of them (85.9%), without any kind of assistance.

Key words: Smoking - Physicians - Cairo

Acknowledgment

I wish to express my deepest gratitude to all those who sincerely helped me in this work.

I am greatly indebted to **Prof. Dr. Ibrahim Hegazy**, Professor of Community Medicine, Cairo University, for his kind supervision guidance, support and encouragement through the whole work.

It is my pleasure to express my unlimited appreciation and deepest gratitude to **Dr. Omaina K. ElSalamony**, Assistant professor of Community Medicine, Cairo University, without her continuous support, guidance and supervision the completion of this work would have been much difficult.

Finally, I want to express my deepest gratitude and thanks to my mother, who has always been a great support for me through my whole life.

Nouran Seif Elhasr

