

تبيكة المعلومات الحامعية

Cierra Territo Cierro Con





تبيكة المعلومات الجامعية



شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيل



جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار في درجة حرارة من 15 - 20 منوية ورطوبة نسبية من 20- 40 $^{\circ}$

To be kept away from dust in dry cool place of 15-25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية





ثبكة المعلومات الجامعية



الأصلية تالفة

The Role of Deferoxamine in Prevention of Aminoglycoside ototoxicity: An EM (Electron Microscopic) Study in Guinea Pigs.

A Thesis submitted for the partial fulfillment of the MD degree in OtoRhinolaryngology

Ву

Mohamed Amir Hassan

M.B.B.Ch., M.S. (OtoLaryngology.)
Ain Shams University, Faculty of Medicine

Supervised by
Prof. Dr. Fouad Abbas Ismail
Professor of OtoLaryngology
Ain Shams University

Prof. Dr. Somia Tawfik Mohamed

Prof. Dr. Badr Eldin Mostafa

Professor of Audiology

Professor of OtoLaryngology

Ain Shams University

Ain Shams University

Dr. Nadia Galal El-Hefnawy Assistant Professor of Pathology

Ain Shams University

Faculty of Medicine
Ain Shams University
2002

N Spire

Borg

بسم اللة الرحمن الرحيم

حدق اللة العظيم

سورة البقرة - الآية ٣٢



ACKNOWLEDGEMENT

I would like to express my sincere gratitude and cordial thanks to Prof. Dr. Fouad Abbas, Professor of Otolaryngology, faculty of medicine, Ain Shams University for his great encouragement and valuable guidance since the beginning of setting the idea of this thesis and for his continuous enthusiasm and patience all through the work, to him I will be always grateful, without him this work could have never been fulfilled.

I would like to express my deep thanks and sincere appreciation to Prof. Dr. Somia Tawfik, Professor and head of the department of Audiology, faculty of medicine, Ain Shams University for her indispensable guidance, great help and kind support.

I am particularly indebted to Prof. Dr. Badr Eldin Mostafa, Professor of otolaryngology, faculty of medicine, Ain Shams University who gave me so much of his valuable time and immense knowledge to produce this work in it's present form.

I am also very grateful to **Dr. Nadia Galal**, Assistant Professor of Pathology, Ain Shams University for her great help in the processing, preparation and interpretation of the electron microscopy specimens.

I would like to thank Dr. Ali Hussein, ENT consultant Shebeen El-Kom Hospital for his help in conducting this experimental work.

Finally, I'm indebted to my family especially my wife, whose care, patience and sincere help have always been the pounce of light to guide me.

	LIST OF CONTENTS H	Page
•	List of tables	i
•	List of figures	.ii
•	List of abbreviations	V
•	Introduction and Rationale	.1
•	Aim of the work	
•	Review of literature	4
	- Functional anatomy of the cochlea	
	- Anatomy of the ear in Guinea pigs	
	- Aminoglycosides	
	- Deferoxamine.	
	- Evaluation of cochlear function in experimental animals	S
•	Methodology	50
•	Audiological Results6	50
•	Histological. Results	82
•	Discussion	20
•	Conclusions	27
•	Recommendation12	28
•	Summary12	29
•	References1	
•	Arabic Summary16	51

- --

List of tables

	Table 1	Comparing between DP amplitude pre and post-injection in	61
1		control group.	
 	Table 2	Comparing between S/N ratio pre and post-injection in	63
		control group.	
	Table 3	Comparing between DP amplitude pre and post-injection in	66
 		gentamicin group.	
	Table 4	Comparing between S/N ratio pre and post-injection in	68
	·	gentamicin group.	
	Table 5	Comparing Control group with gentamicin group (Pre-	70
	•	injection).	
 	Table 6	Comparing Control group with gentamicin group (Post-	71
		injection).	
İ	Table 7	Comparing between DP amplitude pre and post-injection in	74
	, . Y	gentamicin and deferoxamine group.	
; 	Table 8	Comparing between S/N ratio pre and post-injection in	76
		gentamicin and deferoxamine group.	
 	Table 9	Comparing Control group with deferoxamine + gentamicin	78
! !	~ ·	group (Pre-injection).	
;	Table 10	Comparing Control group with deferoxamine + gentamicin	79
		group (Post-injection).	
 -	Table 11	Comparing gentamicin group with deferoxamine +	80
		gentamicin group (Pre-injection).	
	Table 12	Comparing gentamicin group with deferoxamine +	81
		gentamicin group (Post-injection).	
		· · · · · · · · · · · · · · · · · · ·	

List of Figures

Figure 1	Diagram of anatomy of cochlea	4
Figure 2	Diagram of organ of corti	5
Figure 3	Diagram of organ of corti	9
Figure 4	Transmission EM diagram of outer hair cells	10
Figure 5	Transmission EM diagram of inner hair cells	.11
Figure 6	Scanning EM picture of surface of organ of corti	13
Figure 7	Diagram showing the role of tip links between stereocilia in transduction.	14
Figure 8	A= midmodiolar cross-section of the human cochlea. B= Cross-section through the guinea pig cochlea.	18
Figure 9	Mean DPOAE amplitudes Pre and Post-injection in control group.	62
Figure 10	Mean DPOAE signal to noise ratio Pre and Post-injection in control group.	64
Figure 11	Mean DPOAE amplitudes Pre and Post-injection in gentamicin group.	67
Figure 12	Mean DPOAE signal to noise ratio Pre and Post-injection in gentamicin group.	69
Figure 13	Mean DPOAE amplitudes Pre and Post-injection in gentamicin and deferoxamine group.	75
Figure 14	Mean DPOAE signal to noise ratio Pre and Post-injection in gentamicin and deferoxamine group.	77
Figure 15	Semithin section showing overall view of a cochlear turn	85
Figure 16	Semithin section of the organ of corti of control group	86
Figure 17	Semithin section of the organ of corti of control group	87
Figure 18	Semithin section of the organ of corti of control group	88
Figure 19	Semithin section of stria vascularis of control group	89

Figure 20	Semithin section of stria vascularis of control group	90
Figure 21	Ultrathin section of the basilar membrane of control group	91
Figure 22	Ultrathin section of the basilar membrane of control group	92
Figure 23	Ultrathin section of the outer hair cells of control group	93
Figure 24	Ultrathin section of the outer hair cell of control group	94
Figure 25	Ultrathin section of the inner hair cell of control group	95
Figure 26	Ultrathin section of the stria vascularis of control group	96
Figure 27	Semithin section of the organ of corti of gentamicin group	99.
Figure 28	Semithin section of the organ of corti of gentamicin group	100
Figure 29	Semithin section of stria vascularis of gentamicin group	101
Figure 30	Semithin section of stria vascularis of gentamicin group	102
Figure 31	Ultrathin section of outer hair cell of gentamicin group	103
Figure 32	Ultrathin section of outer hair cells of gentamicin group	104
Figure 33	Ultrathin section of outer hair cells of gentamicin group	105
Figure 34	Ultrathin section of outer hair cell of gentamicin group	106
Figure 35	Ultrathin section of inner hair cell of gentamicin group	107
Figure 36	Ultrathin section of stria vascularis of gentamicin group	108
Figure 37	Ultrathin section of stria vascularis of gentamicin group	109
Figure 38	Semithin section of organ of corti of deferoxamine and	111
	gentamicin group	
Figure 39	Semithin section of organ of corti of deferoxamine and gentamicin group	112
Figure 40	Ultrathin section of outer hair cells of deferoxamine and	113
	gentamicin group	
Figure 41	Ultrathin section of outer hair cell of deferoxamine and	114
	gentamicin group	L

Figure 42	Ultrathin section of outer hair cell of deferoxamine and	115
	gentamicin group	
Figure 43	Semithin sections to differentiate between organ of corti of	117
	The three different groups	
Figure 44	Ultrathin sections to differentiate between outer hair cells	119
	of the three different groups	