



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بلله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل



B9v19

Palaeontological and biometrical studies on some Late Cretaceous macroinvertebrate fossils of the Galala area, north Eastern Desert, Egypt

A Thesis

Submitted to the Geology Department, Aswan Faculty of Science
for **Ph.D.** degree in Geology

By

Mohamed Sayed Mohamed Ali Zakhera

B. Sc. in Geology (Assiut Univ.)

M.Sc. in Geology (Assiut Univ.)

Supervised by

Prof. Dr. Abbass I. Kenawy

Professor of Geology

Faculty of Science

Assiut University, Egypt

Prof. Dr. Ahmed S. Kassab

Professor of Geology

Faculty of Science

Assiut University, Egypt

Prof. Dr. T. Setoguchi

Professor of Palaeontology

Head of Geology Dept.

Faculty of Science

Kyoto University, Japan

Dr. Adel W. Felesteen

Lecturer of Geology

Aswan Faculty of Science

South Valley University

Egypt

South Valley University

1999

Medicine

1771

1

Medicine

1771

Dedication

*I dedicate this work to my parents
might ALLAH provides them with a
Healthy Life.*

And

*To my brother Dr. Mahmoud, hoping
a great success to him*



Acknowledgments

All acknowledgements and gratitude are due to **ALLAH** for blessing and support to crown this work by success.

I would like to express my deepest appreciation to Prof. Dr. Abbass Kenawy (Assiut University) for his kind supervision, encouragement and reading the manuscript.

I am greatly indebted to Prof. Dr. Ahmed Kassab (Assiut University) for his joint supervision, suggesting the points of research, guidance in the field, fruitful discussion and comments, and for critically revising the manuscript.

I am greatly indebted to Prof. Dr. Kiyotaka Chinzei Professor of Palaeontology and the Dean of Faculty of Science (Kyoto University, Japan) for his kindness, facilities in various ways, guidance in the laboratory, critical discussion, and for fruitful comments during my stay in Japan.

I wish to express my deepest appreciation to Prof. Dr. H. Setoguchi (Kyoto University, Japan) for his kind support, various facilities and supervision during my study in Kyoto University as Guest Researcher Associate in Doctor program supported by the Egyptian Government.

I am grateful to Dr. Adel W. Felesteen (Aswan Faculty of Science, South Valley University) for his sincere supervision, continuous encouragement, advice and help during the progress of this work.

My grateful thanks are due to Prof. Dr. M. A. El-Maghraby, the Dean of Aswan Faculty of Science, Prof. Dr. H. El Amin, head of the Geology Department and Vice Dean for student affairs and to Prof. Dr. A. I. Koraiem, Vice Dean for higher studies and researches for their unfailing encouragement, kind support, various facilities and help.

Deep thanks are also due to Prof. Dr. H. Maeda (Kyoto University) for his advice and help in making photographs and also to Dr. Keiko Yamaguchi (Kyoto University) for her kind assistance in using the Scanning Electron Microscope.

I am grateful to Prof. Dr. Hayami (Kyushu University, Japan), Prof. Dr. Bandel (Hamburg University, Germany), Prof. Dr. Togo (Hokkaido University, Japan) and to Dr. Malchus (Barcelona University, Spain) for written communication and valuable publications. I would like to express my thanks to Dr. Suzuki, Dr. Kamiya, Dr. Sakai, Mr. Komatsu, Mr. Tsujino, Mrs. Sato, Mrs. Nassan of Kyoto University for their help in various ways.

Thanks are also due to all of my colleagues in Geology Department, Aswan Faculty of Science and to the secretary staff of Geology and Mineralogy Department of Kyoto University for their help during the progress of this work.

Mohamed Zakhera

Notes

Some results of the present Thesis have been presented in an International Conference “the Annual Meeting of the Palaeontological Society of Japan”, whereas others have been published in the International Journal “Neues Jahrbuch für Geologie und Paläontologie”, as follows:

- 1) **Zakhera, M., Kassab, A. and Chinzei, K. (1997):** Morphology and shell microstructure of a Cenomanian oyster from Egypt. Annual Meeting of the Palaeontological Society of Japan, Kyoto University, 29/1/97 – 1/2/97, Extended Abstract, p. 85.

It presents the most important results of studying shell microstructure and functional morphology of some oysters identified in the Thesis.

- 2) **Zakhera, M. and Kassab, A. (1999):** New species of ostreid bivalves from the Upper Cretaceous (Santonian-Maastrichtian) of Egypt.— N. Jb. Geol. Paläont. Mh., 1999 (3), 12 pp., Stuttgart.

It includes the new oyster species as a part of the systematic palaeontology of the bivalves identified in the Thesis.

Contents

	Page
List of Figures.....	iii
List of Tables.....	v
Abstract.....	vi
Chapter One	1
Introduction.....	1
I- Geographic setting.....	1
II- Historical Review.....	1
III- Scope of the Present Study.....	13
IV-Material& Methods.....	14
Chapter Two	17
Lithostratigraphy.....	17
- Galala Formation	17
- Umm Omeiyid Formation.....	19
- Hawashiya Formation.....	20
- Duwi Formation.	22
- Sudr Formation.	23
- Gebel Thilmet Formation.	25
Chapter Three	36
Biostratigraphy	36
-The proposed biozones for Upper Cretaceous	36
- Age assignment.	45
Chapter Four	56
Shell Microstructure	56
- Introduction	56
- Material & Methods.....	57
- Microstructural fabrics	57
- Results of the present study	60
Chapter Five	71
Functional Studies.....	71
- Introduction.....	71
- Morphological and Microstructural analyses.....	72
- Functional interpretation and Inferred mode of life....	73
- Conclusions.....	74

Chapter Six	79
Biometry.....	79
- Introduction.....	79
- Material and Methods.....	80
- Results and interpretation.....	81
- Size frequency distribution.....	81
- Basic statistics.....	86
- Analysis of Variance (ANOVA).....	88
- Bivariate analysis.....	89
- Correlation coefficient.....	93
- Cluster analysis.....	93
- Discussion and conclusions.....	96
Chapter Seven	98
Systematic Palaeontology	98
-Plates	187
Chapter Eight	221
Discussion and Conclusions	221
References	235
Arabic Summary	

List of Figures

Figure	Page
(1): Location map.....	2
(2): Field views of the Upper Cretaceous successions exposed at the study area.....	28
(3): Field views of certain litho-and bio-facies details of the Upper Cretaceous Formations, exposed in the study area.....	30
(4): Field views, showing the characteristic features of the Gebel Thilmet Formation in the study area.	32
(5): Field views of certain features and outcrops of the Upper Cretaceous Formations in the study area.	34
(6): Lithologic symbols used in the present study.	46
(7): Vertical distribution of the identified bivalves and the biozones, Abu Darag section.	47
(8): Vertical distribution of the identified bivalves and the biozones, Gebel Thilmet section.	48
(9): Vertical distribution of the identified bivalves and the biozones, Saint Anthony section.	49
(10): Vertical distribution of the identified bivalves and the biozones, Saint Paul section.	50
(11): Vertical distribution of the identified bivalves and the biozones, Wadi El- Dakhal section.	51
(12): Vertical distribution of the identified bivalves and the biozones, Wadi Tarfa section.	52
(13): Sketch showing the microstructural fabrics of oyster shell.	66
(14): Microstructural types of the examined oyster.	67
(15): Microstructural types of the examined oyster.	69
(16): Two morphotypesw of <i>Ilymatogyra africana</i> (Lmk.) (a) and <i>Rhynchostreon mermeti</i> (Coq.) (b).	71
(17): Sketch showing the degree of coiling umbo of <i>Ilymatogyra africana</i> (Lmk.) and <i>Rhynchostreon</i> <i>mermeti</i> (Coq.).	76
(18): Radial vertical section through right valve of <i>Ilymatogyra africana</i> (Lmk.), showing the Lenticular chambers and the chalky deposits.	76

(19): Radial vertical section in shell wall of <i>Ilymatogyra africana</i> (Lmk.) (a) and <i>Rhynchostreon mermeti</i> (Coq.) (b).	77
(20): The inferred mode of life of <i>Ilymatogyra africana</i> (Lmk.).	78
(21): The inferred mode of life of <i>Rhynchostreon mermeti</i> (Coq.).	78
(22): The morphometric characters measured for <i>Ceratostreon flabellatum</i> (Goldfuss).	81
(23): Size-frequency histograms of some morphometric characters of <i>Ceratostreon flabellatum</i> (Goldfuss).	82
(24): Size-frequency histograms of other morphometric characters of <i>Ceratostreon flabellatum</i> (Goldfuss).	83
(25): Bivariate distribution of L, C, Hu, Lu, Ham, Lam of <i>Ceratostreon flabellatum</i> (Goldfuss).	91
(26): Bivariate distribution of A-H, A-P, GB, Uc, Cu, Cu/H of <i>Ceratostreon flabellatum</i> (Goldfuss).	92
(27): Cluster dendrogram of the measured morphometric characters of <i>Ceratostreon flabellatum</i> (Goldfuss).	94
(28): Cluster dendrogram of the analyzed individuals of <i>Ceratostreon flabellatum</i> (Goldfuss).	95
(29): Distribution and correlation of the fossiliferous Upper Cretaceous rock units exposed in the study area.	232
(30): Cenomanian- Santonian composite section of the Galala area with the proposed bivalve zones.	233
(31): Santonian-Maastrichtian composite section of the Galala area with the proposed bivalve zones.	234