The Early Detection of Subclinical Synovitis by High Frequency Power Doppler Ultrasonography in oligoarticular Juvenile Idiopathic Arthritis

Ehesis

Submitted for partial fulfillment of Master Degree in **Physical Medicine**, **Rheumatology and Rehabilitaion**

BySally Mohamed Abd Elhady Sayed

M.B., B.Ch. Faculty of Medicine, Ain Shams University

Under Supervision of

Prof. Dr. Hala Abd Elhady Saleh Hassan

Professor of Physical Medicine, Rheumatology and Rehabilitation Faculty of Medicine – Ain Shams University

Prof. Dr. Amal Mostafa Ibrahim El Ganzouri

Professor of Physical Medicine, Rheumatology and Rehabilitation Faculty of Medicine – Ain Shams University

Dr. Sahar Fathi Ahmed

Assistant Professor of Physical Medicine, Rheumatology and Rehabilitation Faculty of Medicine – Ain Shams University

Faculty of Medicine
Ain Shams University

2014





First, and foremost, my deepest gratitude and thanks should be offered to "ALLAH", the Most Kind and Most Merciful, for giving me the strength to complete this work.

I would like to express my sincere gratitude to **Prof. Dr. Hala Abd Elhady Saleh Hassan,** Professor of Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine – Ain Shams University for her continuous support and guidance for me to present this work. I really have the honor to work under her generous supervision.

I acknowledge with much gratitude to **Prof. Dr. Amal Mostafa Ibrahim El Ganzouri**, Professor of Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine – Ain Shams University, for her great supervision and unlimited help to provide all facilities to accomplish this work.

I extend my appreciation and thanks to **Dr. Sahar Fathi Ahmed**, Assistant Professor of Physical Medicine, Rheumatology and Rehabilitation, Faculty of Medicine – Ain Shams University, for her great efforts, cooperation, valuable advice, help and encouragement.

I can't forget to thank **Dr. Hossam Mousa Sakr**, Assistant Professor of Radiology, Faculty of Medicine – Ain Shams University, for his great efforts and cooperation in the current work.

Last but not least, thanks to my Parents and my Family for helping me to finish this work.

A Sally Mohamed A. El-Hady

List of Contents

Subject Page S	No.
List of Abbreviations	i
List of Tables	.vii
List of Figures	xiii
Introduction	1
Aim of the Study	5
Review of Literature	••••
Juvenile Idiopathic Arthritis	6
Epidemiology of JIA	.12
Pathogenesis of JIA	.31
Pathology of JIA	.64
Classifications of JIA	.71
Clinical picture of JIA	.76
Investigation of JIA	115
Treatment of JIA	127
Musculoskeletal Ultrasonography	150
Musculoskeletal ultrasonography in juvenile idiopathic arthritis	160
Patients and methods	200
Results	220
Discussion	288
Summary	313
Conclusion	319
Recommendations	320
References	321
Arabic Summary	

List of Abbreviations

A/AT : Achilles tendon

ACPA : antibodies against citrullinated proteins
ACR : American College of Rheumatology

Act : Active

ADAD : Adenosine deaminase domain

ADAM : A Disintegrin And Metalloproteinase

ANA : Antinuclear antibodies

ANGPT1 : Angiopoietin-1

anti-CCP : Antibodies against cyclic citrullinated peptideanti-MCV : Antibodies against mutated citrullinated vimentin

AS : Ankylosing spondylitis **ATA** : Aurintricarboxylic acid

BMI : Body mass index

BOOP : Bronchiolitis obliterans organizing pneumonia

C. elegans: Caenorhabditis elegans

C : Complement C : Cytosine C/Cal : Calcaneum

CACP : Camptodactyly-Arthropathy-Coxa Vara-Pericarditis

CBC : Complete blood countCCR/CXCR : Chemokine receptorsCD : Cluster of differentiation

CePD : Contrast-enhanced Power Doppler

CeUS : Contrast-enhanced US

CHAQ : Childhood health assessment questionnaire

Clin : Clinical

COG : Conserved oligomeric golgiCR : Conventional radiography

CRP : C-reactive protein

CT : Computed tomography

CTLA : Cytotoxic T-lymphocyte antigen

D : Diversity regionD : DimensionalDD : Dijkstra damageDG : Dijkstra growth

DHEA : Dehydroepiandrosterone

DHEA-S: Sulfated conjugate of dehydroepiandrosterone

DI : Dijkstra inflammation

DMARDs: Disease modifying antirheumatic drugs

dnaJ : Deoxy-ribonucleic acid

E : Epiphysis

EBV : Epstein Barr virus

Eff : Effusion

EFV : Extended field of view **ERA** : Enthesitis-related arthritis

ESR : Erythrocyte sedimentation rate

EULAR : European League Against Rheumatism

FE : Femoral epiphysis
FOXP : Forkhead box P
GroEL : Bacterial chaperonin
GT : Greater tuberosity

GWAS : Genome wide association study

Hb : Haemoglobin

HDL : High density lipoproteinHLA : Human leukocyte antigen

HMBPP : (E)-4-hydroxy-3-methyl-2-butenyl pyrophosphate

HMG : High mobility group

hr : Hour

HS : Highly significanthsp : Heat shock protein

IAC : Intraarticular corticosteroid

ICU : Intensive care unitIFN-γ : Gamma interferon

Ig : Immunoglobulin

IL : Interleukin

ILAR : International League of Associations for Rheumatology

ILGF: Insulin-like growth factor

Inact: Inactive

IPP : Isopentenyl pyrophosphateIRF : Interferon regulatory factor

J : Joining region

JAFS : Juvenile Arthritis Functionality Scale

Jak : Janus kinases

JAS : Juvenile ankylosing spondylitis

JCA : Juvenile Chronic Arthritis

JIA : Juvenile idiopathic arthritis

JPsA : Juvenile Phoymatoid Arthritis

JRA : Juvenile Rheumatoid Arthritis JSpA : Juvenile-onset spondylarthritis

K : Kager's fat padk : Kappa agreement

m : Month/s

MAS : Macrophage activation syndrome

MCP : Metacarpophalangeal

MD global: Physician global assessment of overall disease activity

MHC : Major histocompatibility class

MICA/MICB: MHC class I chain related gene A/B
MIF: Macrophage migration inhibitory factor.

ML : Monoclonal antibodies
 MMPs : Matrix metalloproteinases
 MRI : Magnetic resonance imaging
 mRNA : Messenger ribonucleic acid
 MRP : Myeloid related proteins
 MSUS : Musculoskeletal ultrasound

MTP : Metatarsophalangeal

MTX : Methotrexate Neg : Negative NK : natural killer

No : Number of patients/joints

NS : Non significant

NSAIDs : Nonsteroidal anti-inflammatory drugs

OC : Ossification center
Oligo : Oligoarticular

OMERACT: Outcome Measures in Rheumatology Clinical Trial

P : Level of significance

P : Patella

PAF-AH: Platelet activating factor acetylhydrolase

PB : Peripheral blood

PCR : Polymerase chain reaction

PD : Power Doppler

PIP : Proximal interphalangeal

Pos : Positive

PsA : Psoriatic arthritis

PSRA : Post-streptococcal reactive arthritis

PT : Prothrombin time

PTPN: Protein tyrosine phosphatase non receptor **type**

PTT : Partial thromboplastin time

r : Correlation coefficient RA : Rheumatoid arthritis

RAND/UCLA: Research and Development/University of California at Los Angeles

RANKL: Receptor activator of nuclear factor-kappa B ligand

ref. no : Reference number
REL : Reticuloendotheliosis
RF : Rheumatoid factor
ROM : Range of motion

S : Significant S./ Sig : Significance

SCID : Severe combined immune deficiency

SD : Standard deviation SF : Synovial fluid

SJIA : Systemic juvenile idiopathic arthritis

SLE : Systemic lupus erythematosusSNPs : Single nucleotide polymorphismSONAR : Sound navigation and ranging

SpA : Spondylarthritis

STAT : Signal transducer and activator of transcription

T : Tendon

t : Paired t test

TCRs : T cell receptorsTE : Tibial epiphysisTFP : Task Force Panel

TGF : Transforming growth factor

Th : T helper cell

TIE : Endothelium specific receptor tyrosine kinase

TLR : Toll-like receptor

TMJ : Temporo-mandibular joint

TNF : Tumour necrosis factor

TNFAIP: TNF alpha induced protein

TRAF : TNF-receptor association factor

Tregs : T regulatory cells
U test : Mann Whitney Test
UK : United Kingdom

US : Ultraound, ultrasonographyUSA : United States of America

V : Variable region

VAS : Visual analog scale

VLA : Very late activation antigen

WBCs : White blood cells

wk : Week
yr : Year

λs : Sibling recurrence risk

* : Fisher's Exact Test

List of Tables

Eable N	o. Citle	Page No.
Table (1):	Studies of the incidence and prevalence o childhood arthritis	
Table (2):	Criteria for the classification of rheumatoid arthritis	
Table (3):	Criteria for a diagnosis of juvenile chronic	arthritis73
Table (4):	Comparison of EULAR, ACR and ILAR for classification of chronic arthritis of chi	
Table (5):	Systemic juvenile idiopathic arthritis: Inte League of Association for Rheumatology diagnostic criteria	(ILAR)
Table (6):	American Academy of Pediatrics Guide Screening Eye Examinations	
Table (7):	Descriptive data for the age	220
Table (8):	Descriptive data for the gender	220
Table (9):	Descriptive data for disease duration, joints in 1st six months and along the course of the	
Table (10):	Frequency of types of oligoarticular JIA	221
Table (11):	Descriptive data for the medication history	y222
Table (12):	Frequency of the past history of red eye	222
Table (13):	Descriptive data for family history	223
Table (14):	Evaluation of weight, height and body mas	ss index223
Table (15):	Number of knees and ankles involved in o	ur study223
Table (16):	Clinical parameters among knees and ankl	es224
Table (17):	Evaluation of articular index scores	225
Table (18):	Number and percentage of articular index	scores 226
Table (19):	Number and percentage of total articul score.	

Cable N	o. Gitle Page No.
Table (20):	Number and percentage of clinical active and inactive knees and ankles
Table (21):	Description of JAFS score
Table (22):	Number and percentage of modified JAFS score228
Table (23):	Description of laboratory parameters
Table (24):	Number of active joints in each patient230
Table (25):	Description of physician's VAS and parent's VAS230
Table (26):	Evaluation of disease activity score231
Table (27):	Number and percentage of patients in each disease activity score
Table (28):	Number and percentage of plain X- ray using modified Larsen scoring
Table (29):	Number and percentage of plain X- ray using Dijkstra composite scoring
Table (30):	Number and percentage of ultrasonography parameters among knees and ankles234
Table (31):	Description of ultrasonographic scores
Table (32):	Number and percentage of joints as regards ultrasonographic scores
Table (33):	Number and percentage of ultrasonography active and inactive knees and ankles
Table (34):	Description of extra US findings
Table (35):	Comparison between clinical activity and US active joints
Table (36):	Comparison between clinical activity and US inactive joints238

Table N	o. Eitle	Page No.
Table (37):	Number and percentage of knees and clinical inactive and US active joints	
Table (38):	Number and percentage of knees and clinical active and US inactive joints	
Table (39):	Comparison between tenderness score activity of joints.	
Table (40):	Comparison between swelling score activity of joints.	
Table (41):	Comparison between limited ROM scor activity of joints.	
Table (42):	Comparison between total severity score activity of joints.	
Table (43):	Comparison between US synovial hy score and clinical activity of joints	• •
Table (44):	Comparison between US effusion score are activity of joints.	
Table (45):	Comparison between US PD score and activity of joints.	
Table (46):	Correlation between clinical scores and U	JS scores244
Table (47):	Agreement between US synovial hyperpclinical examination as regards swelling.	
Table (48):	Agreement between US effusion and examination as regards swelling	
Table (49):	Agreement between US PD and clinica joints	
Table (50):	Agreement between US active joints and active joints	~

Cable N	o. Eitle	Page No.
Table (51):	Correlation between ESR and CRP with score.	
Table (52):	Comparison between modified Larsen US activity of joints.	
Table (53):	Comparison between DI score and US a joints	•
Table (54):	Comparison between DD score and US a joints	-
Table (55):	Comparison of DG score and US activity	of joints256
Table (56):	Comparison between total Dijkstra of score and US activity of joints	•
Table (57):	Correlation between modified Larsen stotal Dijkstra composite score with total U	
Table (58):	Comparison between US activity and activity.	
Table (59):	Comparison of means of tenderness, limited ROM and total articular index so 2 months	cores after
Table (60):	Comparison of tenderness, swelling an ROM scores after two months	
Table (61):	Comparison of total articular index score months	
Table (62):	Comparison of the mean of modified J. two months.	
Table (63):	Comparison of the mean of laboratory p after 2 months.	
Table (64):	Comparison of the mean of physical parent's VAS after two months	

Cable N	o. Citle	Page No.
Table (65):	Comparison of the mean of disease actival after two months.	-
Table (66):	Comparison of disease activity score a months.	
Table (67):	Comparison of means of plain X ray scor months	
Table (68):	Comparison of plain X- ray using modifies score after two months.	
Table (69):	Comparison of plain X- ray using composite score after two months	
Table (70):	Comparison of means of synovial hyperfusion and PD scores after 2 months	-
Table (71):	Comparison of US parameters after two m	onths269
Table (72):	Comparison between tenderness score activity of joints after two months	
Table (73):	Comparison between swelling score activity of joints after two months	
Table (74):	Comparison between limited ROM score activity of joints after two months	
Table (75):	Comparison between total articular ind and US activity of joints after two months	
Table (76):	Comparison between US synovial hy score and clinical activity of joints a months	after two
Table (77):	Comparison between US effusion so clinical activity of joints after two months	
Table (78):	Comparison between US PD score and activity of joints after two months	