



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

ULTRASONIC COMPARISON BETWEEN DIFFERENT OVULATION INDUCTION PROTOCOLS

THESIS

Submitted for Partial Fulfillment for
Master Degree in Obstetrics and Gynaecology



By

Suzan William Mansour

M.B, B.Ch. Cairo University

Resident of Obstetrics and Gynaecology

Helwan University Hospital

Supervised By

Prof. Aly Mahmoud El-Gazar

Professor of Obstetrics and Gynaecology

Prof. Nabil Gamal El-Orabi

Professor of Obstetrics and Gynaecology

Dr. Mohamed Kamel Alloush

Assistant Professor of Obstetrics and Gynaecology

Benha Faculty of Medicine

Zagazig University

1999

B 7192

ACKNOWLEDGMENT

First and foremost, I feel always indebted to God the most kind and most merciful.

I have no words by which I could express my deepest gratitude, extreme appreciation and sincere thanks to Professor Aly Mahmoud El-Gazar, Professor of Obstetrics and Gynaecology, Professor Nabil Gamal El-Orabi, Professor of Obstetrics and Gynaecology, and Dr. Mohamed Kamel Alloush, Assistant Professor of Obstetrics and Gynaecology, Benha Faculty of Medicine Zagazig University, I would never forget their valuable guidance, precious observations, continuous support and supervision throughout the work.

My deepest thankful gratitude to all staff members of Obstetrics and Gynaecology department in Benha Faculty of Medicine Zagazig University for their kind support throughout this work.

Suzan William Mansour

CONTENTS

INTRODUCTION	5
AIM OF THE WORK	8
REVIEW OF LITERATURE	
• FOLLICLE GROWTH AND DEVELOPMENT	10
• DISORDERS OF OVULATION	27
• DIAGNOSIS OF ANOVULATORY INFERTILITY	45
• ULTRASONOGRAPHY DETECTION OF OVULATION	60
• METHODS OF INDUCTION OF OVULATION	67
• COMPLICATIONS OF OVARIAN STIMULATION	87
SUBJECTS AND METHODS	107
RESULTS	114
DISCUSSION	128
SUMMARY	133
CONCLUSION	140
REFERENCES	142
ARABIC SUMMARY	166

LIST OF FIGURES

Figure 1	12
Figure 2	34
Figure 3	35
Figure 4	36
Figure 5, 6	50
Figure 7	52
Figure I	121
Figure II	122
Figure III	123
Figure 8	124
Figure 9	125
Figure 10	126
Figure 11	127

	INTRODUCTION	
--	-----------------------	--

INTRODUCTION

Induction of ovulation whether medical or surgical is considered to be a management for chronic anovulatory state. Chronic anovulation is a major cause of infertility and in most conditions it is possible to restore normal fertility by careful selection of the appropriate method of induction of ovulation (Franks et al; 1994).

Clomiphene citrate has been available for induction of ovulation for nearly 30 years and still has an important place in the management of women with an ovulatory menses and of estrogenized subjects with amenorrhea. Most women with amenorrhea with a positive progesterone withdrawal test have clinical, biochemical and ultrasound evidence of polycystic ovarian syndrome (Fox et al; 1991).

Gonadotropin-releasing hormone remains the treatment of choice for induction in women with hypogonadotropic hypogonadism of hypothalamic origin (Martin et al; 1993).

Gonadotropin-releasing hormone analogs with gonadotropine in polycystic ovarian syndrome. PCOS is typically associated with hypersecretion of LH and there are now several studies which have shown a link between persistently elevated levels of concentration of LH and a poor outcome of induction of ovulation, so it seems logic to suppress of endogenous LH by administration of GnRH agonist analogs during gonadotropine induced ovarian stimulation may improve the rate of successful pregnancy (Homburg et al; 1990).

It has now been reported that there is a reduced rate of multiple follicular development of purified FSH in very low doses (Polson et al; 1987).

A recent report using human endometrial stromal cells in vitro indicates that FSH, LH and HCG extracted from the urine of either post menopausal women or pregnant women stimulate intracellular Cyclic AMP and induced decidualization (Tang et al; 1993).

Paradoxically, little is known about the pharmaco-dynamics and pharmaco-kinetics of LH and FSH during such stimulation.

Recently, the existence was shown of a sharp FSH threshold concentration above which the ovary responds with gonadotrophins in hypogonadotrophic women, this FSH threshold concentration was independent of the co-administration (Van Weissenbruch; 1990).

	AIM OF THE WORK	
--	-------------------------------	--

AIM OF THE WORK

The aim and the objective of study was done to select the best methods of induction for patients with anovulation variable: PCO, hypothalamic hypogonadotropic hypogonadism, and hyperprolactinamic patients denoted by transvaginal sonography.