

Management of fractures of distal radius

Thesis

Submitted for partial fulfillment of

Master degree in orthopedic surge

Ahmed Ibrahim Fathalla Ibrahim

M.B.B.CH

Faculty of medicine

Al-Azhar university

SUPERVISED BY

PROF.DR.IBRAHIM AHMED HUSSEIN

Professor of orthopedic surgery

Faculty of medicine

Al-Azhar university

PROF.DR.AHMED IBRAHIM KHAMIS.

Ass.Prof.of Orthopedic surgery

Faculty of medicine

Al-Azhar university

Faculty of medicine

Al-Azhar university

Cairo 2014

Acknowledgement

Praise to "**Allah**", the most gracious and the most merciful who guides us to the right way.

I would like to express my deepest gratitude and thanks to **Professor Dr. Ibrahim Hussien**, Professor of Orthopaedic Surgery and Traumatology, Faculty of Medicine, Alazher University for his meticulous guidance, constructive criticism and encouragement were indispensable in accomplishing this thesis.

My extreme appreciation to **Dr. Ahmed Khamis**, Assistant Professor of Orthopaedic Surgery and Traumatology, Faculty of Medicine, Alazher University for his willing help and critical review. His generous advice was illuminative throughout the work.

I would like to thank **all members** of the Department of Orthopaedic Surgery and Traumatology, Faculty of Medicine Alazher University for their support and cooperation.

I am forever grateful to **my family** for their ever after love and support.

Contents

1. Introduction .

2. Aim of the work.

3. Review of literature:

- **Anatomy.**
- **Radiological anatomy**
- **Biomechanics**
- **Mechanism of injury and Classification**
- **Management:**
 - . **Diagnosis**
 - . **Different methods of fixation**
- **Complication**

4. Case presentation.

5. Discussion.

6. Summary.

7. References.

8. Arabic summary.

List of Figures

<i>Figures</i>		<i>Page</i>
Figure 1	View of the articular surfaces of the distal radius and ulna.	6
Figure 2	: A lateral view from the ulnar side shows the plane defined by the volar surface of the distal radius and the extent of the lunate facet anterior to this plane.	7
Figure 3	: Anatomical landmarks of the volar surface of the distal radius.	7
Figure 4	: The palmar vascular anatomy includes a series of two transverse arches that traverse the radial and ulnar arteries.	12
Figure 5	: Cross section at the wrist level illustrating the basic anatomic relations of major structures.	13
Figure 6	: Volar wrist extrinsic and intrinsic ligaments.	15
Figure 7	: Anatomy of the TFCC. (A) End view of the DRUJ. (B) Coronal view of the DRUJ	16
Figure 8	: Wrist joint: Capsular attachment and synovial lining.	18
Figure 9	: PA x-ray of a wrist	20
Figure 10	: Measurement of ulnar variance	21
Figure 11	: Radial width PA x-ray of a wrist	22
Figure 12	: Diagram showing the palmar inclination of the distal radius.	23
Figure 13	: measurement of the AP distance	24
Figure 14	: Assessing volar and dorsal rim on PA view.	25
Figure 15	: Three-dimensional computed tomography scans of a distal radial fracture measurement of the AP distance	28
Figure 16	: Ulnar variance is measured on a standardized PA radiograph.	34

Figure 17	: Tension patterns in the dorsal radioulnar ligament (DRUL) and the palmar radioulnar ligament (PRUL).	37
Figure 18	: Frykman classification	43
Figure 19	: Fernandez classification	47
Figure 20	: AO Classification.	51
Figure 21	: Schematic representation of the three-point contact cast.	62
Figure 22	: Numerous percutaneous pinning techniques	66
Figure 23	: Plain radiograph and a photo of a patient show the nonbridging wrist fixator.	73
Figure 24	: Plain radiograph showing AO external fixator fixing fracture of the distal end radius	74
Figure 25	: Dynamic wrist Fixator	75
Figure 26	: Plain radiograph showing the radiolucent wrist fixator in combination with k wire fixation.	76
Figure 27	: Plain radiograph and photo of a patient showing the hybrid wrist fixator.	77
Figure 28	: A. Intramedullary nail B.X-ray of implanted intramedullary nail.	80
Figure 29	The three-column concept of the structural anatomy of the distal radius and ulna.	84
Figure 30	: Fragment-specific implants.	84
Figure 31	: Volar non-locking T-plate "Synthes".	91
Figure 32	: Force distribution of a plate osteosynthesis without angular stability.	93
Figure 33	: Locking compression plate with combination hole.	94
Figure 34	: The Hand Innovations DVR plate using 2.0-mm distal locking buttress pins with terminal thread and 3.5-mm non-locking shaft screws.	98

Figure 35	: T-shaped LCP (Mathys Medical Ltd, Bettlach, Switzerland)A volar locking plate with combi-holes and a locking screw.	99
Figure 36	: Second generation DVR plate.	99
Figure 37	: Early osteotomy and use a volar fixed angle plate to correct dorsal	113
Figure 38	: angulation and radial inclination	114
Figure 39	: pre-operative AP and lateral X-RAYS	115
Figure 40	: CT view	115
Figure 41	: post-operative X-RAYS	116
Figure 42	: 2months after removal fixation126	116
Figure 43	: Hand movement with fixation	117
Figure 44	: Early osteotomy and use a volar fixed angle plate to correct dorsal	118
Figure 45	: angulation and radial inclination	118
Figure 46	: pre-operative AP and lateral X-RAYS	119
Figure 47	: CT view	119
Figure 48	: post-operative X-RAYS	120
Figure 49	: 2months after removal fixation126	121
Figure 50	: Hand movement with fixation	121
Figure 51	:Immediate postoperative x-rays	122
Figure 52	: Three weeks postoperative.	122
Figure 53	: One month following k-wires and fixator removal	123

Figure 54	: Pre-operative AP and lateral X-RAYS	124
Figure 55	: Immediate postoperative x-rays	124
Figure 56	: After 5 months post- operative AP and lateral X-RAYS	125
Figure 57	: Hand movement after physiotherapy	125
Figure 58	: Pre-operative AP and lateral X-RAYS	126
Figure 59	: Immediate postoperative x-rays	126
Figure 60	: Post-operative APand lateral X-RAYS after 4 months of operation	127
Figure 61	: Scar of incision	127
Figure 62	: Hand movement after physiotherapy	128

List of tables

Table 1 : Classification of BINDRA for comminuted intra-articular fractures of distal Radius.

Table 2 : FRYKMAN classification of fracture of distal radius.

Table 3 : MCMURTRY AND JUPITER classification.

Table 4 : MELONE classification.

Table 5 : Main different points between volar fixed-angle locked plating and external fixation with percutaneous wiring .

List of abbreviations

AO	Arbeitsgemeinschaft fur Osteosynthesefragen
AP	Anteroposterior
BR	Brachioradialis
CRPS	Complex regional pain syndrome
DRF	Distal radial fractures
DRUJ	Distal radioulnar joint
EPL	Extensor Pollicis Longus
FCR	Flexor carpi radialis
FFH	Fall from a height
FPL	Flexor pollicis longus
IFZ	Intermediate fibrous zone
LCP	Locking compression plate
LHS	Locking head screw
ORIF	Open reduction and internal fixation
PMMA	Polymethyl methacrylate
PQ	Pronator quadratus
RSC	Radioscaphocapitate
SRL	Short radiolunate
TCL	Transverse carpal ligament
TFCC	Triangular fibrocartilage complex

Introduction

Distal radius fractures remain an injury that fosters considerable interest and debate.⁽¹⁾ Distal radial fractures account for 20% of fractures among medicare enrollees and are second only to hip fractures in terms of incidence in the elderly population. These fractures represent approximately one-sixth of all fractures that are treated in emergency departments.⁽²⁾

In the older population, the mechanism of injury usually involves a fall into the outstretched hand in an elderly patient whose bone quality is diminished by osteoporosis. In younger patients, high-energy injury mechanisms lead to wide displacement and marked comminution in bone of normal quality.⁽³⁾

The goal of surgical fixation in the unstable distal radius fracture is to restore intra-articular and extraarticular anatomic alignment.⁽⁴⁾ Persistent step or gap deformity of the articular surface after distal radius fracture may predispose to the development of posttraumatic arthritis of the radiocarpal joint. Although the clinical manifestations of post traumatic arthritis may not be disabling, yet the goal of surgical treatment of intra-articular distal radius fractures remains anatomic reduction.⁽⁵⁾

Many methods are available to treat distal radius fractures, each with its own benefits and potential complications, which must be weighed when choosing which procedure to use.⁽⁶⁾

Many fixation techniques have been described including pin and plaster fixation, percutaneous pinning, intramedullary pinning, external fixation (bridging or nonbridging, static or dynamic), injectable bone cement, and various forms of internal fixation with customized implants.⁽⁷⁾

AIM OF THE WORK

The aim this prospective study to evaluate different types of fracture distal radius in different ages with different methods of treatment and follow up of these cases .

Anatomy

A) Osseous anatomy

The lower end of the radius is large, quadrilateral, and provided with two articular surfaces; one below, for the carpus, and another at the medial side, for the ulna. The carpal articular surface is triangular, concave, smooth, and divided by a slight antero-posterior ridge into two parts. The lateral, triangular, articulating with the scaphoid bone and called the scaphoid fossa; the medial, quadrilateral, with the lunate bone and called the lunate fossa. The articular surface for the head of the ulna is called the ulnar notch (sigmoid cavity) of the radius. These two articular surfaces are separated by a prominent ridge, to which the base of the triangular articular disk is attached; this disk separates the wrist-joint from the distal radioulnar articulation (**Figure1**).

The distal radius has three non-articular surfaces: volar, dorsal, and lateral. The dorsal surface is convex, affords attachment to the dorsal radiocarpal ligaments, and is marked by three grooves and a prominent ridge "Lister's tubercle". The lateral surface is prolonged obliquely downward into the styloid process, which gives attachment by its base to the tendon of Brachioradialis, and by its apex to the radial collateral ligament of

the wrist-joint. The lateral surface of this process is marked by a flat groove, for the tendons of the Abductor pollicis longus and Extensor pollicis brevis.⁽⁸⁾

The volar surface, is broad, flat, affords attachment to the volar radiocarpal ligaments and the pronator quadratus. The volar lunate facet projects an average of approximately 3mm anterior to the relatively flat volar surface (**Figure 2**).⁽⁹⁾ Plates placed over this prominence can risk tendon irritation and can be more palpable to the patient.⁽¹⁰⁾

The concave surface of the volar radius is limited distally by a transverse ridge or the **watershed line**. Distal to the watershed line, the radius slopes in a dorso-distal direction and receives the proximal attachments of the volar wrist capsule and the volar capsular ligaments. This ridge lies close (2 mm) to the joint line on its ulnar aspect and well proximal (10–15 mm) from the joint line on its radial aspect. Fixation implants must be placed proximal to and their profile must not project a **Volar Radial Tuberosity** is a prominence on the radial margin of the pronator quadratus (PQ) fossa. Plates that are placed too radial will overlie this tuberosity, will be pronated, and will not lie flat on the radius and may become palpable to the patient, requiring subsequent

removal. **Volar Radial Ridge** a ridge of varying prominence extends proximally from the volar radial tuberosity and marks the radial limit of the pronator quadratus fossa (**Figure 3**).⁽¹⁰⁾




Figure 1: View of the articular surfaces of the distal radius and ulna.⁽¹¹⁾