

شبكة المعلومات الجامعية







شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد أعدت دون أية تغيرات




يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار المفلام بعيدا عن الغبار %٤٠-٢٠ منوية ورطوية نسبية من ٢٠-١٠ في درجة حرارة من ٢٥-١٥ منوية ورطوية نسبية من ٢٥-١٥ to be Kept away from Dust in Dry Cool place of 15-25- c and relative humidity 20-40%











EFFECT OF GIBBERELLIC ACID AND SOME **BIOFERTILIZERS APPLICATION ON YIELD AND** FRUIT CHARACTERISTICS AT HARVEST OF EGYPTIAN LIME (CITRUS AURANTIFOLIA B.)

BY Mohamed Ahmed Hussein Hassan

B.Sc. Agric. Sci., (Pomology, Assiut University (1998)

Thesis

Submitted in Partial Fulfillment of the Requirement For the degree of

MASTER OF SCIENCE

IN

Agricultural Sciences (Horticulture Pomology) **Department of Horticulture Faculty of Agriculture Assiut University**

2005

Supervised by:

Prof. Dr. Hamdy M.M. Marzouk Dr. Farouk M. A. Mostafa Hort. Dept., Assiut University

Hort. Dept., Assiut University

CRYJJY



APPROVAL SHEET

The members of the committee appointed to examine the dissertation presented by:

Mohamed Ahmed Hussein Hassan

Title: Effect of gibberellic acid and some biofertilizers application on yield and fruits characteristics at harvest of Egyptian lime (Citrus aurantifolia B.)

found it satisfactory and have been approved for M.Sc. degree in Horticultural Science (Pomology)

Prof. Dr. Hamdy M.M. Marzouk Handy M. M. Massgorth Hort. Dept. Assiut. Univ.

Prof. Dr. Samir Z. El-Agamy Hort. Dept. Assiut.

Prof. Dr. Farouk H. Abd El-Aziz . F. H. . Abdela 2.1.2.....
Hort. Dept. Minia. Univ.

Prof. Dr. Farouk M. A. Mostafa Hort. Dept. Assiut. Univ. Farouk Mostafa

Committee in charge

Date: 3º/ 7/2005

ACKNOWLEDGEMENT


ACKNOWLEDGMENT

I would like to express my deepest grateful and appreciation to *Prof. Dr. Hamdy M.M. Marzouk*. Professor of Pomology, Horticulture Department, Faculty of Agriculture, Assiut University, for his supervision, valuable criticism, providing facilities, technical advice and preparing this manuscript.

With sincere respect and gratitude, I would like to express my deep thanks to *Dr. Farouk M. A. Mostafa El-Borai*, Associate Professor of Pomology, Horticulture Department, Faculty of Agriculture, Assiut University, for his supervision, encouragement, Counsel in carrying out the research and in preparation of this manuscript.

I am greatly indebted to *Prof. Dr. Samir Z. El-Agamy*, *Prof. Dr. Abd El Fattah M. El-Salhy*, *Prof. Dr. Ahmed Mokhles Abdo*, *Associate Prof. Dr. Talat K. El-Mahady*, *Dr. Hassan A. Abd El-Galiel*, Associate Prof. of Pomology, staff members of Horticulture Department, Faculty of Agriculture, Assiut University, for kind advice, help, support and their encouragement.

Mohamed Ahmed Hussien



Contents	Page
1. Introduction	1
2. Review of literature	3
2.1. Effect of some biofertilizers and GA ₃ on yield components of some citrus fruit trees	3
2.2. Effect of some biofertillizers and GA ₃ acid on fruit number/tree of some citrus fruit trees	7
2.3. Effect of some biofertilizers and GA ₃ on fruit weight of some citrus fruit trees	10
2.4. Effect of some biofertilizers and GA ₃ on fruit volume of some citrus fruit trees:	13
2.5. Effect of some biofertilizers and GA ₃ on fruit length and diameter of some citrus fruit trees	15
2.6. Effect of some biofertilizers and GA ₃ on fruit shape of some citrus fruit trees	16
2.7. Effect of some biofertilizers and GA ₃ on peel weight% of some citrus fruit trees	17
2.8. Effect of some biofertilizers and GA ₃ on juice weight% and volume of some citrus fruit trees:	20
2.9. Effect of some biofertilizers and GA ₃ on TSS% of some citrus fruit trees:	23
2.10. Effect of some biofertilizers and GA ₃ on acidity of some citrus fruit trees:	26
2.11. Effect of some biofertilizers and GA ₃ on TSS/Acid ratio of some citrus fruit trees:	29
2.12. Effect of some biofertilizers and GA ₃ on Vitamin C. of some citrus fruit trees	31
	ĺ

Contents	Page
3. Materials and methods	34
4. Results and discussion	40
4.1. Yield components	40
4.1.1 Yield weight	40
4.1.2. Number of fruits/tree	45
4.2. Physical characteristics of fruits	51
4.2.1. Fruit weight	51
4.2.2. Fruit volume	55
4.2.3. Fruit length	59
4.2.4. Fruit diameter	63
4.2.5. Fruit shape	67
4.2.6. Peel weight percentage	70
4.2.7. Juice weight percentage	74
4.2.8. Juice volume	78
4.3. Chemical characteristics of fruits.	82
4.3.1. Total soluble solids	82
4.3.2. Total acidity	85
4.3.3. TSS/Acid ratio	88
4.3.4. Vitamin C content	92
5. Summary and conclusion	97
6. Literature cited	104
7. Arabic summary	

÷. ·,

LIST OF TABLES

Table No.	Title	Page
A	Some physical and chemical properties of the experimental soil site	35
t	Yield weight (kg)/tree in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	43
2	Number of fruits/tree in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	49
3	Average fruit weight (g) in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	53
4	Average fruit volume (cm³) in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	57
5	Fruit length (mm) in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	61
6	Fruit diameter(mm) in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	65
7	Fruit shape (length/diameter ratio) in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons	68
8	Average peel weight%/fruit in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2003 and 2004 seasons.	72
'	Average juice weight%/fruit in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2002/2003 and 2003/2004 seasons	74
۱ (Average juice volume(cm³)/fruit in response to GA ₃ and some piofertilizers applied to Egyptian lime in 2002/2003 and 2003/2004 seasons	80

Table No.	Title	Page
11	Total soluble solids%(TSS%) in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2002/2003 and 2003/2004 seasons	83
12	Titeretable acidity%(as g. of citric acid/100g juice) in fruit juice in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2002/2003 and 2003/2004 seasons.	86
13	TSS/Acid ratio in fruit juice in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2002/2003 and 2003/2004 seasons	.90
14	V.C (mg of ascorbic acid)/100ml juice in fruits in response to GA ₃ and some biofertilizers applied to Egyptian lime in 2002/2003 and 2003/2004 seasons.	95

LIST OF FIGURE

Figure No.	Title	Page
1	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on fruit yield (kg-tree) of Egyptian lime in 2002 / 2003 and 2003/2004 seasons	44
2	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20 ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on number of fruits /tree of Egyptian lime in 2002/2003 and 2003 /2004 seasons.	50
3	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on average fruit weight(g)/fruit of Egyptian lime in 2002/2003 and 2003/2004 seasons	54
4	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20 ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on average fruit volume(cm ³)/fruit of Egyptian lime in 2002/2003 and 2003/2004 seasons	58
5	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20 ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on average fruit length(mm) of Egyptian lime in 2002/2003 and 2003/2004 seasons.	62
6	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20 ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on average fruit diameter(mm)of Egyptian lim in 2002/2003 and 2003/2004 seasons.	66
7	Effect of GA ₃ 5, 10ppm (2weeks after anthesis) and 10, 20 ppm (2 weeks pre-harvest date) sprayed on treated trees with some biofertilizers on average fruit shape (length/diameter ratio) of Egyptian lime in 2002/2003 and 2003/2004 seasons.	69
	.	