

شبكة المعلومات الجامعية







شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار في درجة حرارة من ١٥-٥٠ مئوية ورطوبة نسبية من ٢٠-٠٠% To be Kept away from Dust in Dry Cool place of 15-25- c and relative humidity 20-40%



بعض الوثائـــق الإصليــة تالفــة



بالرسالة صفحات لم ترد بالإصل

EARLY VITRECTOMY FOR DIABETIC VITREOUS HAEMORRHAGE IN TYPE 2 DIABETES MELLITUS

Thesis Submitted for partial fulfillment of M.D. degree In OPHTHALMOLOGY

ByKhaled El-Ghonemy El-Kotb (M.B., B.Ch., M.Sc.)

Discussed By

rof. Dr. El-Saieed Ebraheem El-Dsouky
Professor of Ophthalmology

Faculty of Medicine, Tanta University

Prof. Dr. Mostafa Kamal Nassar
Professor of Ophthalmology,
Faculty of Medicine, Menoufiya University.

Prof. Dr. Hassan Gamal El-Din Farahat

Professor of Ophthalmology, Faculty of Medicine, Menoufiya University.

FACULTY OF MEDICINE MENOUFIYA UNIVERSITY 2005

			i
			•

EARLY VITRECTOMY FOR DIABETIC VITREOUS HAEMORRHAGE IN TYPE II DIABETES MELLITUS

Thesis
Submitted for partial fulfillment of M.D. degree
In OPHTHALMOLOGY

ByKhaled El-Ghonemy El-Kotb
(M.B., B.Ch., M.Sc.)

Supervised By

Prof. Dr. Saber Hamed El-Sayed

Professor and head of Ophthalmology Department Faculty of Medicine, Menoufiya University

Prof. Dr. Omar Mohammed Afiffi

Professor of Ophthalmology, Faculty of Medicine, Zagazig University.

Prof. Dr. Mostafa Hamed Nabih

Professor of Ophthalmology, Faculty of Medicine, Cairo University.

Prof. Dr. Hassan Gamal El-Din Farahat

Professor of Ophthalmology, Faculty of Medicine, Menoufiya University.

Prof. Dr. Dominic McHugh

Consultant of Ophthalmology, King's College Hospital, London, UK.

FACULTY OF MEDICINE MENOUFIYA UNIVERSITY 2005

	-		



صدى الله العظيم البقرة ٣٢

LIST OF TABLES

Table	Page
Table (1) Prevalence of Retinopathy by Duration of Diabetes	10
Table (2) Prevalence of Retinopathy by Severity Level	10
Table (3) Ten-Year Incidence of any retinopathy, Progression of	11
Retinopathy, Progression to Proliferative Retinopathy Table (4) Causes of vitreous haemorrhage in different studies	17
Table (5) Natural Evolution of Massive Vitreous Haemorrhage In Patients with Diabetic Retinopathy	28
Table (6) Current Indications for Vitrectomy in Diabetic Eye Disease	44
Table (7) Objectives of vitrectomy for severe diabetic retinopathy	49
Table (8) Preoperative eye examination of studied patients.	82
Table (9) Iatrogenic and acute postoperative complications.	84
Table (10) Intraoperative findings in the studied	85
patients in comparison to DRVS findings	
Table (11) Lens condition at different times of follow up	86
Table (12) Retina condition at different times of follow up	88
Table (13) Occurrence of vitreous haemorrhage during different times of	90
follow up	92
Table (14) Visual acuity of studied patients before operation and at one month after	<i>72</i> .
Table (15) Visual acuity at six month	93
after operation and before operation	
Table (16) Visual acuity of studied patients before	94
operation and at 12 months after operation	0.5
Table (17) Visual acuity of studied patients before operation and at 18 months of operation	95
Table (18) Visual acuity of studied patients before operation and at 24	97
months of operation Table (19) Comparison of the distribution of best preoperative visual acuity	99
and visual acuity at 12 months	100
Table (20) Comparison of the distribution of best preoperative visual acuity and at 24 months	100
Table (21) Comparison of visual acuity score before operation and at 12	101
months Table (22) Comparison of visual acuity score before operation and after 24	102
months	102
Table (23) Comparison of visual acuity score at different times of follow up	103
Table (24) Comparison between proportions of patients with $VA \ge 6/12$ or no light perception between early vitrectomy group of DRVS (1985) and	105
current study at 12 and 24 months	40-
Table (25) Comparison of proportions of patients with visual acuity $\geq 6/12$ and no light perception between deferral group of the DRVS (1985) and the	107
current study at 12 and 24 months	

Table (26) Comparison between the early vitrectomy group of DRVS and	109
Table (27) Comparison between the determ group of the Dict of and current	111
study regarding incidence of retinal detachment	

....

LIST OF FIGURES

Figure	Page
Figure (1)Algorithm for diagnosing diabetes. PG, venous plasma glucose; challenge, 75g oral glucose tolerance test based on WHO (1999)guidelines.	5
Figure (2) Different possible locations for vitreous haemorrhage.	14
Figure (3) Various pathologic mechanisms of vitreous haemorrhage, (1) mechanical disruption of normal retinal vessels; (2) bleeding from diseased retinal vessels or abnormal neovessels; (3a&3b) extension of haemorrhage through the retinal from other sources.	16
Figure (4) Tangential traction is released by segmenting the formed vitreous and proliferative tissue between areas of retinal adhesions.	52.
Figure (5) A delaminating technique is used to excise areas of proliferative attachment to the retina along the inner surface of the internal limiting membrane with vitreous scissors.	53
Figure (6) An en bloc vitrectomy technique is performed with scissors passing through a tunnel in the formed vitreous	55
Figure (7) Anterior to posterior vitreous traction forces separate the formed vitreous from the retina when the areas of vitreoretinal traction are released using en bloc dissection	56
Figure (8) Series Ten Thousand® Vitrectomy machine (Alcon)	72
Figure (9) Accurs® vitrectomy machine (Alcon)	72
Figure (10) BSS® solution (Alcon)	72
Figure (11) Zeiss OPMI VISU 200®	72
Figure (12) Topcon OMS 600®	72
Figure (13) Perfluoropropane C3F8 lable	72
Figure (14) Different probes used in vitrectomy	73
Figure (15) The curved diode laser probe	73
Figure (16) Different vitreous scissors and forceps	74
Figure (17) Lenses used for vitrectomy in KCH, London	74
Figure (18) The diode laser (810 nm) meachine OcuLight SLx®, Iris medical	75
Figure (19) Vitrectomy pack for Accurs® machine, Alcon	75
Figure (20) Preoperative lens status in studied patients	83
Figure (21) Preoperative vitreous haemorrhage in studied patients	83
Figure (22) Lens condition at different times of follow up	87
Figure (23) Retinal condition at different times of follow up	89
Figure (24) Occurrence of vitreous haemorrhage	91
during different times of follow up Figure (25) Visual acuity of studied patients before operation and at 18 months of operation	96

Figure (26) Visual acuity of studied patients before	98
operation and at 24 months of operation	
Figure (27) Comparison of visual acuity score at different times of follow up	104
Figure (28) Comparison between proportions of patients with $VA \ge 6/12$ or	106
no light perception between early vitrectomy group of DRVS (1985) and	
current study at 12 and 24 months	100
Figure (29) Comparison of proportions of patients with visual acuity $\geq 6/12$	108
and no light perception between deferral group of the DRVS(1985) and the current study at 12 and 24 months	
Figure (30) Comparison between the early vitrectomy group of DRVS and	110
current study regarding incidence of retinal detachment	
Figure (31) Comparison between the deferral group of the DRVS and	112
current study regarding incidence of retinal detachment	
Figure (32) Preoperative ocular ultrasounds showing vitreous haemorrhage	113
with no sings of tractional retinal detachment in patients with type 2 diabetes mellitus.	
Figure (33) Patient No 2, 1 st postoperative day	114
Figure (34) Patient No 3, 1 st postoperative day	114
Figure (35) Patient No 5, 6 months postoperative visit	114
Figure (36) Patient No 20, 1 st postoperative day	114
Figure (37) Patient No 20, 3 months postoperative visit	114
Figure (38) Patient No 20, 18 months postoperative visit	114
Figure (39) View of wide field lense photographed via the operating	115
microscope	
Figure (40) Photograph of case No. 1 in the first	115
postoperative day showing diode laser scores	116
Figure (41) 1st postoperative fundus photograph of patient no. 26 with vitreous haze secondary to mild recurrent vitreous haemorrhage	110
Figure (42) Fundus photograph of patient no. 28	116
with C3F8 in 1st postoperative day	
Figure (43) Photograph of case no. 40, 3 months postoperatively	117
Figure (44) Photograph of the same patient 6 months postoperatively	117
Figure (45) Photograph of patient no. 42, one month postoperatively	118
Figure (46) Photograph of patient no. 45, 24 months postoperatively	118
Figure (47) Photograph of anterior segment of	119
patient no. 33 showing NVI (iris neovessels).	
Figure (48) Photograph of the same patient	119
with PCIOL 18 months postoperatively	120
Figure (49) Posterior subcapsular cataract of patient no. 26, 12 months postoperatively	120
Figure (50) Fundus photograph of patient no. 37 with	120
recurrent retinal detachment with impending phthisis bulbi	

G,

,			
	•		