

# 





ثبكة المعلومات الجامعية





### جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15-20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of 15-25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية







### BIOLOGICAL STUDIES ON *Neochetina eichhorniae*AND *Neochetina bruchi*, TWO SPECIFIC WEEVILS FOR CONTROLLING WATER HYACINTH

By

Islam Rashed Marzouk El-Zoghby

B.Sc. Agric. (Plant Protection), Fac. of Agric., Moshtohor, Zagazig Univ. (Benha Branch) (1999)

#### **THESIS**

Submitted in partial fulfillment of the requirements for the degree

of

Master of Agric. Science

In

**Economic Entomology** 

(Biological Control)

Plant Protection Department
Faculty of Agriculture, Moshtohor
Benha Branch, Zagazig University



2005

• 

#### **SUPERVISION COMMITTEE**

## AND *Neochetina bruchi*, TWO SPECIFIC WEEVILS FOR CONTROLLING WATER HYACINTH

By

Islam Rashed Marzouk El-Zoghby
B.Sc. Agric. (Plant Protection), Fac. of Agric., Moshtohor,
Zagazig Univ. (Benha Branch) (1999)

#### Under the supervision by:

Prof. Dr. Fawzy Faiek Shalaby

Prof. of Economic Entomology,

Plant Protection Dept., Fac. of Agric. Moshtohor,

Zagazig University (Benha Branch)

Prof. Dr. Yahia Hussein Fayad

Chief Researcher of Biological Control,

Dept. of Biological Control, Plant Protection Research Institute, ARC

Prof. Dr. Adel Abd El-Hameid Hafez

Prof. of Economic Entomology, Plant Protection Dept., Fac. of Agric. Moshtohor.

Zagazig University (Benha Branch)

A.A. Hafez

Plant Protection Department Faculty of Agriculture, Moshtohor Benha Branch, Zagazig University

#### **APPROVAL SHEET**

## AND *Neochetina bruchi*, TWO SPECIFIC WEEVILS FOR CONTROLLING WATER HYACINTH

By

Islam Rashed Marzouk El-Zoghby B.Sc. Agric. (Plant Protection), Fac. of Agric., Moshtohor, Zagazig Univ. (Benha Branch) (1999)

This thesis for M. Sc. degree has been Approved by:

Prof. Dr. Mohamed Fouad S. Tawfik

Mit i Ta 3

A. A. Halez

Prof. of Economic Entomology, Fac. Agric., Cairo University (Chairman)

Prof. Dr. Ezzat Farag El-Khayat

Prof. Dr. Economic Ent., Head of Plant Protection Dept., Fac. Agric., Moshtohor, Zagazig University, Benha Branch

Prof. Dr. Fawzy Faiek Shalaby

Prof. of Economic Entomology, Fac. Agric., Moshtohor, Zagazig University


Prof. Dr. Adel Abd El-Hameid Hafez

Prof. of Economic Entomology, Fac. Agric., Moshtohor, Zagazig University

Prof. Dr. Yahia Hussein Fayad

Chief Researcher (Professor) of Biological Control, Plant Protection Research Institute, ARC

Date of examination Sun. 29/5/2005.



#### ACKNOWLEDGEMENT

#### Ultimate Thanks to ALLAH

The writer wishes to express his deep thanks and gratitude to the committee members who supervised this work; Dr. Fawzy Faiek Shalaby, Emeritus Professor of Economic Entomology, Plant Protection Department, Faculty of Agriculture at Moshtohor, Benha branch, Zagazig University; Prof. Dr. Yahia Hussein Fayad, Emeritus Chief Researcher, Department of Biological Control, Plant Protection Research Institute, (PPRI) A.R.C.; and Dr. Adel Abd El-Hameid Hafez Professor of Economic Entomology, Plant Protection Department in the former Faculty and University. To them, the author is grateful for suggesting the problem, supplying all the needed facilities, keen & close supervision and valuable scientific advices& criticism throughout the whole period of study, and also for their great efforts for revising the manuscript.

Deep thanks are also due to Dr. *Amal Ahmed El-Zoghby*, Senior Researcher, Department of Biological Control, (PPRI) who was very helpful throughout the period of achieving the presented experiments. Her scientific advices and help are greatly acknowledged. Also, thanks to Prof. Dr. *Amira Abd El-Hameid Ibrahim*, Emeritus Chief Researcher in the same Department for her encouragement.

Thanks are also due to Prof. Dr. *Mohammed M. Abu-Setta*, Chief Researcher, Scale Insects & Mealy-Bugs Research Department (PPRI) who paid great efforts for achieving the statistical analysis of the obtained data. To him, the writer is indebted with thankfulness and gratefulness.

The author wishes also to express deep thanks to Dr. Ahmed Ramadan Ibrahim, Researcher in the Department of Vegetable Crop Pests (PPRI) for his encouragement and help during conducting this work.

Great thanks to the staff- members and colleagues in the Plant Protection Department, Faculty of Agriculture at Moshtohor for morality and continuous encouragement during the period of the study.

It is of great importance to express deep thanks to the Egyptian – French Project for the Biological Control of Water Hyacinth for providing *Neochetina* spp. and other materials needed for the study.

Finally, deep thanks to all members of my family for patience and cordial encouragement throughout the whole period of the study.

#### **CONTENTS**

	Page
1. INTRODUCTION	1
2. REVIEW OF LITERATURE	4
1- Distribution of water hyacinth	4
2- Effect of N. bruchi on water hyacinth	8
2.1. On flowering	8
2.2. Damage to leaves and shoots	10
3- Water loss by water hyacinth	13
4- Mass rearing of the weevils	17
5- Biological and morphological studies on Neochetina spp	21
6- Management of water hyacinth.	28
6.1. Biological control	28
A- With arthropods	28
B- With pathogens	37
6.2. Mechanical control	40
6.3. Chemical control	41
6.4. Integrated control	43
3. MATERIALS AND METHODS	48
1- Cages used for growing water hyacinth	48
2- Obtaining weevil species and artificial infestations to water	
hyacinth plants	51
3- Effect of N. bruchi adults in saving water loss through water	
hyacinth plants	52
4- Estimating the effect of N. bruchi on flowering of water	
hyacinth	54

	Page
5- Biological and morphological studies on Neochetina spp	54
5.1. Neochetina bruchi	54
5.2. N. bruchi & N. eichhorniae	55
6- Mounting mouth parts of Neochetina spp. larvae	55
7- Statistical analysis	55
4. RESULTS AND DISCUSSION	56
1- Role of N. bruchi in reducing water loss in water hyacinth	
infested areas	56
2- Effect of <i>N. bruchi</i> on flowering of water hyacinth	62
3- Biological and morphological studies on <i>N. bruchi</i> and <i>N.</i>	
eichhorniae	68
A- Egg stage	68
B- Larval stage	72
C- Papal stage	80
D- Adult stage	85
4- Productivity of eggs by <i>N. bruchi</i> and adults longevity	88
5- Reproduction potential and life table parameters	89
5. SUMMARY	92
	98
6. CONCLUSION	102
7. REFERENCES	,