Contents

Subjects	Page
List of abbreviations	II
List of Figures	IV
List of Tables	V
• Introduction	1
Aim of the work	5
• Review of Literature	
♦ Chapter (1): Cesarean Section	6
◆ Chapter (2): Postoperative Infective Complication	ons30
♦ Chapter (3): Antibiotic Prophylaxis	58
Patients and Methods	80
• Results	95
• Discussion	114
• Summary	120
• Conclusion	125
• Recommendations	126
• References	127
Arabic Summary	

List of Abbreviations

ACOG: American College of Obstetricians and

Gynecologists.

ASHP: American Society of Health-System

Pharmacists.

BMI: body mass index.

CBC: Complete Blood Count.

CD: Cesarean delivery.

CDC: Centers for Disease Control and Prevention.

CDI: Clostridium difficile infection.

Cfu: Colony forming unit.

CI: Confidence interval.

CMACE: Centre for Maternal and Child Enquiries.

CS: Cesarean section.

HAART: Highly Active Anti-Retroviral Therapy.

HIV: Human Immunodeficiency Virus.

HSV: Herpes Simplex Virus.

IDSA: Infectious Diseases Society of America.

IU: International Unit

MRSA: Methicillin-resistance S. aureus .

List of Abbreviations

NICE: National Institute for Health and Clinical

Excellence.

NICHD: National Institute of Child Health and Human

Development. SMFM: Society of Maternal-

Fetal Medicine.

RCT: Randomized Controlled Trial

RR: Relative risk.

SHEA: Society for Healthcare Epidemiology of

America.

SIGN: Scottish Intercollegiate Guidelines Network.

SIS: Surgical Infection Society.

SSI: Surgical site infection.

UK: United Kingdom.

UTI: Urinary tract infection.

WHO: World Health Organization.

List of Figures

Figure	Title	Page
Figures in Review		
<u>1</u>	Normal healing response in full-thickness wounds.	39
Figures in Results		
1	Incidence of wound infection in both study groups.	100
<u>2</u>	Grading of wound infection in both study groups.	101
<u>3</u>	Timing of development of wound manifestations in patients with wound involvement in both study groups.	103
4	Severity of wound manifestations in patients with wound involvement in both study groups.	105
5_	Kaplan-Meier curves for time to wound infection of any grade. Hazard ratio (group B/group A), 0.90 (95% CI 0.56 to 1.43; p-value 0.632).	107
<u>6</u>	Incidence of other unwanted maternal outcomes in both study groups.	111
<u>7</u>	Incidence of unwanted neonatal outcomes in both study groups.	113

List of Tables

Table	Title	Page
Tables in Review		
1	Common Indications for Cesarean Delivery.	7
<u>2</u>	Antimicrobial prophylaxis for gynecologic and obstetric surgery in adults.	66
Tables in Results		
**************************************	Patients' characteristics in both study groups.	96
<u>2</u>	Surgical data and hospital length of stay in both study groups.	97
<u>3</u>	Incidence and grading of wound infection in both study groups.	99
<u>4</u>	Timing and severity of wound manifestations in patients with wound involvement in both study groups.	102
<u>5</u>	Kaplan-Meier analysis for the time to wound infection of any grade.	106
<u>6</u>	Incidence of other unwanted maternal outcomes in both study groups.	108
7	Incidence of unwanted neonatal outcomes in both study groups.	112



Introduction





Aim of the Work





Review of Literature





Chapter 1

Cesarean Section





Chapter 2

Postoperative Infective Complications





Chapter 3

Antibiotic Prophylaxis





Patients and Methods





Results





Discussion





Summary

