

**EFFECT OF GROWTH PROMPTING ON
PRODUCTIVE AND REPRODUCTIVE
PERFORMANCE OF HEIFERS**

By

MOHAMED ABD EL-FATAH EL-SAMAHY

B.Sc. Agric. Sc. (Animal Production), Ain Shams University, 2003

A thesis submitted in partial fulfillment

of

**the requirements for the degree of
MASTER OF SCIENCE**

in

**Agricultural Science
(Animal Nutrition)**

**Department of Animal Production
Faculty of Agriculture
Ain Shams University**

2011

Approval Sheet

**EFFECT OF GROWTH PROMPTING ON
PRODUCTIVE AND REPRODUCTIVE
PERFORMANCE OF HEIFERS**

By

MOHAMED ABD EL-FATAH EL-SAMAHY

B.Sc. Agric. Sc. (Animal Production), Ain Shams University, 2003

This thesis for M.Sc. degree has been approved by:

Dr. Hassan Mahmoud El-Gendy

Prof. Emeritus of Animal Nutrition, Faculty of Agriculture, Tanta
University

Dr. Salwa Mahmoud Hamdy

Prof. Emeritus of Animal Nutrition, Faculty of Agriculture, Ain
Shams University

Dr. Mahmoud Mohamed Aly Khorshed

Associate Prof. of Animal Nutrition, Faculty of Agriculture, Ain
Shams University

Date of Examination: 9/ 11/ 2010

EFFECT OF GROWTH PROMPTING ON PRODUCTIVE AND REPRODUCTIVE PERFORMANCE OF HEIFERS

By

MOHAMED ABD EL-FATAH EL-SAMAHY

B.Sc. Agric. Sc. (Animal Production), Ain Shams University, 2003

Under the supervision of:

Dr. Mahmoud Mohamed Aly Khorshed

Associate Prof. of Animal Nutrition, Department of Animal
Production, Faculty of Agriculture, Ain Shams University
(Principal Supervisor)

Dr. Mohamed Ali Abd El-Moniem El-Ashry

Prof. Emeritus of Animal Nutrition, Department of Animal
Production, Faculty of Agriculture, Ain Shams University

Dr. Samer Ali Ibrahim

Head of Research of Animal Husbandry, Department of Cattle
Breeding Research, Animal Production Research Institute,
Agricultural Research Center

تأثير دفع النمو على الأداء الإنتاجي والتناسلي للعجلات

رسالة مقدمة من

محمد عبد الفتاح بسيوني السماحي

بكالوريوس علوم زراعية (إنتاج حيواني)، جامعة عين شمس، ٢٠٠٣

للحصول على

**درجة الماجستير في العلوم الزراعية
(تغذية حيوان)**

قسم الإنتاج الحيواني

كلية الزراعة

جامعة عين شمس

جامعة عين شمس
كلية الزراعة

رسالة ماجستير

اسم الطالب : محمد عبد الفتاح بسيونى السماحي
عنوان الرسالة: تأثير دفع النمو على الأداء الإنتاجي والتناسلي للعجلات
اسم الدرجة : ماجستير في العلوم الزراعية (تغذية حيوان)

لجنة الإشراف :

د. محمود محمد علي خورشيد
أستاذ تغذية الحيوان المساعد، قسم الإنتاج الحيواني، كلية الزراعة، جامعة عين شمس
(المشرف الرئيسي)

د. محمد علي عبد المنعم العشري
أستاذ تغذية الحيوان المتفرغ، قسم الإنتاج الحيواني، كلية الزراعة، جامعة عين شمس

د. سمير علي ابراهيم
رئيس بحوث رعاية الحيوان، قسم بحوث تربية الأبقار، معهد بحوث الإنتاج الحيواني،
مركز البحوث الزراعية

تاريخ التسجيل: ٢٠٠٥/٩/٥

الدراسات العليا

أجيزت الرسالة بتاريخ
٢٠١٠ / ١١ / ٩

ختم الإجازة

موافقة مجلس الجامعة
٢٠١٠ / /

موافقة مجلس الكلية
٢٠١٠ / /

صفحة الموافقة على الرسالة

تأثير دفع النمو على الأداء الإنتاجي والتناسلي للعجلات

رسالة مقدمة من

محمد عبد الفتاح بسيوني السماحي

بكالوريوس علوم زراعية (إنتاج حيواني)، جامعة عين شمس، ٢٠٠٣

للحصول على

**درجة الماجستير في العلوم الزراعية
(تغذية حيوان)**

وقد تمت مناقشة الرسالة والموافقة عليها

اللجنة :

د. حسن محمود الجندى

أستاذ تغذية الحيوان المتفرغ، كلية الزراعة، جامعة طنطا

د. سلوى محمود حمدى

أستاذ تغذية الحيوان المتفرغ، كلية الزراعة، جامعة عين شمس

د. محمود محمد علي خورشيد

أستاذ تغذية الحيوان المساعد، كلية الزراعة، جامعة عين شمس

تاريخ المناقشة: ٢٠١٠/١١/٩

ABSTRACT

Mohamed Abd El-Fatah Basuony El-Samahy: Effect of Growth Prompting on Productive and Reproductive Performance of Heifers. Unpublished M.Sc. Thesis, Department of Animal Production, Faculty of Agriculture, Ain Shams University, 2010.

The objective of this experiment was to evaluate the effects of altered forage: concentrate ratio and DL-methionine analog supplementation on growth performance of growing heifers until confirmed pregnant. Twenty-four Holstein heifers with an average of age 5.6 ± 0.36 mo and weight 116 ± 6 kg, were fed restricted intakes formulated to allow for 800 g/d (NRC, 1989). Experimental diets were formulated to contain either 50: 50 (low forage) or 75: 25 (high forage) forage: concentrate ratio diets (DM basis). Forage comprised of corn silage and constant amounts of berseem hay (1.5 and 2 kg/ h/ d for low and high forage, respectively). Treatments were fed with or without methionine hydroxy analog supplementation (MHA, 2.1 g / day/ 100 kg BW). Live body weight was taken biweekly, while some body measurements and blood samples were taken monthly. Blood plasma was taken twice weekly after insemination to determine the plasma progesterone concentration. Digestibility trial was performed at 370 kg BW, during the digestibility trial heifers were fed individually in tie-stall. Actual intakes of nutrients were lower than expected intakes due to lower corn silage DM ratio. DM, OM, CP, and NFE digestibility were significantly ($P < 0.05$) improved by reducing the forage portion of the diet, however CF digestibility had a significant ($P < 0.05$) opposite direction. While, supplementation of MHA resulted in trends toward increased DM digestibility. Ruminal TVFA were significantly higher in low forage groups, however, ruminal PH was higher in low forage groups, while MHA trend to decreased TVFA only. Insignificant differences were observed for

plasma total protein triglyceride, creatinine, urea, in addition to negligible difference for albumin, globulin, and A/G ratio, while significant ($P < 0.05$) decreasing in plasma cholesterol was recorded for low forage without MHA treatment, also high forage without MHA group showed a trend to decreased plasma urea. Throughout the feeding period, ADG was not affected across all treatment rations (0.681 LF, 0.685HF, 0.695 LFM, and 0.663 HFM, $SE \pm 0.033$ kg/d). Gain of heifers body weight and measurements were not different among treatments. However, low forage groups had better feed-conversion for DM, ME and TDN, while CP and DCP conversion were better in high forage groups. Age at 330 kg BW was recorded an average of 15.98 mo (15.85 LF, 16.26 HF, 15.41 LFM, and 16.38 HFM, $SE \pm 0.53$ mo), as well as BW at age 14 mo and age and BW at AI, or conception not affected by treatments. Total and daily feed cost for high forage groups was significantly ($P < 0.05$) better than low forage groups, but total and gain feed cost from 150 to 330 kg BW did not affect by treatment may be due to the deep gap within treatments, especially in high forage groups. Thus, we can feed growing heifers in this tested forage strategy under quality recommended for feedstuffs, and used satisfactory method for feeding. Notwithstanding, MHA addition did not has effect on growing heifers performance, nevertheless MHA addition may has influences on DM digest and blood metabolism, and may be the amount used under this experimental was not enough to improve growth performance.

Keywords: growing heifers, forage: concentrate ratio, DL-methionine, forage strategy, nutrients digestibility, rumen parameters, blood plasma parameters, growth and reproductive performance, plasma progesterone.

ACKNOWLEDGMENT

All thanks are due to ALLAH (الله), the most gracious, most beneficent most merciful for the help and guidance to achieve goals and make them possible.

I wish to express my sincere appreciation and deepest gratitude to **Dr. M. A. El-Ashry**, Professor Emeritus of Animal Nutrition, Animal Production Department, Faculty of Agriculture, Ain Shams University, for continuous supervision support, guidance and constant throughout most stages of this work.

I like to express my thanks to **Dr. M. M. Khorshed** Assistant Professor of Animal Nutrition, Animal Production Department, Faculty of Agriculture Ain Shams University, for help and encouragement throughout the course of the study, and also to **Dr. S. A. Ibrahim**, Assistant Professor of Animal Breeding, Animal Production Research Institute, Agriculture Research Center, for continuous help and advice during the experimental work.

Many thanks are also due to the chief of El-Karada Animal Experimental Station **Dr. M. A. Tag**, Assistant Professor of Animal Rearing, Animal Production Research Institute, Agriculture Research Center, for his advice, providing most necessary facilities required for the experimental work. I need to especially thank to **Dr. N. E. El-Bordeny** Assistant Professor of Animal Nutrition, Animal Production Department, Faculty of Agriculture Ain Shams University, for advice and great help in the laboratory work, and I wish to thank **Dr. Sayed Wahdan** for her veterinary assistance required for the experimental work.

Deep thanks for all nutrition staff members of Animal Production and Poultry Production departments, Faculty of Agriculture, Ain Shams University, and to the staff members of the Animal Production Research Institute, Agriculture Research Center, for they offered to make this work possible.

DEDICATION

This thesis is dedicated to souls of my fathers **Mr. Abd El-Fatah El-Samahy** and Prof. **Dr. Mohamed Abd El Moniem El-Ashry**, may **ALLAH** (الله) save their souls. Their constant support, encouragement and love gave me the strength and ability to believe in myself and believe that anything is possible.

Certainly, grateful dedication should be also extended especially to my mother, my sister and my friends for the continuous encouragement and help during all my life.

CONTENTS

	Page
LIST OF TABLES	IV
LIST OF FIGURES	V
LIST OF ABBREVIATIONS	VI
I. INTRODUCTION	1
II. REVIEW OF LITERATURE	3
GROWING HEIFERS	3
FEED STRATEGY	5
DL-Methionine / MHA	7
2.1. Feed intake & Digestion coefficients	11
2.2. Rumen liquor parameters	23
2.3. Blood metabolites parameters	32
2.4. Growth parameter	38
2.4.1. Body weight & Daily gain	39
2.4.2. Body measurements	46
2.5. Reproductive performance	52
2.5.1. Reproductive parameters	52
2.5.2. Level of Progesterone	59
III. MATERIALS AND METHODS	63
3.1. DL- Methionine	63
3.2. Experimental animals	63
3.3. Experimental rations	64
3.4. Experimental animals management	65
3.5. Digestibility trials	65
3.5.1. Feces collection	65
3.5.2. Feedstuffs and feces analysis	66
3.6. Feed conversion	66
3.7. Rumen liquor	66
3.7.1. Sampling	66

II

3.7.2. Rumen fluid analysis	67
3.8. Blood parameters	67
3.8.1. Blood sampling	67
3.8.2. Blood plasma analysis	67
3.9. Monitoring growth	68
3.9.1. Body weight	68
3.9.2. Measuring body	68
3.10. Reproductive performance	68
3.11. Economical efficiency	68
3.12. Statistical analysis	69
II. RESULTS AND DISCUSSION	70
4.1. Effect of experimental treatments on apparent digestibility coefficients and nutritive values	70
4.2. Effect of experimental treatments on feed intake	74
4.3. Effect of experimental treatments on some Rumen liquor parameters	83
4.3.1. Ruminal pH	83
4.3.2. Ruminal total volatile fatty acids (TVFA)	85
4.4. Effect of experimental treatments on blood plasma metabolites	86
4.4.1. Blood plasma proteins (Total protein, Albumin, Globulin, and Albumin: Globulin ratio)	86
4.4.2. Blood plasma Urea	89
4.4.3. Blood plasma creatinine and urea: creatinine ratio	90
4.4.4. Blood plasma triglyceride	90
4.4.5. Blood plasma cholesterol	91
4.5. Effect of experimental treatments on Growth performance	93
4.5.1.1. Body weight & Daily gain	93
4.5.1.2. Feed conversion of experimental treatments	95
4.5.2. Body measurements	98
4.6. Effect of experimental treatments on reproductive	103

III

performance	
4.6.1. First estrus, First insemination, and conception	103
4.6.2. Level of progesterone	106
4.7. Economic efficiency of experimental treatments	109
V. SUMMARY AND CONCLUSION	113
VI. REFERENCES	116
ARABIC SUMMARY	

IV

LIST OF TABLES

Table		Page
1	Chemical composition of feed stuffs.	70
2	Effect of the experimental rations on apparent digestion coefficients and nutritive values % (DM basis)	72
3	Daily feed intakes component of two forage levels (DM basis), according to means of feedstuff analysis of digested trial	77
4	Daily digested component intakes of treatment rations	80
5	Effect of the experimental rations on some rumen liquor parameters, at 0, 3, and 6 hours after beginning of feeding	83
6	Effect of the experimental rations on Blood plasma proteins (Total protein, Albumin and Globulin)	88
7	Effect of the experimental rations on Blood plasma urea, creatinine, triglyceride, and cholesterol	92
8	Effect of the experimental treatments on LBW and average daily gain	94
9	Effect of the experimental treatments on feed conversion	97
10	Effect of the experimental rations on some body measurements and average daily gain	100
11	Effect of the experimental rations on some reproductive parameters	106
12	Effect of the experimental rations on plasma progesterone, at and after conception AI	108
13	Effect of the experimental rations on feed cost	110
14	Effect of the experimental rations on economic efficiency	111
15	Stander deviation value from some parameter that may be confused with economic evaluation	112

LIST OF FIGURES

Figure		page
1	Schematic of the conversion of D-HMB, L-HMB and D-methionine to L-met.	10
2	The related of plane ration with body weight, for growing heifers	64
3	Relationship between LBW and distribution sources of (OM, CP, CF, and NFE) intake according to feed strategy	79
4	Relationship between LBW and daily intake from digested materials, TDN, and ME, and CP: ME ratio	82
5	Effect of experimental treatments on ruminal pH	84
6	Effect of experimental treatments on ruminal TVFA	85
7	Effect of experimental treatments on LBW at fixed ages	95
8	Effect of experimental treatments on plasma progesterone after conception	109