

Role of PET/CT in diagnosis, staging and follow up of HCC

Essay

Submitted for partial fulfillment of
Master Degree in Radiodiagnosis

By

Osama Mohamed Ahmed Montaser

M B, B Ch

Faculty of medicine Cairo Univerisity

Supervised By

Prof Dr :Abeer Abd Elmaksood Hafez

Professor of Radiodiagnosis

Faculty of Medicine

Ain shams university

Dr. Yosra Abdelzaher Abdallah

Lecturer of Radiodiagnosis

Ain shams university

Radiodiagnosis Department

Faculty of Medicine

Ain shams University

2014

Acknowledgement

First and foremost, I submit all my gratitude to ALLAH to whom I owe every success in my life.

I would like to express my sincere appreciations and profound gratitude to Prof. Dr.Abeer Abd Elmaksood Hafez, Professor of Radiodiagnosis, Faculty of Medicine –Ain shams University, for her help, kind guidance, continuous support & giving me such an honor to work under her supervision.

I would like to express my respect, appreciation, and thanks for Dr.yosra Abdelzaher, Lecturer of Radiodiagnosis, Ainshams university, for her assistance, encouragement, invaluable guidance, constructive criticism and great help in supervising this work,

*To my parents,
And my wife.*

*FOR
THEIR HELP,
SUPPORT,
GREAT CARE,
ENCOURAGEMENT
AND CONTINUOUS PUSH*

Abstract

PET/CT is superior to PET and CT alone, and/or magnetic resonance imaging (MRI), in the diagnosis and treatment of various primary or metastatic cancers.

Dual modality PET/CT scanning provides accurately fused morphologic (CT) and functional (PET) data sets. A very small tumor is well detected by PET but can be missed by CT. On the other hand, a large tumor with minimal functional deviations may be seen on a CT image, but may not be detected by PET. In both situations, PET/CT would localize the tumor accurately. Thus, PET/CT is a more accurate test than either of its individual components.

PET/CT has advantages over other imaging methods; it can differentiate benign from malignant lesions, staging and restaging tumors, detect functional changes before there is any change in clinical or radiological size of a mass, better in identifying cancer that has spread, making up treatment plan and monitoring tumor response, distinguish viable metabolically active tissue from scars, and it is indicated for restaging in patients with suspected recurrent and metastatic disease.

Key words :

PET CT - Hepatitis C Virus - Aflatoxin B .

LIST OF ABBREVIATION

LLS: Left lobe segment donation

FCAT: Federative Committee on Anatomical Terminology

RHV: Right hepatic vein

MHV: Middle hepatic vein

LHV: Left hepatic vein

IVC: Inferior vena cava

MP: Main portal vein

CM: centimeter

SMA :Superior mesenteric artery

CT:Computed tomography

HU:Housfield unit

FL: Falciform ligament

PET: Positron emitted tomography

FDG: Fluoro Deoxy Glucose

PET CT: Position emission tomography with computed tomography

CNS: Central nervous system

CVS:Cardio vascular system

¹⁸F FDG: ¹⁸Fluorine labeled 2 fluoro 2 deoxy glucose

GCSF :Granulocyte colony-stimulating factor

HCC:Hepato cellular carcinoma

HCV: Hepatitis C Virus

RNA: Ribonucleic acid

HBV: Hepatitis B Virus

AFB: Aflatoxin B

AFM1: Aflatoxin M1

H&E: Hematoxylin and Eosin

NAFLD: Non-alcoholic fatty liver disease

TNM: Tumor Node Metastasis

UICC: International Union against Cancer

MRI: Magnetic resonant imaging

KEV: Kilo electron volt

MEV: Milli electron volt

PDGF: Platelet Derived Growth Factor

VEGF: Vascular Endothelial Growth Factor

BFGF: Basic Fibroblast Growth Factor

GLUT: Glucose Transporters

PMTs: Photomultiplier tubes

NaI (TI): Thallium-doped sodium iodide

BGO: Bismuth Germinate

LSO: Lutetium Oxyorthosilicate

GSO: Gadolinium Silicate

SPECT: Single Photon Emission Computed tomography

KV: Killo Volt

LOR: Line of response

¹⁸F: Fluorine 18

¹³N: Nitrogen 13

¹¹C: Carbon 11

¹⁵O: Oxygen 15

⁸²Rb: Rubidium 82

β⁺: Positron

ν: neutrino

γ : Photon

N: neutron

P: proton

H⁺: Hydrogen ion

Z: Atomic number

mCi: Millicurie

IV: Intravenous

AC/AL: Attenuation correction/Alignment

SUV: Standardized uptake value

CT FOV: Computed Tomography Field of View

AFP: Alfa-fetoprotein

US: Ultrasonography

CO₂: Carbon Dioxide

¹¹C-ACT: ¹¹ carbon acetate

BCLC: Barcelona Clinic Liver Cancer

¹⁸FDG-6-P: ¹⁸ F-FDG-6-phosphate

¹⁸ FDGal: ¹⁸ fluoro-2-deoxy-D-galactose

FCH : Flurocholine

¹¹C-choline: ¹¹ carbon choline

MDCT: Multi detector computed tomography

TACE: Transarterial chemoembolization

RFA: Radiofrequency ablation

PEI: Percutaneous ethanol injection

PVT: Portal vein thrombosis

PV: Portal vein

LIST OF FIGURES

Figure 1.1 The surfaces and external features of the liver.....	6
Figure 1.2 showing the Relations of the liver.....	8
Figure 1.3 showing Segmentation of the liver – Couinaud.....	12
Figure 1.4 Drawing illustrates the segmental anatomy of the liver.....	13
Figure 1.5 Segmental anatomy according to Couinaud.....	13
Figure 1.6 H. Bismuth’s functional classification of the liver.....	15
Figure 1.7 Dissection to show the relations of the hepatic artery, bile duct and portal vein to each other in the lesser omentum: anterior aspect.....	19
Figure 1.8: The portal vein and its tributaries (semi-diagrammatic).....	20
Figure 1.9 Arrangement of the hepatic venous territories.....	20
Figure 1.10 Biliary system anatomy.....	22
Figure 1.11(a–d) CT scan through the liver with Couinaud’s segments divided and numbered.....	25
Figure 1.12 Computerized tomography of the liver. The vessels can be recognized and used as landmarks to define the different segments.....	26
Figure 2.1 Hepatocellular carcinoma. A cirrhotic liver with multiple tumor nodules scattered throughout.....	35
Figure 2.2 Hepatocellular carcinoma. A cirrhotic liver with a solitary encapsulated tumor.....	35
Figure 3.1 Uptake of FDG.....	44
Figure 3.2 Annihilation reaction.....	45
Figure 3.3 Radial blurring.....	47
Figure 3.4 Mean positron range and annihilation angle blurring.....	48
Figure 3.5 Coincidence imaging.....	49
Figure 3.6 A schematic illustration of a PET CT system.....	51
Figure 3.7 PET/CT image consisting of coronal whole-body CT image.....	53
Figure 3.8 Photograph (side view) of a hybrid PET-CT scanner shows the PET (P) and CT (C) components.....	53

Figure 4.1 Normal distribution of FDG.....	66
Figure 4.2 Typical scout image obtained during an FDG PET/CT study.....	77
Figure 4.3 Typical imaging protocol for combined PET/CT.....	78
Figure 4.4 Display screen of the syngo software platform shows fused PET/CT....	80
Figure 4.5 High-density metallic implants generate streaking artifacts.....	86
Figure 4.6 Attenuation correction artifacts due to oral contrast agents.....	87
Figure 4.7 Attenuation correction artifacts due to IV contrast.....	87
Figure 4.8 Truncation artifact.....	88
Figure 4.9 Misregistration artifacts.....	89
Figure 5.1 Patient with left hepatic mass incidentally detected on ultrasound. PET/CT was requested for further assessment, and revealed FDG avidity of the hepatic mass.....	95
Figure 5.2 PET/CT shows a solitary focal hypermetabolic focus in the right lobe of the liver.....	95
Figure 5.3 47-year-old male patient on whom was performed a liver CT and ^{18}F FDG / ^{11}C acetate PET/CT for the evaluation of suspicious recurrent Rt HCC.....	96
Figure 5.4 Detection of HCC with ^{18}F -FDG PET/CT and ^{11}C acetate PET/CT on transaxial sections of liver and chest.....	96
Figure 5.5 (A–C) Transaxial PET/CT images of 71-y-old man in whom HCC of Edmondson and Steiner’s grade 1 was diagnosed.....	97
Figure 5.6 FDGal PET/CT image (a) and CECT image in arterial phase (b) of a large necrotic HCC with adjacent viable tumor tissue.....	100
Figure 5.7 FCH (a) and FDG (b) PET/CT images of patient with diffuse recurrence of HCC involving all the liver.....	100
Figure 5.8 A 45-year-old man with moderately differentiated HCC in the right posterior lobe and a hemangioma in the left lobe of the liver confirmed by the histopathological examination after surgery A , Multiphase contrast-enhanced MRI. B , Transaxial ^{18}F FDG PET/CT. C , Transaxial ^{11}C -choline PET/CT	102
Figure 5.9 Transaxial PET/CT images of 63-y-old man in whom HCC of Edmondson and Steiner’s grade 2 was diagnosed. ^{11}C -acetate PET maximal-intensity image (A) and integrated PET/CT images (B).....	106
Figure 5.10 Mediastinal metastasis by ^{18}F -FDG and ^{11}C -ACT PET/CT.....	107

Figure 5.11 Multifocal bone metastases by ^{18}F -FDG and ^{11}C -ACT PET/CT.....	107
Figure 5.12 a Axial contrast-enhanced computed tomography (CT) image 90 days after a radiofrequency ablation (RFA) procedure in the right liver lobe b Corresponding axial positron emission tomography PET/CT image: c Corresponding axial PET image d Axial contrast enhanced CT image 230 days after the RFA procedure in the right liver lobe e Corresponding PET/CT image.....	111
Figure 5.13 A 50-year woman who had HCC resection 2 years before had an intrahepatic HCC recurrence and received combined RFA and TACE treatment. A highly metabolically active lesion was detected on the top of the lesion by PET and PET/CT fused images (white arrows, A-C and G).....	112
Figure 5.14 Images of 53-y-old woman with viable (20% necrosis) lipiodolized HCC. Multiphasic contrast-enhanced CT images (A and B), ^{18}F -FDG PET/CT images (C and D).....	115
Figure 5.15 Images of 55-y-old man with nonviable lipiodolized HCC. Multiphasic contrast-enhanced CT images (A and B), ^{18}F -FDG PET/CT images (C and D).....	115
Figure 5.16 38-year man, who had TACE 1 month before. Contrast-enhanced arterial-phase axial CT image (A, B). The portal phases (C). PET and PET/CT fused images (D, E and F).....	116
Figure 5.17 A 55-year woman suffering from HCC received TACE treatment. (A, B) PET images. (D, E, F) PET/CT fused imaging. Contrast-enhanced CT follow-up after 1 month later (G, H, I).....	117
Figure 5.18 representative images for the three grades of lipiodol deposition in the tumor after TACE	118
Figure 5.19 representative image characteristics of FDG uptake by viable HCC after TACE treatment	118
Figure 5.20 Two lymph node metastatic foci below the left diaphragm and the recurrent lesion in the transplanted liver graft 10 months after liver transplantation using CT, PET, and PET/CT.....	121
Figure 5.21 Metastatic tumor embolism in the inferior vena cava 24 months after liver transplantation using CT, PET, and PET/CT.....	121
Figure 5.22 An intramedullary metastatic focus in the left upper femoral segment 3 months after liver transplantation using CT, PET, and PET/CT.....	122
Figure 5.23 Contrast CT scan showing right lobe and caudate lobe mass and portal vein thrombus in a 55-year-old man. During the arterial phase (A), during the portal phase (B), PET (C) and PET/CT fused images (D).....	124

Figure 5.24 MRI scan showing a left lobe mass and portal vein thrombus in an 80-year-old man (A), CT (C) , PET (B) and PET/CT fused images (D).....125

Figure 5.25 Contrast CT scan showing right lobe and caudate lobe mass and Portal vein thrombus of the right branch in the same patient. During the arterial phase (A), the portal phase (B), PET (C) and PET/CT fused images (D).....125

Figure 5.26 Contrast CT scan showing a diffuse tumor and portal vein thrombus in a 44 year-old man. During the arterial phase (A and C) and PET/CT fused images (B and D).....126

Figure 5.27 Contrast CT scan demonstrating a left lobe mass and portal vein thrombus in a 60-year-old woman. During the arterial phase (A), portal phase (B), PET (C) and PET/CT fused images (D).....126

LIST OF TABLES

Page no.

Table 1.1: The summary of the classifications of the liver segments.....	16
Table 2.1: Relative incidence of primary liver cancer in men in different parts of the world.....	28
Table 2.2: TNM staging of liver tumors.....	39
Table 2.3: The Barcelona Clinic Liver Cancer staging classification....	40
Table 3.1: Commonly used PET scintillator detectors and their properties.....	46
Table 3.2: Radioactive isotopes used in PET.....	55

CONTENTS

	Page
Introduction.....	1
Aim of the Work.....	3
Chapter 1: Anatomy of the Liver.....	4
Chapter 2: Pathology of Hepatocellular carcinoma	27
Chapter 3: Physical Principles of PET/CT.....	41
Chapter 4: Technique of PET/CT examination.....	56
Chapter 5: Diagnostic value of PET/CT in evaluation of HCC.....	91
Summary.....	127
References.....	129
Arabic Summary.....	

INTRODUCTION

Cancer is a major cause of death in the developed world, and is becoming a significant issue for developing countries (*Jones et al., 2006*).

Hepatocellular carcinoma (HCC) is the 5th most common cancer world wide & responsible for up to 1 million deaths annually, its incidence is increasing worldwide because of the dissemination of hepatitis B and C virus infections (*Huang et al., 2009*).

Hepatocellular carcinoma (HCC) is globally the commonest liver primary, and cholangiocarcinoma the second commonest primary liver tumour. Cholangiocarcinoma accounts for 3% of all gastrointestinal cancers. Mesenchymal liver tumours are rare, but include hepatic angiosarcoma and primary hepatic lymphoma (*Vauthey and Blumgart, 2009*).

The most common malignant tumors in the liver are metastases from wide variety of neoplasms, that most frequently are carcinomas from colorectal, breast, and lung primaries. Often discovered as solitary, liver metastases can be effectively treated with surgery (*Arciero and Sigurdson, 2008*).

Surgical treatment ,including hepatic resection and liver transplantation ,are considered as the most effective treatment of HCC. Interventional treatment have been applied to patients with inoperable HCC. Despite initial remission of HCC after surgical and interventional treatment ,recurrence is common. Since patients with recurrent HCC may be amenable to potentially curative resection, early detection of intrahepatic recurrence and /or extra hepatic metastases is extremely important and can facilitate successful retreatment at an early stage (*Sun et al., 2009*).

Modern cross sectional structural imaging techniques like ultrasonography, computed tomography (CT) and magnetic resonance imaging(MRI) provide high resolution images that aid in accurate detection, delineation and anatomic localization of the liver malignancies. However, characterization of lesions into benign and malignant etiologies is often not possible from structural imaging techniques alone. Although functional imaging techniques like positron emission tomography (PET) with radiolabeled ¹⁸F labeled