

Immediate Implant Placement In Fresh Extraction Sockets With Topical Statins And Nano Crystalline Bone Graft

(Clinical and Biochemical study)

Thesis submitted to the faculty of dentistry

Ain Shams University

In partial fulfillment of the requirements of PHD degree in

Oral Medicine , Periodontology and Oral Diagnosis

By

Sherif Abdelrahman Lofty Amer

Teaching Assistant of Oral Medicine, Periodontology

and Oral Diagnosis Department

Faculty of Oral and Dental Medicine

Future University

2018

Supervisors

Prof.Dr. Khaled Atef Abdelghaffar

Minister of higher education and scientific research

Professor Of Oral Medicine,Periodontology

and Oral Diagnosis, Faculty of Dentistry

Ain Shams University

Prof.Dr.Nivine Ibrahim Ragy

Professor of Oral Medicine,Periodontology

and Oral Diagnosis

Vice dean of Faculty of Oral and Dental Medicine

Future University In Egypt

Prof.Dr.Olfat Gamel Shaker

Professor of Medical Biochemistry and Molecular Biology

Faculty of Medicine

Cairo University

Abbreviations

AgP	Aggressive periodontitis
BD	Bone density
BIC	Bone to implant contact
BMC	Bone mineral content
BMD	Bone mineral density
BMP	Bone morphogenic proteins
BV	Bone volume
CAL	Clinical attachment level
CBCT	Cone beam computed tomography
CRP	C-reactive protein
CS	Calcium sulfate
CT	Computed tomography
3D	Three dimensional
DFDBA	Demineralized freeze dried bone allograft
ELISA	Enzyme-linked immunosorbent assay
FOV	Field of view
GBR	Guided bone regeneration
GCF	Gingival crevicular fluid

GI	Gingval index
HA	Hydroxyapetite
HMG	Hydroxy methyl glutaryl
HMGB	High mobility group box
huCRP	Human C-reactive protein
IAN	Inferior alveolar nerve
ICAM	Intracellular adhesion molecule
IL	Interleukin
LDL	Low density lipoprotein
mA	Milliampere
MMP	Matrix metalloproteinase
MPO	Myeloperoxidase
NADPH	Nicotinamide adenine dinucleotide phosphate
NBT	New bone thickness
Ncm	Newton centimeter
nCS	nano calcium sulfate
PBI	Papillary bleeding index
PD	Probing depth
PGA	propylene glycol alginate
PISF	Peri implant sulcular fluid

PLGA	poly(lactic-co-glycolic acid
PTFE	Polytetrafluoroethylene
RA	Rheumatoid arthritis
RANKL	Receptor activator of nuclear factor kappa-B ligand
ROI	Region of interest
SMV	Simvastatin
SRB	Scaling and root planing
TCP	Tri-calcium phosphate
Th-2	T-helper
TMJ	Tempromandibular joint
TNF	Tumor necrosis factor
VEGF	Vascular endothelail growth factor
WKM	Width of keratinized mucosa

List of Figures

<i>No</i>	<i>Heading</i>	<i>Page</i>
		<i>No</i>
<i>1</i>	Molecular structure of simvastatin	<i>44</i>
<i>2</i>	Cholesterol biosynthesis stages and statin action .	<i>46</i>
<i>3</i>	Surgical armamentarium	<i>73</i>
<i>4</i>	NeoBiotic implant kit	<i>73</i>
<i>5</i>	Human IL-13 ELISA kit components	<i>77</i>
<i>6</i>	Periopaper	<i>78</i>
<i>7</i>	Peri-implant sulcular fluid collection	<i>79</i>
<i>8</i>	Buffer concentrations	<i>82</i>
<i>9</i>	Cross sectional view of implant with four levels of bone thickness measurements.	<i>85</i>
<i>10</i>	Gingival cuff after three months	<i>88</i>
<i>11</i>	Cementation of the restoration	<i>88</i>
<i>12</i>	Occlusal view of the tooth to be extracted (upper left 1st premolar) .	<i>90</i>
<i>13</i>	Facial view of the tooth to be extracted	<i>90</i>

14	The use of periotome for severing the gingival and periodontal fibers .	91
15	Caliberated probe is used for measuring the root length .	91
16	The hard and soft tissue condition before osteotomy preparation .	92
17	Sounding of bone with a periodontal probe and detection of the buccal bone level .	92
18	Occlusal view of the implant in place .	93
19	Periodontal probe for measuring the gap distance .	93
20	Addition of nano-crystalline calcium sulfate bone graft .	94
21	Healing abutment connected to the implant and suturing .	94
22	Cross-sectional CBCT view for measuring the buccal bone thickness at baseline.	95
23	Cross-sectional CBCT view for measuring the buccal bone thickness after 6 months.	95
24	Occlusal view of tooth to be extracted (upper left 1st premolar).	96

25	Facial view of the tooth to be extracted	96
26	The use of periosteal elevator for severing the gingival and periodontal fibers.	97
27	Calibrated probe is used for measuring the root length .	97
28	The hard and soft tissue condition before osteotomy preparation.	98
29	Sounding of bone with a periodontal probe and detection of the buccal bone level.	98
30	Measuring of the mesio-distal width with a periodontal probe.	99
31	Occlusal view of the implant in place . A residual gap distance was measured with a periodontal probe.	99
32	Addition of simvastatin gel to fill the gap distance.	100
33	Healing abutment connected to the implant and suturing.	100
34	Cross-sectional CBCT view for measuring the buccal bone thickness at baseline.	101
35	Cross-sectional CBCT view for measuring the buccal bone thickness after 6 months.	101
36	Occlusal view of the tooth to be extracted	102

(upper right 1st premolar).

37	Facial view of the tooth to be extracted	102
38	The use of periosteal elevator for severing the gingival and periodontal fibers .	103
39	The hard and soft tissue condition before osteotomy preparation .	103
40	Calibrated probe is used for measuring the root length.	104
41	Sounding of bone with periodontal probe and detection of the buccal bone level.	104
42	Measuring of the mesio-distal width with a periodontal probe.	105
43	Occlusal view of the implant in place . A gap distance was measured with a probe.	105
44	Healing abutment connected to the implant.	106
45	Simvastatin gel and nano-crystalline calcium sulfate to fill the gap distance.	106
46	Suturing with interrupted sutures	107
47	Cross-sectional CBCT view for measuring the buccal bone thickness at baseline.	107
48	Cross-sectional CBCT view for measuring the buccal bone thickness after 6 months.	108

49	Occlusal view of the tooth to be extracted (upper left 1st premolar).	109
50	The hard and soft tissue condition before osteotomy preparation .	109
51	Caliberated probe is used for measuring the root length.	110
52	Paralleling pin into osteotomy site.	110
53	Occlusal view of the implant in place.	111
54	Healing abutment connected to the implant.	111
55	Cross-sectional CBCT view for measuring the buccal bone thickness at baseline.	112
56	Cross-sectional CBCT view for measuring the buccal bone thickness after 6 months.	112
57	Bar chart showing standard deviation of patients age among the groups .	118
58	Bar chart showing the mean values of (PBI) for all groups over the study follow up period .	122
59	Bar chart showing the mean values of (GI) for all groups over the study follow up period .	125

60	Bar chart showing the mean and standard deviation values of (PD) for all groups over the study follow up period .	129
61	Bar chart showing the mean values of (IL-13) for all groups over the study follow up periods .	135
62	Bar chart showing the mean values of bone thickness for all groups at baseline .	139
63	Bar shart showing the mean values of bone thickness for all groups after 6 months .	146

List Of Tables

<i>No</i>	<i>Heading</i>	<i>Page</i>
------------------	-----------------------	--------------------

	<i>No</i>
1 IL-13 kit components	80
2 Showing the demographic data for group (A)	114
3 Showing the demographic data for group (B)	115
4 Showing the demographic data for group (C)	116
5 Showing the demographic data for group (D)	117
6 Showing the mean and standard deviation (SD) values of patient ages for all groups .	118
7 The mean and standard deviation values of (PBI) for all groups at baseline .	119
8 The mean and standard deviation values of (PBI) for all groups after 3 months .	120
9 The mean and standard deviation values (PBI) for all groups over the study follow up period .	121
10 The mean and standard deviation (SD) values of (GI) for all groups at baseline .	123
11 The mean and standard deviation values of (GI) for all groups over the study follow up period .	124

12	Multiple comparisons of gingival index (GI) between groups after 3 months .	125
13	The mean and standard deviation values of probing depth (PD) for all groups at baseline .	126
14	The mean and standard deviation values of probing depth (PD) for all groups over the study follow up period .	127
15	Multiple comparisons of probing depth (PD) between groups after 3 months .	128
16	The mean and standard deviation values of IL-13 for all groups at baseline	130
17	The mean and standard deviation values of (IL-13) for all groups after 2 weeks .	131
18	Multiple comparisons of (IL-13) between groups after 2 weeks .	131
19	Comparison between mean and standard deviation (SD) values of (IL-13) at baseline and after 2 weeks .	132
20	The mean and standard deviation values of (IL-13) for all groups after 3 months .	133
21	Multiple comparison of (IL-13) for all groups after 3 months .	134
22	Comparison between mean and standard deviation (SD) values of (IL-13) at 2	134

weeks and 3 months .

- | | | |
|-----------|--|------------|
| 23 | The mean and standard deviation values of bone thickness at T(0) for all groups at baseline . | 136 |
| 24 | The mean and standard deviation values of bone thickness at T(1) for all groups at baseline . | 137 |
| 25 | The mean and standard deviation values of bone thickness at T(2) for all groups at baseline . | 138 |
| 26 | The mean and standard deviation values of bone thickness at T(3) for all groups at baseline . | 139 |
| 27 | The mean and standard deviation values of bone thickness at T(0) for all groups after 6 months . | 140 |
| 28 | Multiple comparisons of bone thickness at T(0)for all groups after 6 months . | 141 |
| 29 | The mean and standard deviation values of bone thickness at T(1) for all groups after 6 months . | 142 |
| 30 | Multiple comparisons of bone thickness at T(1) for all groups after 6 months . | 143 |
| 31 | The mean and standard deviation values of bone thickness at T(2) for all groups after 6 months . | 144 |

- 32** Multiple comparisons of bone thickness at T(2)for all groups after 6 months . **145**
- 33** The mean and standard deviation values of bone thickness at T(3) for all groups after 6 months . **146**

