

**Percutaneous Balloon Pulmonary Valvuloplasty
for Pulmonary Valve Stenosis in Neonates, Infants
and Children: 10 Years' experience at Cairo
University Pediatric Hospital**

Thesis subjected to partial fulfillment of Master Degree of Pediatrics

By

Khaled Mohie El-din Saad, M.B., B.C

Under supervision of

Prof. Dr. Sonia Ali El-Saiedi

Professor of Pediatrics

Cairo University

Prof. Dr. Amr Mohamed Malash

Professor of Pediatrics

Cairo University

Dr. Doaa Mohamed Abd El-Aziz

Lecturer of Pediatrics

Cairo University

Faculty of medicine

Cairo University

2015

ABSTRACT

Background: Balloon pulmonary valvuloplasty (BPV) is the treatment of choice for patients with pulmonary valve stenosis (PS). **Methods:** An observational retrospective cross sectional study was done on cases underwent BPV from 2004-2013 in cardiac cath unit- pediatric cardiology department- Cairo university. The registry was queried for cases of isolated BPV. Multivariable models were built to report the predictors of outcome of balloon pulmonary valvuloplasty and its complications both during and early after the intervention. **Results:** 1200 cases were included in the study and subdivided into age groups (Neonates: 282 cases, Infants: 362 cases and Children: 556 cases). Procedural success was achieved in 82.7% procedures, being defined as drop of right ventricular systolic pressure to less than or equal to 50% of the baseline measurements. Procedural success was more common in children age group (84.5%) compared to other age groups. Among the unsuccessful group which represents 17.3% of the study group, 13.7 % underwent redo of BPV while 3.6% undergone surgery. Complications observed in 8.5% of cases being the commonest in the neonatal age group with 21.6%. **Conclusions:** BPV is the treatment of choice for pulmonary valve stenosis in all age groups and the immediate post-BPV pressure gradient drop was the most important predictor of success of BPV.

Key words:

(Valvular Pulmonary stenosis, Balloon valvuloplasty, Redo, Surgical valvotomy).

DEDICATION

I would like to dedicate this Master dissertation to **my wife and my family** Record. There is no doubt in my mind that without their continued support and counsel I could not have completed this process.

I would like to acknowledge the inspirational instruction and guidance of **Prof .Dr. Sonia Ali El-Saiedi** and **Prof .Dr. Amr Malash** with the great efforts and assistance of **Dr. Doaa Abd El-Aziz**. They have given me a deep appreciation, feedback and advice.

CONTENTS

	Page
List of Abbreviations	4
List of Tables	6
List of Figures	10
Introduction and aim of work	13
Review of literature	14
Patients and methods	71
Results	81
Discussion	106
Summary and conclusion	122
Recommendations	126
References	127
Arabic summary	144

LIST OF ABBREVIATIONS

ACT: Activated Clotting Time.

ASD: Atrial Septal Defect.

Bal: Balloon.

BPV: Balloon Pulmonary Valvuloplasty.

Cath: Catheterization.

CC: Cubic Centimeter.

CT: Computed Tomography.

Echo: Echocardiography.

IM: Intra-muscular.

IV: Intra-venous.

IU: International Unit.

kg: Kilogram.

LPA: Left Pulmonary Artery.

Max: Maximum.

mg: Milligram.

min: Minute.

mm: Millimeter.

mmHg: Millimeters of Mercury.

MR: Mitral Regurge.

MRI: Magnetic Resonance Imaging.

N: Number.

PA: Pulmonary Artery.

PAP: Pulmonary Artery Pressure.

PASP: Pulmonary Artery Systolic Pressure.

PDA: Patent Ductus Arteriosus.

PFO: Patent Foramen Ovale.

PG: Pressure Gradient.

PO: Per Oral.

PR: Pulmonary Regurge.

PS: Pulmonary Stenosis.

PV: Pulmonary Valve.

PVS: Pulmonary Valve Stenois.

RA: Right Atrium.

RPA: Right Pulmonary Artery.

RV: Right Ventricle.

RVP: Right Ventricular Pressure.

RVSP: Right Ventricular Systolic Pressure.

SD: Standard Deviation.

TR: Tricuspid Regurge.

TV: Tricuspid Valve.

VSD: Ventricular Septal Defect.

LIST OF TABLES	Page
Table (1): Pre catheterization feeding order.	52
Table (2): The distribution of patients among the study period.	81
Table (3): Distribution of studied patients in each age group.	83
Table (4): Echocardiographic findings associated with valvular pulmonary stenosis in the studied group before balloon pulmonary valvuloplasty.	85
Table (5): Echocardiographic findings of pulmonary valve annulus and pressure gradient across the pulmonary valve before BPV in each age group.	86

Table (6): Associated cardiac anomalies obtained by echocardiography before BPV of the studied group.	87
Table (7): Description of pressures obtained by catheterization in each age group.	88
Table (8): Description of pulmonary valve annulus in each age group.	89
Table (9): The difference of the values of diameter of the pulmonary valve annulus obtained by echo and cath before balloon dilatation in each age group.	90
Table (10): Correlation between pressure gradient before BPV by echocardiography and cath.	91
Table (11): Correlation between the body weight and balloon size in each age group.	92
Table (12): Correlation between balloon size and	92

pulmonary valve annulus by cath in each age group.	
Table (13): Descriptive data of the balloon used in (BPV) in each age group.	93
Table (14): Distribution of successful cases of (BPV) in each age group.	94
Table (15): The details of the pressure gradient difference before and after balloon dilatation in each age group.	95
Table (16): Distribution of redo cases in each age group.	97
Table (17): Distribution of surgery cases in each age group.	98
Table (18): Descriptive data of complications during or immediately after (BPV) in each age group.	100
Table (19): Variables measured by cath. and	102

echocardiography and their correlation to the risk of redo of BPV.	
Table (20): Variables measured by cath. and echocardiography and their correlation to the risk of surgery after BPV.	103
Table (21): The studied variables measured by cath and their association with the risk of redo of BPV.	104
Table (22): The studied variables measured by cath and their association with surgery.	104

LIST OF FIGURES	Page
Figure (1): Selected cine frames of two balloon catheters placed across the pulmonary valve.	62
Figure (2): Selected cineradiographic frames of a balloon dilatation catheter placed across a stenotic pulmonary valve.	67
Figure (3): Selected frames from lateral views of right ventricular (RV) cineangiograms before (A)and after (B) balloon pulmonary valvuloplasty.	69
Figure (4): Distribution of patients within the study period.	82
Figure (5): Age groups of the studied patients.	84

Figure (6): The percent of patients with almost similar values of the pulmonary valve annulus by Echo and angiography.	90
Figure (7): Distribution of successful cases of (BPV) in each age group.	95
Figure (8): The details of the pressure gradient difference before and after BPV.	96
Figure (9): Distribution of redo cases in each age group.	98
Figure (10): Distribution of surgery cases in each age group.	99
Figure (11): Descriptive data of complications during	101

or immediately after (BPV) in each age group.	
---	--

INTRODUCTION AND AIM OF WORK

INTRODUCTION AND AIM OF WORK

The traditional method of treatment for congenital pulmonic valve stenosis was surgical valvotomy until 1982, when **Kan et al, (1982)** introduced the technique of percutaneous balloon pulmonary valvuloplasty. The short-term (**Kan et al, 1984; Rocchini et al, 1984; Khan et al, 1986; Piechaud et al, 1987; Rey et al, 1988; Santro et al, 1988; Schmaltz et al, 1989; Stanger et al, 1990; and Maalouk et al, 1995**) and mid-term (**Al Kasab et al, 1988; Mullins et al, 1988; Rao et al, 1988; O'Connor, 1992 and Shrivastava et al, 1993**). Results of balloon pulmonary valvuloplasty have been so good that nowadays it has become the preferred method of therapy for moderate to severe pulmonary valve stenosis in children and adults. However, data documenting long term effectiveness of this procedure are scarce (**Mc Crindle et al, 1991; Masura et al, 1993 and Mc Crindle et al, 1994**).

The aim of this work is to determine the outcome of trans-catheter balloon dilatation of the pulmonary valve in children with pulmonary valve stenosis with a wide range in age and severity, also to report the predictors of outcome of balloon pulmonary valvuloplasty and its complications both during and early after the intervention.