



Different Types of Graph Labelings

A thesis
submitted for the Ph.D. degree of pure mathematics
as a fulfillment for requirements of
the Ph.D. degree in pure mathematics

Submitted to

Mathematics Department

Faculty of Science

Ain Shams University

Cairo Egypt

2013

By:

Maher Abdulkader Salim

(M.Sc. Degree in Mathematics - Ain Shams University 2009)

Supervised By:

Professor Mohammed Abd-El Azim Seoud

Professor of Pure Mathematics

Mathematics Department

Faculty of Science

Ain Shams University

Dr. Alaa E.I. Abdel Maqsoud

Lecturer of Pure Mathematics

Mathematics Department

Faculty of Science

Ain Shams University

ACKNOWLEDGMENT

*First and Foremost, Thanks are to **GOD**, The Most Beneficent and Merciful.*

*Since the success of any project depends largely on the encouragement and guidelines of others. I take this opportunity to express my sincere gratitude to the person who has been instrumental in the successful completion of this dissertation ; I would like to show my greatest appreciation to **Prof. Dr. Mohammed Abdel Azim Seoud**, Professor of Pure Mathematics Department, Faculty of Science, Ain Shams University, I can't say thank you enough for his fruitful supervision, tremendous support and continuous help. I feel motivated and encouraged every time I attend his meeting. Without his encouragement and guidance this dissertation would not have materialized.*

*Thanks and gratitude to **Dr. Alaa El-din Ismail Mohamed Abdel Maqsoud**, Teacher of Pure Mathematics Department, Faculty of Science, Ain Shams University, for his continuous encouragement.*

With great respect I would like to say thanks to every member in Mathematics department, I really feel a proud to achive my thesis in this department, college and university.

كلمة شكر

أشكر الله سبحانه وتعالى على توفيقى فى إتمام هذه العمل وأسأله أن يكون فى ميزان حسناتى.

أعنتم هذه الفرصة للتعبير عن خالص امتنانى للمشرف الرئيس الأستاذ الدكتور محمد عبد العظيم سعود على تفضله بالإشراف علىّ إشرافاً مثمراً أغنى الرسالة وأغنى عقلية إنجاز البحث العلمى لىّ ، فلقد كان نعم المعلم و الناصح و الناقد و المصوّب أثناء إعداد الرسالة و كان بجانبى وساندى طوال الوقت ولقد أفدت من علمه وأدبه.

كما أشكر المشرف الثانى الدكتور علاء عبد المقصود على تشجيعه الدائم.

وأوجه بالشكر موصولاً لأبى وأمى لأ نهما السبب فى اىصالى إلى هذه المرتبة. وأشكر زوجتى لصبرها على الغربة وعونها الدائم لى ، كذلك أشكر ولدىّ وقرّة عيني عبد القادر واللىث على تحملهما انشغالى عنهما.

Table of contents

Chapter One Introduction.....	3
1.1 Some fundamentals in graph theory.....	3
1.1.1 Definitions.....	3
1.1.2 Some important types of graphs	7
1.1.3 Definitions of two operations on graphs	9
1.2 Introduction to graph labeling.....	10
Chapter Two On Odd Mean Graphs	17
2.1 Introduction:	17
2.2 Preliminary results	19
2.3 Two necessary conditions.....	20
2.4 All odd mean graphs of order ≤ 6	26
Chapter Three Further Results On Triangular Sum Graphs	35
3.1 Introduction	35
3.2 Triangular sum graphs.....	36
3.3 Embedding graphs as induced subgraphs of triangular sum graphs.	46
Chapter Four Upper Bounds Of Four Types Of Graph Labelings.....	55
4.1 Introduction	55
4.2 Prime cordial labeling	56
4.3 k -equitable graph labeling.....	59
4.4 (k, d)-odd mean labeling	61
4.5 Z₂⊕Z₂ -cordial labeling	64
Chapter Five Further Results On Edge - Odd Graceful Graphs.....	67
5.1 Introduction:	67
5.2 Main results:	68
5.3 Two theorems of non edge - odd graceful graphs.....	78
5.4 Some edge - odd graceful complete graphs.	79
Chapter Six Some Results And Remarks On Harmonic Mean And Mean Cordial Labelings.....	83
6.1 Introduction	83
6.2 Harmonic mean labeling	84
6.3 Mean cordial labeling.....	93
References.....	101

Chapter One

Introduction

1.1 Some fundamentals in graph theory:

1.1.1 Definitions[30]:

A **simple graph** G consists of a non-empty finite set $V(G)$ of elements called **vertices** (or nodes), and a finite set $E(G)$ of distinct unordered pairs of distinct elements of $V(G)$ called **edges**. We call $V(G)$ the **vertex set** and $E(G)$ the **edge set** of G . An edge $\{v, w\}$ is said to join the vertices v and w , and is usually abbreviated to vw . For example, *Figure 1.1* represents the simple graph G whose vertex set $V(G)$ is $\{u, v, w, z\}$, and whose edge set $E(G)$ consists of the edges uv, uw, vw and wz .




Figure 1.1

In any simple graph there is at most one edge joining a given pair of vertices. However, many results that hold for simple

graphs can be extended to more general objects in which two vertices may have several edges joining them. In addition, we may remove the restriction that an edge joins two *distinct* vertices, and allow loops -edges joining a vertex to itself.

The resulting object, in which loops and multiple edges-are allowed, is called a general graph - or, simply, a graph. Thus every simple graph is a graph, but not every graph is a simple graph.

Anyway through chapters 2,3,4 and 5, wherever we say “**graph**” we mean “**simple graph**”.

Also it is essential to mention that we refer to a graph G as $G(V(G), E(G))$ or $G(p, q)$, where

$p = |V(G)|$ (= **order of G**), $q = |E(G)|$ (= **size of G**).

Isomorphism

Two general graphs G_1 and G_2 are **isomorphic** if there is a one-one correspondence between the vertices of G_1 and those of G_2 such that the number of edges joining any two vertices of G_1 , is equal to the number of edges joining the corresponding vertices of G_2 . Thus the two graphs shown in Figure 1.2 are isomorphic under the correspondence $u \leftrightarrow l, v \leftrightarrow m, w \leftrightarrow n, x \leftrightarrow p, y \leftrightarrow q, z \leftrightarrow r$




Figure 1.2

Connectedness

We can combine two graphs to make a larger graph. If the two graphs are $G_1 = (V(G_1), E(G_1))$ and $G_2 = (V(G_2), E(G_2))$, where $V(G_1)$ and $V(G_2)$ are disjoint, then their **union** $G_1 \cup G_2$ is the graph with vertex set $V(G_1) \cup V(G_2)$ and edge family $E(G_1) \cup E(G_2)$ (see Figure 1.3).

Thus a graph is **connected** if it cannot be expressed as the union of two graphs, and **disconnected** otherwise. Clearly any **disconnected** graph G can be expressed as the union of connected graphs.




Figure 1.3

Adjacency

We say that two vertices v and w of a graph G are **adjacent** if there is an edge vw joining them, and the vertices v and w are then **incident** with such an edge. Similarly, two distinct edges e and f are **adjacent** if they have a vertex in common (see Figure 1.4).




Figure 1.4

The **degree** of a vertex v of G is the number of edges incident with v , and is written $\deg(v)$; in calculating the degree of v , we usually make the convention that a loop at v contributes 2 (rather than 1) to the degree of v . A vertex of degree 0 is an **isolated vertex** and a vertex of degree 1 is an **end-vertex**.

Note: *in any graph the sum of all the vertex-degrees is an even number* - in fact, twice the number of edges, since each edge contributes exactly 2 to the sum. This result is called the **handshaking lemma**. An immediate corollary of the handshaking lemma is that *in any graph the number of vertices of odd degree is even*.

Subgraphs

A **subgraph** of a graph G is a graph, each of whose vertices belongs to $V(G)$ and each of whose edges belongs to $E(G)$. Thus in *Figure 1.5* the graph (a) is a subgraph of the graph (b), but is not a subgraph of the graph (c), since the latter graph contains no 'triangle = C_3 '.




Figure 1.5

1.1.2 Some important types of graphs[30]

Null graphs

A graph whose edge-set is empty is a null graph. We denote the null graph on n vertices, (of “order” n), by N_n or \bar{K}_n .

Complete graphs

A simple graph in which each pair of distinct vertices are adjacent is a complete graph. We denote the complete graph of order n by K_n . Realize that K_n has $\binom{n}{2} = n(n-1)/2$ edges.

Cycle graphs and path graphs

A connected graph that is regular of degree 2 is a cycle graph.

We denote the cycle graph of order n by C_n . The graph obtained from C_n by removing an edge is the path graph of order n , denoted by P_n . The graphs C_6 and P_6 are shown in *Figure 1.6*. C_3 is called **Triangle**.




Figure 1.6

Bipartite graphs

If the vertex set of a graph G can be split into two disjoint sets A and B so that each edge of G joins a vertex of A and a vertex of B , then G is a **bipartite graph**. Alternatively, a bipartite graph is one whose vertices can be coloured black and white in such a way that each edge joins a black vertex (in A) and a white vertex (in B). A **complete bipartite graph** is a bipartite graph in which each vertex in A is joined to each vertex in B by just one edge. We denote the bipartite graph with r black vertices and s white vertices by $K_{r,s}$; $K_{1,3}$, $K_{2,3}$ and $K_{3,3}$ are shown in *Figure 1.7*. Note that $K_{r,s}$ has $r + s$ vertices and rs edges.




Figure 1.7

It is useful here to mention the lemma stating that every bipartite graph contains no **odd cycles** (= cycles of odd order).

As a similar argument of a bipartite graph, a graph G is **k -partite graph**, $k \geq 3$, if it is possible to partition $V(G)$ into k subsets V_1, V_2, \dots, V_k (called partite sets) such that every element of $E(G)$ joins a vertex of V_i to a vertex of $V_j, i \neq j$.

A **complete k -partite graph** G is a k -partite graph with partite sets V_1, V_2, \dots, V_k , having the added property that if $u \in V_i$ and $v \in V_j, i \neq j$, then $uv \in E(G)$. If $|V_i| = n_i$, then this graph is denoted by K_{n_1, n_2, \dots, n_k} .

Trees

A **forest** is a graph that contains no cycle and a connected forest is a **tree**.

The complement of a simple graph

If G is a simple graph with vertex set $V(G)$, its **complement** \bar{G} is the simple graph with vertex set $V(G)$ in which two vertices are adjacent if and only if they are *not* adjacent in G .

1.1.3 Definitions of two operations on graphs[6]

- 1) The **corona** $G_1 \odot G_2$ of two graphs G_1 and G_2 was defined by Frucht and Harary as the graph H obtained by taking one copy of G_1 (which has p vertices) and p copies of G_2 , and then joining the i^{th} vertex of G_1 to every vertex

in the i^{th} copy of G_2 . (See Figure 1.8).




Figure 1.8 Two graphs and their coronas

2) The **Cartesian product** $G = G_1 \times G_2$ has $V(G) = V(G_1) \times V(G_2)$, and two vertices (u_1, u_2) and (v_1, v_2) of G are adjacent if and only if either

$u_1 = v_1$ and $u_2 v_2 \in E(G_2)$ or

$u_2 = v_2$ and $u_1 v_1 \in E(G_1)$. See Figure 1.9




Figure 1.9 The product of two graphs

1.2 Introduction to graph labeling

A graph labeling is an assignment of integers to the vertices or edges, or both, subject to certain conditions.

Most graph labeling methods trace their origin to one introduced by Rosa [15] in 1967, or one given by Graham and Sloane [5] in 1980. Rosa [15] called a function f a β -valuation of a graph G with q edges if f is an injection from the vertices of G to the set $\{0, 1, \dots, q\}$ such that, when each edge xy is assigned the label $|f(x) - f(y)|$, the resulting edge labels are distinct. Golomb [4] subsequently called such labelings graceful and this is now the popular term. Rosa introduced β -valuations as well as a number of other labelings as tools for decomposing the complete graph into isomorphic subgraphs. In particular, β -valuations originated as a means of attacking the conjecture of Ringel [14] that K_{2n+1} can be decomposed into $2n + 1$ subgraphs that are all isomorphic to a given tree with n edges. Although an unpublished result of Erdős says that most graphs are not graceful (cf. [5]), most graphs that have some sort of regularity of structure are graceful. Sheppard [26] has shown that there are exactly $q!$ gracefully labeled graphs with q edges. Balakrishnan and Sampathkumar [1] have shown that every graph is a subgraph of a graceful graph. Rosa [15] has identified essentially three reasons why a graph fails to be graceful: (1) G has “too many vertices” and “not enough edges,” (2) G “has too many edges,” and (3) G “has the wrong parity.” An infinite class of graphs that are not graceful for the second reason is given in [2]. As an example of the third

condition Rosa [15] has shown that if every vertex has even degree and the number of edges is congruent to 1 or 2 (mod 4) then the graph is not graceful. In particular, the cycles C_{4n+1} and C_{4n+2} are not graceful.

Seoud and Wilson [19] have shown that $C_3 \cup K_4$, $C_3 \cup C_3 \cup K_4$, and certain graphs of the form $C_3 \cup P_n$ and $C_3 \cup C_3 \cup P_n$ are not graceful. Seoud and Youssef [22] investigated the gracefulness of specific families of the form $G \cup K_{m,n}$. They obtained the following results: $C_3 \cup K_{m,n}$ is graceful if and only if $m \geq 2$ and $n \geq 2$; $C_4 \cup K_{m,n}$ is graceful if and only if $(m,n) \neq (1,1)$; $C_7 \cup K_{m,n}$ and $C_8 \cup K_{m,n}$ are graceful for all m and n ; $mK_3 \cup nK_{1,r}$ is not graceful for all m, n and r ; $K_i \cup K_{m,n}$ is graceful for $i \leq 4$ and $m \geq 2$; $n \geq 2$ except for $i = 2$ and $(m,n) = (2,2)$; $K_5 \cup K_{1,n}$ is graceful for all n ; $K_6 \cup K_{1,n}$ is graceful if and only if n is not 1 or 3.

Harmonious graphs naturally arose in the study by Graham and Sloane [5] of modular versions of additive bases problems stemming from error-correcting codes. They defined a graph G with q edges to be harmonious if there is an injection f from the vertices of G to the group of integers modulo q such that when each edge xy is assigned the label $f(x) + f(y) \pmod{q}$, the resulting edge labels are distinct. When G is a tree, exactly one label may be used on two vertices. Analogous to

the “parity” necessity condition for graceful graphs, Graham and Sloane proved that if a harmonious graph has an even number q of edges and the degree of every vertex is divisible by 2^k then q is divisible by 2^{k+1} . Liu and Zhang [12] have generalized this condition as follows: if a harmonious graph with q edges has degree sequence d_1, d_2, \dots, d_p then $\gcd(d_1, d_2, \dots, d_p, q)$ divides $q(q - 1)/2$. They have also proved that every graph is a subgraph of a harmonious graph. Determining whether a graph has a harmonious labeling was shown to be NP-complete by Auparajita, Dulawat, and Rathore in 2001 (see [10]).

Seoud and Youssef [25] define a *closed helm* as the graph obtained from a helm by joining each pendant vertex to form a cycle and a *flower* as the graph obtained from a helm by joining each pendant vertex to the central vertex of the helm. They prove that closed helms and flowers are harmonious when the cycles are odd. A *gear graph* is obtained from the wheel W_n by adding a vertex between every pair of adjacent vertices of the n -cycle. Seoud and Youssef [23] have shown that the one point union of a triangle and C_n is harmonious if and only if $n \equiv 1 \pmod{4}$ and that if the one-point union of two cycles is harmonious then the number of edges is divisible by 4. The question of whether this latter condition is sufficient is open. Seoud and Youssef [24] proved that $P_n \oplus \overline{K_2}$ is graceful