



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



# شبكة المعلومات الجامعية

## التوثيق الالكتروني والميكرو فيلم

# جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

## قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها  
علي هذه الأفلام قد اعدت دون أية تغيرات



## يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of  
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



# بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات  
لم ترد بالأصل



# **Value Of Polymerase Chain Reaction In The Diagnosis Of Candida Endophthalmitis**

Thesis submitted in partial fulfillment of MD degree of  
Ophthalmology

By  
**Ahmed Fathy Gabr**  
MSc Ophthalmology



## **Supervisors**

**Prof. Dr. Aida Ali Hussein**  
Professor of Ophthalmology  
Benha Faculty of Medicine

**Prof. Dr. Wahid Mahmoud Yousef**  
Professor of Ophthalmology  
Benha Faculty of Medicine

**Dr Sawsan Abdul Rahman Yousef**  
Assist. Prof. of Microbiology  
Benha Faculty of Medicine

**Dr. Amr Ibrahim Sharawy**  
Assist. Prof. of Ophthalmology  
Benha Faculty of Medicine

B 1-10

**Faculty of Medicine  
Benha University  
2006**

بسم الله الرحمن الرحيم  
قالوا سبحانك لا علم لنا إلا ما علمتنا  
إنك أنت العليم الحكيم

صدق الله العظيم

## ***Acknowledgement***

I would like to thank Professor Dr. Aida Ali Hussein, Professor of Ophthalmology, Benha Faculty of Medicine, to whom my gratitude is beyond expression for her continuous, faithful, and valuable support.

I owe so much to Professor Dr. Wahid Mahmoud Yousef, Professor of Ophthalmology, Benha Faculty of Medicine, for his constant advice and encouragement.

I want to express my gratitude to Dr. Sawsan Abdul Rahman Yousef, Assist. Prof. of Microbiology, Benha Faculty of Medicine, for her great help and honest advice.

I'm grateful also to Dr Amr Ibrahim Sharawy, Assist. Prof. of Ophthalmology, Benha Faculty of Medicine, for his enormous help and support.

Finally, my thanks to all staff-members of departments of Ophthalmology and Microbiology, Benha Faculty of Medicine, and to all those who helped me in preparing this study.



# **Contents**

<b>1- Introduction</b>	<b>(1)</b>
<b>2- Aim of the work</b>	<b>(3)</b>
<b>3- Review of literature</b>	<b>(4)</b>
• Historical review	(4)
• Classification of endophthalmitis	(5)
• Incidence of fungal endophthalmitis	(5)
• Pathology of fungal endophthalmitis	(6)
• Clinical settings in fungal endophthamitis	(9)
• Complications of fungal endophthalmitis	(29)
• Diagnostic techniques in fungal endophthalmitis	(30)
• Polymerase chain reaction (PCR) assay	(40)
<b>4- Material and methods</b>	<b>(50)</b>
• Preparation of the animal	(50)
• Preparation of the fungus suspension	(50)
• Test of viability	(50)
• Inoculation of the fungus	(50)
• Follow up	(51)
Collection of intraocular samples	<b>(54)</b>
Microscopic examination by Gram stain	<b>(56)</b>
Culture on Sabouraud agar	<b>(56)</b>
Polymerase chain reaction	<b>(57)</b>

<b>5- Results</b>	(62)
• Gram stain	(68)
• Culture on Sabouraud's agar	(72)
• Polymerase chain reaction assay	(77)
<b>6- Case reports</b>	(83)
<b>7- Discussion</b>	(93)
<b>8- Conclusion and Summary</b>	(103)
<b>9- References</b>	(105)
<b>10- Arabic summary</b>	(127)

## **List of figures**

<b>Fig. (1) :</b>	<b>Multifocal Candida retinochoroiditis with extension into the vitreous</b>	<b>(22)</b>
<b>Fig. (2) :</b>	<b>Aspergillus endophthalmitis</b>	<b>(22)</b>
<b>Fig. (3) :</b>	<b>Schematic of the mechanism of amplification of genomic DNA (A&amp;B).</b>	<b>(41)</b>
<b>Fig. (4) :</b>	<b>Schematic of the mechanism of amplification of genomic DNA (C).</b>	<b>(42)</b>
<b>Fig. (5) :</b>	<b>Candida albicans colonies on Sabouraud agar during the test of viability.</b>	<b>(53)</b>
<b>Fig. (6) :</b>	<b>Collection of aqueous humor sample from an infected eye.</b>	<b>(53)</b>
<b>Fig. (7) :</b>	<b>Collection of vitreous fluid sample from an infected eye.</b>	<b>(55)</b>
<b>Fig. (8) :</b>	<b>Eppendorf tube.</b>	<b>(55)</b>
<b>Fig. (9) :</b>	<b>Safety cabinet (LABCONCO) containing micropipettes, repeater, Eppendorf tubes, and minicentrifuge.</b>	<b>(58)</b>
<b>Fig.(10) :</b>	<b>Thermal controller for the Real-time PCR.</b>	<b>(58)</b>
<b>Fig. (11):</b>	<b>Electrophoresis unit for agarose gel.</b>	<b>(60)</b>
<b>Fig. (12):</b>	<b>Ultraviolet transilluminator.</b>	<b>(60)</b>
<b>Fig. (13):</b>	<b>Infected eye of rabbit No 2 three days postinoculation showing conjunctival hyperemia and vitreous reaction.</b>	<b>(63)</b>
<b>Fig. (14):</b>	<b>Infected eye of rabbit No 3 three days postinoculation showing conjunctival edema. Posterior synechiae, and vitritis.</b>	<b>(63)</b>
<b>Fig. (15):</b>	<b>Infected eye of rabbit No 6 ten days postinoculation showing severe conjunctival injection and exudation, severe corneal edema, and hypopyon.</b>	<b>(64)</b>
<b>Fig. (16):</b>	<b>Infected eye of rabbit No 13 fourteen days postinoculation showing severe conjunctival injection, severe corneal edema and vascularization, with masking of anterior chamber details.</b>	<b>(64)</b>
<b>Fig. (17):</b>	<b>Infected eye of rabbit No 8 fourteen days postinoculation showing severe conjunctival injection, corneal edema, vascularization and central corneal melting.</b>	<b>(65)</b>



- Fig. (18):** Infected eye of rabbit No 9 ten days postinoculation showing severe conjunctival injection and discharge, severe corneal edema, hypopyon, and loss of red reflex. (65)
- Fig. (19):** Fundus photography of infected eye of rabbit No 6 three days postinoculation showing vitreous reaction partially masking fundus details. (66)
- Fig. (20):** Infected eye of rabbit No 5 four days postinoculation showing vitreous (cotton balls) reaction. (66)
- Fig. (21):** Combined A and B scan of infected eye of rabbit No 13 eight days postinoculation showing moderate vitreous opacities and membranes. (67)
- Fig. (22):** Combined A and B scan of infected eye of rabbit No 17 twelve days postinoculation showing dense vitreous opacities. (67)
- Fig. (23):** Infected eye of rabbit No 1 second day postinoculation showing injury of the posterior lens capsule. (69)
- Fig. (24):** Infected eye of rabbit No 19 four days postinoculation showing vitreous reaction and vitreous hemorrhage. (69)
- Fig. (25):** Gram stain for vitreous fluid sample from infected eye of rabbit No. 8 showing *Candida albicans* as rounded oval, Gram positive organism. (70)
- Fig. (26):** Gram stain for *Candida albicans* obtained from culture on Sabouraud's agar. (70)
- Fig. (27):** Culture on Sabouraud's agar five days postinoculation showing dome-shaped, creamy white colonies of *Candida albicans*. (73)
- Fig. (28):** The mean duration required for growth of *Candida albicans* on Sabouraud's agar (early and late samples). (76)
- Fig. (29):** PCR results of samples from infected eye of rabbit No 1 showing the amplified products after agarose gel electrophoresis using panfungal primer. (78)
- Fig. (30):** PCR results of samples from control eye of rabbit No 19 showing the amplified products after agarose gel electrophoresis using panfungal primer. (78)
- Fig. (31):** PCR results of vitreous samples from infected eyes showing the amplified products after agarose gel electrophoresis using *Candida* specific primer. (80)
- Fig. (32):** PCR results of samples from control eye of rabbit No 5 showing the amplified products after agarose gel electrophoresis using panfungal primer. (80)

<b>Fig. (33):</b>	<b>PCR results of samples from infected eye of rabbit No 21 showing the amplified products after agarose gel electrophoresis using panfungal primer.</b>	<b>(81)</b>
<b>Fig. (34):</b>	<b>Comparison between average sensitivity of PCR, Gram stain, and culture technique for aqueous humor samples.</b>	<b>(82)</b>
<b>Fig. (35):</b>	<b>Comparison between average sensitivity of PCR, Gram stain, and culture technique for vitreous fluid samples.</b>	<b>(82)</b>
<b>Fig. (36):</b>	<b>Severe vitreous reaction with no red reflex at start of pars plana vitrectomy.</b>	<b>(84)</b>
<b>Fig. (37):</b>	<b>Area of chorioretinitis and retinal deposits during vitrectomy.</b>	<b>(84)</b>
<b>Fig. (38):</b>	<b>Pallor of optic disk with sheathing of blood vessels.</b>	<b>(86)</b>
<b>Fig. (39):</b>	<b>Culture of vitreous fluid sample on Sabouraud agar showing <i>Aspergillus niger</i> hyphae with small blackish spores at the center.</b>	<b>(86)</b>
<b>Fig. (40):</b>	<b>PCR results of vitreous sample of patient No I showing the amplified products after agarose gel electrophoresis.</b>	<b>(87)</b>
<b>Fig. (41):</b>	<b>Intraoperative picture showing severe vitreous reaction (cotton balls) at the lower part of the vitreous.</b>	<b>(89)</b>
<b>Fig. (42):</b>	<b>Cotton balls reaction and retinal deposits at the posterior pole.</b>	<b>(89)</b>
<b>Fig. (43):</b>	<b>Immediate postoperative picture showing sutured corneal wound, iris laceration, and completely clear media.</b>	<b>(91)</b>
<b>Fig. (44):</b>	<b>Fundus photography of silicon filled eye 2 months postoperatively showing resolution of infection and laser scars at the lower retina.</b>	<b>(91)</b>
<b>Fig. (45):</b>	<b>PCR results of vitreous sample of patient No II showing the amplified products after agarose gel electrophoresis.</b>	<b>(92)</b>
<b>Fig. (46):</b>	<b>PCR results of vitreous sample of patient No II showing the amplified products after agarose gel electrophoresis.</b>	<b>(92)</b>



## List of tables

Tab. (1) :	Follow up sheet.	(52)
Tab. (2) :	Results of Gram stain for aqueous humor samples two days after appearance of signs of inflammation.	(68)
Tab. (3) :	Results of Gram stain for vitreous fluid samples two days after appearance of signs of inflammation.	(71)
Tab. (4) :	Results of Gram stain for aqueous humor samples two weeks after appearance of signs of inflammation.	(71)
Tab. (5) :	Results of Gram stain for vitreous fluid samples two weeks after appearance of signs of inflammation.	(72)
Tab. (6) :	Results of culture for aqueous humor samples two days after appearance of signs of inflammation.	(72)
Tab. (7) :	(7): Results of culture for vitreous fluid samples two days after appearance of signs of inflammation.	(73)
Tab. (8) :	Results of culture for aqueous humor samples two weeks after appearance of signs of inflammation.	(74)
Tab. (9) :	Results of culture for vitreous fluid samples two weeks after appearance of signs of inflammation.	(74)
Tab. (10):	The mean duration needed to report culture as positive for aqueous humor samples.	(75)
Tab. (11):	The mean duration needed to report culture as positive for vitreous fluid samples.	(75)
Tab. (12):	The mean duration required for growth of <i>Candida albicans</i> on Sabouraud's agar from early and late samples.	(75)
Tab. (13):	Results of polymerase chain reaction for aqueous humor samples two days after appearance of signs of inflammation.	(77)
Tab. (14):	Results of polymerase chain reaction for vitreous fluid samples two days after appearance of signs of inflammation.	(77)
Tab. (15):	Results of polymerase chain reaction for aqueous humor samples two weeks after appearance of signs of inflammation.	(79)
Tab. (16):	Results of polymerase chain reaction for vitreous fluid samples two weeks after appearance of signs of inflammation.	(79)

## Abbreviations

AH	Aqueous humor
AMB	Amphotericin B
bp	Base pairs
Can-tec	Candida detection system
CBC	Complete blood count
CFU	Colony Forming Unit
CNV	Choroidal neovascularization
CSF	Cerebrospinal fluid
DNA	Deoxyribonucleic Acid
EDTA	Ethylenediamine tetraacetic acid
ERG	Electroretinography
ESR	Erythrocyte sedimentation rate
EVS	The Endophthalmitis Vitrectomy Study
fg	Femtogram
HIV	Human Immunodeficiency Virus
IOLs	Intraocular lenses
ITS	Internal transcribed spacer
L-AMB	Liposomal amphotericin B
LY121019	Cilofungin
LY303366	Terphenyl-substituted echinocandin B
MICs	Minimal Inhibitory Concentrations
MIC <sub>90</sub>	Minimum inhibitory concentrations of 90% of isolates
MK-0991	Caspofungin acetate
mM	milliMole
ng	Nanogram