Dermoscopy Aids in the Diagnosis of Erythematous Scaly Patches and Plaques

Thesis

Submitted for partial fulfillment of Master Degree in Dermatology, Venereology and Andrology

By

Sara Mohammed Mahfouz Mohammed

M.B., B.Ch.

Faculty of Medicine – Ain Shams University

Under Supervision of

Prof. Dr. Hoda Ahmed M. Moneib

Professor of Dermatology, Venereology and Andrology Faculty of Medicine – Ain Shams University

Dr. Al Hassan M. El Hefnawy

Lecturer of Dermatology, Venereology and Andrology Faculty of Medicine – Ain Shams University

Faculty of Medicine
Ain Shams University
2014



Acknowledgements

First, and foremost, my deepest gratitude and thanks should be offered to "ALLAHI", the Most Kind and Most Merciful, for giving me the strength to complete this work.

I would like to express my sincere gratitude to **Prof. Dr. Hoda Ahmed M. Moneib**, Professor of Dermatology, Venereology and Andrology, Faculty of Medicine – Ain Shams University, for her continuous support and guidance for me to present this work. It really has been an honor to work under her generous supervision.

I acknowledge with much gratitude Dr. Al Hassan M. El Hefnawy and Dr. Azza Esmat, Lecturers of Dermatology, Venereology and Andrology, Faculty of Medicine – Ain Shams University, for his great supervision and unlimited help to provide all facilities to accomplish this work.

Last but not least, thanks to my Parents and my Husband, for helping me to finish this work.

Sara Mohammed Mahfouz Mohammed

List of Contents

| Subject | Page No. |
|--|----------|
| List of Abbreviations | i |
| List of Tables | iii |
| List of Figures | iv |
| Introduction | 1 |
| Aim of the Work | |
| Review of Literature | |
| Chapter (1): Dermoscopy | 4 |
| Chapter (2): Classification of Erythematous Scaly Lesions Diagnosed by Dermoscopy | 41 |
| Patients, Materials and Methods | 84 |
| Results | |
| Discussion | 133 |
| Conclusions and Recommendations | 142 |
| Summary | 144 |
| References | 146 |
| Arabic Summary | |

List of Abbreviations

AIDS : Acquired immunodeficiency disease Syndrome

AK : Actinic keratosis

BCC : Basal cell carcinoma

BD : Bowen's disease

BWS : Blue-white structures

CCLE: Chronic Cutaneous Lupus Erythematosus

CD : Chronic dermatitis

CI : Confidence interval

CMM : Cutaneous malignant melanoma

CTD : Connective tissue disease

DLE : Discoid lupus erythematosus

FUMHD: Febrile ulceronecrotic Mucha-Habermann disease

IEC : Intraepidermal carcinoma

KA : Keratoacanthoma

LED : Light emitting diodes

LK : Lichenoid keratosis

LM : Lentigo Maligna

LP : Lichen planus

LPLK : Lichen planus-like keratosis

MF : Mycosis fungoides

Non-PSL: Non-Pigmented skin lesions

PABAK : Prevalence-adjusted bias adjusted kappa

pBD : Pigmented Bowen's disease

PLEVA : Pityriasis lichenoides et varioliformis acuta

List of Abbreviations (Cont...)

PP : Plaque psoriasis

PR : Pityriasis Rosea

PSL : Pigmented skin lesions

sBCC : Superficial basal cell carcinoma

SCC : Squamous cell carcinoma

SD : Seborrheic dermatitis

SE : Standard error

SFTs : Short fine telangiectasias

SK : Seborrheic keratosis

SPSS : Statistical package for social science

UVR : Ultraviolet rays

VD : Videodermatoscope

WS : Wickham striae

List of Tables

| Table N | o. Title | Page | No. |
|-------------------|--|-------|-----|
| Table (1): | Dermoscopic algorhythm illustrating mel versus non- melanocytic lesions. | - | |
| Table (2): | Dermoscopy Colors and location of me | lanin | 19 |
| Table (3): | Vascular structures seen by dermoscop | y | 31 |
| Table (4): | Histopathologic correlation of the predermoscopic structures | _ | |
| Table (5): | Recommended classification of clini histological BCC subtypes | | |
| Table (6): | Agreement between Dermoscopic and pathologic diagnosis | | |
| Table (7): | Crosstabulation of dermoscopic and pathologic diagnoses of the same lesion | | |
| Table (8): | Agreement between dermoscopic and pathologic diagnoses | | |

List of Figures

| Figure No | o. Title | Page No. |
|---------------------|--|-----------|
| Figure (1): | Optics of dermoscope | 7 |
| Figure (2): | Dermoscopes with image capturing fa | acility10 |
| Figure (3): | Dermatoscope | 11 |
| Figure (4): | Dermaphot | 12 |
| Figure (5): | Stereomicroscope | 13 |
| Figure (6): | Videodermatoscope | 14 |
| Figure (7): | Colors in dermoscopy depending location of melanin in the skin | |
| Figure (8): | Reticular pattern | 20 |
| Figure (9): | Globular pattern | 20 |
| Figure (10): | Cobblestone pattern | 20 |
| Figure (11): | Homogeneous pattern | 20 |
| Figure (12): | Starburst pattern | 20 |
| Figure (13): | Starburst pattern | 20 |
| Figure (14): | Parallel pattern | 21 |
| Figure (15): | Multicomponent pattern | 21 |
| Figure (16): | Lacunar pattern | 21 |
| Figure (17): | Pigment network in melanocytic tum | ors26 |
| Figure (18): | Streaks | 28 |
| Figure (19): | Blue-whitish veil | 28 |
| Figure (20): | Vascular patterns in tumoral lesions. | 31 |
| Figure (21): | Milia like cysts | 32 |
| Figure (22): | Comedo-like openings | 33 |

List of Figures (Cont...)

| Figure No | o. Title | Page No. |
|---------------------|--|----------|
| Figure (23): | Leaf-like areas | 34 |
| Figure (24): | Scabies | 39 |
| Figure (25): | Actinic porokeratosis. | 39 |
| Figure (26): | Sebaceous hyperplasia | 40 |
| Figure (27): | Hemangioma and angiokeratoma | 40 |
| Figure (28): | Pyogenic granuloma | 40 |
| Figure (29): | Arborizing vessels and atypical red in seborrhoeic dermatitis | |
| Figure (30): | Schematic representations of dern patterns and underlying histological f | - |
| Figure (31): | Dermoscopic and histopathological of PLEVA | _ |
| Figure (32): | Model of dermoscopic time morphological changes of discoid erythematosus | d lupus |
| Figure (33): | Association between dermoscopic and pathology of basal cell carcinoma | |
| Figure (34): | Scheme of the dermoscopic structu in basal cell carcinoma | |
| Figure (35): | Seborrheic keratosis | 71 |
| Figure (36): | Lichenoid keratosis | 75 |
| Figure (37): | a) Dermoscope view lens b) Dermoscope contact lens | |
| Figure (38): | Attachment piece | 86 |
| Figure (39): | Definition of dermatoscopic features. | 89 |

List of Figures (Cont...)

| Figure No | . Title | Page No. |
|---------------------|--|----------|
| Figure (40): | Agreement between dermoscopy and histopathology | |
| Figure (41): | Non-pigmented AK | 97 |
| Figure (42): | Pigmented AK on the scalp | 99 |
| Figure (43): | Morpheic type of BCC | 101 |
| Figure (44): | Basal cell carcinoma | 103 |
| Figure (45): | Bowen's disease | 106 |
| Figure (46): | Non-pigmented Bowen's disease | 107 |
| Figure (47): | Pigmented Bowen's disease | 107 |
| Figure (48): | Keratoacanthoma | 109 |
| Figure (49): | Mycosis fungoides | 111 |
| Figure (50): | Plaque psoriasis | 113 |
| Figure (51): | Plaque psoriasis | 114 |
| Figure (52): | Scalp psoriasis | 115 |
| Figure (53): | Pityriasis rosea | 117 |
| Figure (54): | Lichen planus | 119 |
| Figure (55): | Acute/subacute eczema | 121 |
| Figure (56): | seborrheic dermatitis | 122 |
| Figure (57): | Chronic eczema | 124 |
| Figure (58): | Fungal infection. | 125 |
| Figure (59): | Vessels by dermoscopy for histopathological lesions | |
| Figure (60): | Vascular pattern by dermoscopy for histopathological lesions | |

List of Figures (Cont...)

| Figure No | o. Title | Page No. |
|---------------------|--|----------|
| Figure (61): | Background color by dermoscopy for histopathological lesions | |
| Figure (62): | Scale color by dermoscopy for histopathological lesions | |
| Figure (63): | Scale distribution by dermoscopy for histopathological lesions | |
| Figure (64): | Special features by dermoscopy for histopathological lesions | |

Introduction

Dermatoscopy (also known as dermoscopy, incident light microscopy, epiluminescence microscopy and skinsurface microscopy) is an inexpensive in vivo and non-invasive technique that permits the visualization of morphologic features of the epidermis and papillary dermis that are not visible to the naked eye. Previously, dermoscopy had been used in differentiating malignant pigmentary disorders, but the use of dermoscopy has recently been extended to the diagnosis of nonpigmentary skin disorders by defining the characteristic vasculature (*Zalaudek et al., 2006a*).

Intraepidermal carcinoma (IEC), superficial basal cell carcinoma (sBCC), and psoriasis may all present as a well-demarcated, variably scaly erythematous patch or plaque. Their differentiation poses a diagnostic challenge, particularly in the patient with both psoriasis and solar-damaged skin. When clinical diagnosis is not possible, biopsy is used to differentiate these entities. Dermatoscopy is valuable in the diagnosis and differentiation of IEC, sBCC, and psoriasis because of consistent dermatoscopic morphology (*Pan et al., 2008*).

Oligolesional psoriasis on the legs, where IEC is common, can be a difficult clinical diagnosis and dermatoscopy provides a quick, simple, and noninvasive aid. The major benefit from improved dermatoscopic differentiation of these common nonpigmented tumors (sBCC and IEC) is a reduction in the need for skin biopsy. In addition, if the tissue is not altered by surgical biopsy, monitoring the response to treatment with topical and light based skin cancer therapies may be easier. Just as these vascular patterns are useful in making a positive diagnosis, the loss of characteristic vascular pattern is also useful in providing dermatoscopic evidence of cure after therapy (*Micantonio et al.*, 2007).

Several previous studies, of limited size, have described some dermatoscopic features of IEC, sBCC, and psoriasis. The most significant dermatoscopic features of IEC were a clustered vascular pattern, glomerular vessels, and hyperkeratosis. When all 3 features were observed together, the diagnostic probability for IEC was 98% (Argenziano et al., 2004; Malvehy at al., 2006). sBCCs were characterized by a scattered vascular pattern, arborizing microvessels, telangiectatic or atypical vessels, milky-pink background, and brown dots/globules; the diagnostic probability was 99% if 4 of these 6 features were identified (Giacomel and Zalaudek, 2005; Argenziano et al., 2004). For psoriasis, the significant features identified were a homogenous vascular pattern, red dots, and light-red background, yielding a diagnostic probability of 99% if all 3 features were present (*Pan et al.*, 2008).

Aim of the Work

The aim of the current work is to evaluate the efficacy of dermoscopy in diagnosing scaly erythematous patches and plaques (eg. psoriasis, psoriasiform lesions) comparing dermoscopic diagnosis with histopathological results, as well as mentioning the dermoscopic features of each disease.

Chapter (1): **Dermoscopy**

Dermoscopy has gained popularity in recent years as an aid to the clinical diagnosis of many pigmented skin lesions (PSL) and non-pigmented skin lesions (non-PSL). The improvement in diagnostic accuracy of PSL has been well established (*Argenziano et al.*, 2003) but recent work has also suggested an improved diagnosis of lesions that are typically non-pigmented using this technique (*Zalaudek et al.*, 2005a). This is because dermoscopy allows the uncovering of key structures in non-PSL, including vascular patterns that are usually not visible to the naked eye (*Zalaudek et al.*, 2006a).

Historical background

The 1980s can be considered as the heyday of dermoscopy, with the definition of criteria for dermoscopy and the first Consensus Conference on Skin Surface Microscopy (*Bahmer et al.*, 1990).

However, the origin of the dermoscopic technique goes back as far as the 17th century, when Kohlhaus in 1663 first examined ungual matrix vessels under the microscope (*Gilje et al.*, 1953).

In 1878 Abbe improved the technique of skin surface microscopy by using immersion oil between the lens and the skin to improve the image clarity (*Carli et al.*, 2001b).