

شبكة المعلومات الجامعية







شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار في درجة حرارة من ١٥-٥٠ مئوية ورطوبة نسبية من ٢٠-٠٠% To be Kept away from Dust in Dry Cool place of 15-25- c and relative humidity 20-40%



بعض الوثائـــق الإصليــة تالفــة



بالرسالة صفحات لم ترد بالإصل

ais's

Looesly packed versus densely packed alloplast in the treatment of periodontal intrabony defects (A clinical and radiographical study)

Thesis

Submitted in partial fulfillment for

The requirement of

Master degree In

Oral Medicine, Oral Diagnosis and

Periodontology

By

Heba Samir Mahmoud

BDS 2004 (Cairo University)

Oral medicine, Oral Diagnosis and Periodontology Department Faculty of Oral and Dental Medicine

SUPERVISORS

Prof. Dr. Mona Salah El-Din Darhous

Professor and Head of Oral Medicine, Oral Diagnosis and
Periodontoloy
Faculty of Oral and Dental Medicine
Cairo University

Prof.Dr. Hala Zakaria

Professor of Oral Radiology

Faculty of Oral and Dental Medicine

Cairo University

Dr. Eman Yossef El-Firt

Lecturer of Oral Medicine, Oral Diagnosis and
Periodontology
Faculty of Oral and Dental Medicine
Cairo University

Acknowledgement

First and foremost, I would like to thank **God**, for allowing me to complete this subject and having made everything possible by giving me the strength and courage to do this work.

Also, words fail to express my feelings of gratitude and deep appreciation to **professor Dr. Mona Salah El-Din Darhous**, professor of Oral Medicine, Oral Diagnosis and Periodontology, Cairo University, for her great support, continuous teaching and guidance during this study.

I wish to extend my great thanks and appreciation to **Professor Dr. Hala Zakaria** Professor of Oral Radiology, Cairo University, for her wonderful participation and cooperation.

I would also like to express my warmest thanks to **Dr. Eman Yossef** El – Firt, Lecturer of Oral Medicine, Oral Diagnosis and Periodontology, Cairo University, for her patience, valuable advice and continuous support throughout the whole study.

Finally, I would to extend my thanks to all staff members of Oral Medicine, Oral Diagnosis and Periodontology, Cairo University, for their considerable cooperation, kind feelings and support.

Table of content

Page

Introduction Review of literature1
Aim of the study43
Subjects and methods
Results59
Discussion95
Conclusion
Summery109
References113
Arabic summery

List of Abbreviation

ADC	Analog digital conversion			
BMP	Bone morphogenetic protein			
BRM	Bone repair materials			
CAL	Clinical attachment level			
CCD	Charge- coupled device			
CEJ	Cemento-enamel junction			
CMOS	Complementary metal oxide semi			
	conductor			
CMOS-APS	Complementary metal oxide semi conductor active pixel sensor.			
СРС	Calcium phosphate cement			
DBM	Demineralized bone matrix			
EMD	Enamel matrix derivatives			

E-PTFE	Expanded Ploy tetrafluoroethylene
	membrane
GAG	Glycosaminoglycan
GI	Gingival index
GTR	Guided tissue regeneration
HA	Hydroxyapatite
HTR	Hard tissue replacement
MWF	Modified Widman Flap
NBM	Natural bone mineral
NSAIDs	Non steroidal anti- inflammatory
	drugs.
OFD	Open flap debridement
PD	Periodontal diseases
PI	Plaque index
PPD	Probing pocket depth
PRP	Platelet rich plasma

PSP	photo-stimulable phosphor
RVG	RadioVisioGraphy
SD	Standard deviation
в-тср	Beta-Tricalcium phosphate

List of figures

Figure number	Figure title	pages
Fig. (1)	Photograph showing Rinn XCP film holder consisting of interchangeable plastic bite blocks, a plastic aiming ring and a metallic indicator arm.	49
Fig. (2)	Photograph showing an acrylic occlusal stent.	49
Fig. (3)	Photograph showing the imaging plat.	49
Fig. (4)	Photograph showing measurement of density by Digora system for a defect mesial to the upper second premolar.	50
Fig.(5)	Photograph showing Linear measurement for a defect distal to the upper second premolar.	50
Fig. (6)	photograph showing KASIOS ®- tricalcium phosphate dental alloplast(150 - 500µm)	52
Fig. (7)	Immediate postoperative radiograph showing mesial and distal defects related to the upper second premolar with loosely packed tricalcium phosphate alloplast.	55
Fig.(8)	Immediate postoperative radiograph showing a distal defect related to the lower first molar with densely packed tri-calcium phosphate alloplast.	55

Fig. (9)	Clinical preoperative photograph showing 6 mm probing pocket depth distal to upper second premolar	60
Fig.(10)	Preoperative radiograph showing intrabony defects mesial & distal to the upper second premolar.	60
Fig. (11)	Reflected mucoperiosteal flap & intrabony defect mesial & distal to the upper second premolar.	60
Fig. (12)	Clinical photograph showing defect mesial to the upper second premolar filled with loosely packed alloplasts.	60
Fig. (13)	Six months postoperative phorograph showing 4 mm pocket depth distal to the upper second premolar.	61
Fig. (14)	Postopertive radiograph showing evidence of bone fill mesial & distal to the upper second premolar and remmenamt of the graft can still be detected.	61
Fig. (15)	Clinical preoperative photograph showing a probing pocket depth of 6 mm mesial to the lower first molar.	62
Fig. (16)	Preoperative radiograph showing the intrabony defect mesial to the lower first molar.	62
Fig. (17)	Reflected mucoperiosteal flap & the intrabony defect mesial to the lower first molar.	62

Fig. (18)	Clinical photograph showing defects mesial & distal to the lower first molar filled with loosely packed alloplast.	62
Fig.(19)	Six months postoperative photograph showing 2mm probing pocket depth mesial to the lower first molar.	63
Fig. (20)	Postoperative radiograph showing evidence of bone fill mesial to the lower first molar.	63
Fig.(21)	Preoperative photograph showing a probing pocket depth of 6mm mesial to the upper first premolar.	64
Fig.(22)	Preoperative radiograph showing the intrabony defect mesial& distal to the upper first premolar.	64
Fig.(23)	Reflected mucoperiosteal flap with the intrabony defects mesial & distal to the upper first premolar.	64
Fig.(24)	Clinical photograph showing the defects filled with densely packed alloplast.	64
Fig.(25)	Six months postoperative photograph showing 5 mm probing pocket depth mesial to the upper first premolar.	65
Fig.(26)	Postoperative radiograph showing evidence of bone fill mesial & distal to the upper second premolar.	65
Fig.(27)	Clinical preoperative photograph showing a probing pocket depth of 8 mm distal the lower first molar.	66
<u> </u>		