



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات

لم ترد بالأصل

B1. 519

Effect Of Early Feeding Practices And Mode Of Delivery On Neonatal Jaundice

Thesis

Submitted For Partial Fulfillment Of Master Degree (M. Sc.) In Pediatrics.

By

WALEED MOHAMED EL AZAB ABUL SOUD

M.B.B. Ch

Supervisors

Prof. AZZA M. A. M. ABUL FADL

*Professor of Pediatrics
Benha Faculty of Medicine
Benha University*

Prof. MOHAMED M. RASHAD

*Professor of Pediatrics
Benha Faculty of Medicine
Benha University*

Prof. MAHA SALAH ELDIN YOUSEF

*Professor of Clinical Pathology
Benha Faculty of Medicine
Benha University*

Dr. MOAZ EBN GABL AHMED

*Ass. Consultant of Pediatrics
Benha University Hospitals
Benha University*

**Faculty of Medicine
Benha University
2005**

Acknowledgement

First and foremost, I think Allah, the merciful and the beneficent.

I would like to express my sincere gratitude and thanks to Prof. Azza Abul Fadl, Professor of Pediatrics, Benha Faculty of Medicine, whose guidance, help and sincere supervision were the cornerstone in building up this thesis. She kindly supervised all of details of this work and revised all the work throughout its various steps. She offered me much of her help for unlimited experience in this research.

I would like to thank Prof. Mohamed Rashad, Professor of Pediatrics, Benha Faculty of Medicine, for his great help, encouragement, and constructive advise , He kindly offered much guidance , patience and supervised details of this work and revised all the work .

I also wish to thank Prof. Maha Salah Eldin Yousef, Professor of Clinical Pathology, Benha Faculty of Medicine, for her constant supervision, sincere advise and guidance.

I also wish to thank Dr Moaz Ebn Gabl Ahamed, Ass. Consultant of Pediatrics, Benha University Hospitals, for his continuous support and helpful guidance.

Finally I would like to express my deep thanks to my family and all my friends who really helped me all through this work by their sincere advise and encouragement.

Contents

	Page
Introduction and Aim Of The Work.	1
Review of Literature	4
• Chapter 1 : Feeding Practices	4
- Exclusive Breastfeeding.	5
- Recommendations on Breastfeeding	6
- Benefits of Breastfeeding	12
- Contraindications to Breastfeeding	16
- Lactation Management	19
- Breast Pumping.	28
- Breastfeeding Devices	31
- Baby Friendly Hospital Initiative(BFHI)	34
• Chapter 2 : Neonatal Jaundice	42
- Pathophysiology	43
- Causes of Jaundice	47
- Risk Factors for Development of Hyperbilirubinemia	52
- Bilirubin Encephalopathy & Kernicterus.	52
- Management of Neonatal Jaundice	54
• Chapter 3 : Intranatal Period	66
- Normal Birth	67
- Cesarean Section	92
- Hospital Stay for Healthy Term Infant	99
Methods and Subjects	104
Results	107
Discussion	123
Conclusions	131
Recommendations.	133
Summary	134
References	137
Annex	165
Arabic Summery	

List Of Tables

	Page
Table 1 : Exclusive Breastfeeding Rates For Infants In USA.	6
Table 2 : Egypt DHS studies	6
Table 3 : Baby's Age to Weight Gain	20
Table 4 : Baby's Age to Length Increase and Head Circumference Increase.	21
Table 5 : Causes Of Neonatal Jaundice	49
Table 6 : Comparison Of Jaundice	51
Table 7 : Laboratory Evaluation of Jaundiced Infant of 35 or More Weeks Gestation	58
Table 8 : Timing of Follow-up of Jaundiced Infants	60
Table I : Sex Distribution Among The Study Groups.	107
Table II : Sex and Peak Bilirubin Level	108
Table III : Feeding Practices.	109
Table IV : Breast Milk Expression	110
Table V : Breastfeeding Practices and Peak Bilirubin Level	111
Table VI : Mode of Delivery and Peak Bilirubin Level.	112
Table VII : Rooming In and Peak Bilirubin Level	113
Table VIII: Time of 1 st Breastfeed and Peak Bilirubin Level.	114
Table IX : Time of 1 st Skin & Eye Contact and Peak Bilirubin Level	115
Table X : Frequency of Feeding and Peak Bilirubin Level.	116
Table XI : Prelacteals and Peak Bilirubin Level	117
Table XI : Bottle Feeding and Peak Bilirubin Level	118
Table XII : Supplements and Peak Bilirubin Level.	119
Table XIII : Pacifiers and Peak Bilirubin Level.	120

List Of Figures

	Page
Figure 1 : Latch -On	22
Figure 2 : Madonna Or Cradle Hold.	23
Figure 3 : Breast Support.	23
Figure 4 : Wide open "ah".	23
Figure 5 : Cradle Hold.	26
Figure 6 : Side Lying Position.	26
Figure 7 : Clutch Hold.	27
Figure 8 : Cylinder Pump.	32
Figure 9 : 1 Pull or Push Per Second Pump.	32
Figure 10: Bulb Pump.	32
Figure 11: Large Electric Pump	32
Figure 12: Feeding Tube Devices Lact-Aid.	33
Figure 13: Nomogram for designation of risk in newborns.	58
Figure 14: Guidelines for phototherapy in hospitalized infants.	61
Figure 15: Guidelines for exchange transfusion in infants.	62
Figure 16: Positions that a woman may adopt during labour.	80
Figure 17: Effacement and dilatation of the cervix.	81
Figure 18: Positions that a woman may adopt during childbirth.	86
Figure I : Sex and Peak Bilirubin Level.	108
Figure II : Breastfeeding Expression	111
Figure III : Breastfeeding Practices and Peak Bilirubin Level	112
Figure IV : Mode of Delivery and Peak Bilirubin Level.	113
Figure V : Rooming In and Peak Bilirubin Level.	114
Figure VI : Time of 1 st Breastfeed and Peak Bilirubin Level.	115
Figure VII : Time of 1 st Skin & Eye Contact and Peak Bilirubin Level	116
Figure VIII : Frequency of Feeding and Peak Bilirubin Level.	117
Figure IX : Prelacteals and Peak Bilirubin Level	118
Figure X : Bottle Feeding and Peak Bilirubin Level	119
Figure XI : Supplements and Peak Bilirubin Level.	120
Figure XI : Pacifiers and Peak Bilirubin Level	121
Figure XIII: Follow up of Feeding practices and Peak Bilirubin Level	121
Figure XIII: Follow up of mode of delivery and Peak Bilirubin Level	122

List Of Abbreviations

AAP	: American Academy of Pediatrics.
AABB	: American Association of Blood Banks .
BFHI	: Baby Friendly Hospital Initiative.
CDC	: Center for Disease Control and Prevention growth charts.
CMV	: Cyto Meglo Virus.
CS	: Cesarean Section.
ETCO	: End Tidal Carbon Monoxide .
FIGO	: International Federation of Obstetricians and Gynaecologists
G6PD	: Glucose 6 Phosphate Deficiency.
HIV	: Human Immunodeficiency Virus.
ICM	: International Confederation of Midwives.
IDDM	: Insulin Dependant Diabetes Mellitus.
LLL	: La Leche League.
NCHC	: National Center for Health Statistics growth charts.
NICU	: Neonatal Intensive Care Unit.
RH	: Rhesus Factor.
S Ig A	: Serum Immuno globulin A.
SPSS	: Statically Package Social Science .
PROM	: Premature Rupture Of Membranes.
TBA's	: Traditional Birth Attendants
TCB	: Transcutaneous Bilirubinometry.
TENS	: Transcutaneous Electrical Nerve Stimulation
TSB	: Total Serum Bilirubin.
UDPGT	: Uridine Diphospho Glucuronyl Transferase
UNICEF	: United Nations Children's Fund
WHA	: World Health Assembly
WHO	: World Health Oragnization.

Introduction
&
Aim of the Work

Introduction and Aim of Work

Neonatal jaundice remains the most common and, perhaps, the most vexing problem in full-term infants during the immediate postnatal period. Despite the numerous articles published about this subject, many aspects of neonatal hyperbilirubinemia remain unexplained. Changes in the presentation and course of neonatal jaundice associated with breastfeeding have been recognized for some time, and have been the subject of scientific studies for the past twenty years (*Maisels, 2000*).

Though the data still tend to be conflicting, a reasonable summary of current knowledge suggests that breastfeeding is associated with an increased incidence of neonatal jaundice compared with formula feeding, jaundice in breast-fed infants tends to be more severe and that the duration of measurable hyperbilirubinaemia may extend for weeks or even months in some cases (*Zetterstrom, 1999*). A fairly typical infant could be the one who at 2 wk of age still has a serum bilirubin that hovers around 300–350 mmol/L, but who, beyond his obvious jaundice, appears perfectly healthy and has a satisfactory weight curve.

This phenomenon, as observed in our practice, requested an increased number of lab tests as well as repeated visits for infants who might need neither. To test or not to test, to release from further follow-up as healthy or to schedule a new appointment because you are not one hundred percent certain that the infant's serum bilirubin might not be >400 mmol/L in the next couple of days and whether you should be concerned because there is evidence suggesting that prolonged exposure to high serum bilirubin levels is associated with the risk of developmental sequelae or you should breath a sigh of relief because bilirubin as an antioxidant is probably good for the little baby are questions related to breastfeeding-associated jaundice that have become a common occurrence in the well baby nursery as well as in the follow-up of healthy infants born in our hospitals. (*Hammermann et al., 1998*).

In fact, our answers to such highly relevant clinical questions are still based more on suppositions than on solid clinical scientific proofs. There is accumulating evidence that kernicterus, the much feared sequel of neonatal jaundice, now occurs more frequently than used to be the case a decade ago (*Ebbesen, 2000 & Hansen, 2000*).

Some believe that this is caused by a combination of early release from maternity hospitals and lessened awareness of the potential danger of severe hyperbilirubinaemia (*Ebbesen, 2000*).

While these are probable contributory factors, there are also data suggesting that increased postnatal weight loss, a phenomenon associated with breastfeeding, is seen in a number of infants with extreme neonatal jaundice (*Hansen, 1997*).

So, is neonatal jaundice associated with breastfeeding totally innocuous, or there is a cause for concern? With this commentary, I hope I have made a case for the urgent need for increased scientifically based knowledge about breastfeeding-associated jaundice in the neonates. Is there more than one syndrome of jaundice associated with breastfeeding and breast milk? This question was recently reviewed by *Maisels (2000)* who concluded that the evidence to support the presence of two distinct syndromes is meager. He was referring to a suggested terminology whereby jaundice associated with breastfeeding in the first 2–4 days of life was called “the breastfeeding jaundice syndrome”, while that which appears at 4–7 days of life was called “the breast-milk jaundice syndrome”.

Inadequate breastfeeding, particularly in the first days of life, can as be associated with elevated levels of bilirubin (*Gartner, 2001*). This manifestation is known as breast-nonfeeding jaundice and is the neonatal manifestation of the adult disorder, starvation jaundice (*Whitmer & Gollan, 1983*). In this condition, marked by inadequate milk and caloric intake, but not necessarily by dehydration, there may be a delay in

bilirubin clearance resulting from low stool output (*De Carvalho, Robertson, & Klaus, 1985*) with an increase in the intestinal absorption of bilirubin.

Exaggerated jaundice due to poor breastfeeding should not be considered physiologic. It has been shown that there is neither a difference in the percentage of weight loss nor a difference in the percentage of neonates with elevated levels of bilirubin during the first days of life between those who were adequately breastfed on demand and those who were bottle-fed (*Rubaltelli, 1993*).

Despite the knowledge of an association between exaggeration of neonatal jaundice and breastfeeding, it will be a mistake to assume, without careful consideration, that because a neonate is breastfed and is jaundiced, breastfeeding is the sole or main cause of jaundice. It will be also a mistake to believe that if the jaundice is associated with breastfeeding, it can never be harmful (*Riordan, 2004*).

The mode of delivery also influences neonatal jaundice. There is a probable association between delivery by cesarean section and the method of feeding. Women subjected to cesarean section breastfeed infrequently during the first 48 hours after delivery (*Giovanna, 2001*).

After all, breast milk is the undisputed natural food for the newborn. The fact that this form of alimentation is frequently associated with limited food availability (and thus intake) in the first days of life is something for which the full term newborn is well prepared. It is interesting that the transition from the use of breast-milk substitutes at various periods during the 20th century to the current emphasis on breastfeeding is forcing us to reconsider some of our knowledge regarding the normal physiology of neonatal jaundice (*Gartner, 2001*).

Hintz et al., (2001) stated that for centuries, neonatal jaundice has caused medical practitioners to reflect and speculate as well as to offer