Contents

Subjects	Page
List of abbreviations	II
List of Figures	
List of Tables	
• Introduction	1
• Objectives	7
• Review of Literature	
♦ History of Rhinoplasty	8
♦ Anatomy	15
♦ Anthropometry and evaluation	42
♦ Non surgical rhinoplasty recent advance	ce56
♦ Contraindications, Adverse Effect and Complications	
 Review of Various Techniques in performing Non surgical rhinoplas 	sty94
Patients and Methods	116
• Results	126
• Illustrative cases	130
• Discussion	136
• Summary	142
• Conclusion	146
• Recommendations	148
• References	150
• Arabic Summary	

List of Abbreviations

	BDDE	.Butane I	Diol D	D iglycidy	yl Ether
--	-------------	-----------	---------------	-------------------	----------

BTxA.....Botulinum Toxin A

CaHA.....Calcium hydroxyl Apatite

HA.....Hyaluronic Acid.

NASHA......Non-Animal Stabilized Hyaluronic
Acid

PLLA.....Poly-L-Lactic Acid

PMMA.....PolyMethylMethAcrylate

SLR.....Single Lens Reflex

SMAS.....Superficial Muscular Aponeorotic

System

S-STE.....Skin-Soft Tissue Envelop

List of Figures

Fig no.	title		pag	ie no.
Figure	(1): Dieffenbach's met alter the size of the no			
Figure	(2): Facial relative skir	thickness	map	17
Figure	(3): Anatomical dissect muscles: nasal-labial a resection	rea after sk	in	21
Figure	(4): Anatomical dissect muscles: radix after sk			
Figure	(5): Anatomical muscles: nasal resection	pyramid	after	skin
Figure	(6): Anatomical dissec muscles: nasal pyrami resection	d after skin		25
Figure	(7): Anatomical dissect septi and myrtiformis			
Figure	(8): Nasal SMAS anato	omical disse	ection	28
Figure	(9): Nasal SMAS ligan	nents	•••••	30

Figure ((10): Nasal SMAS deep layer30
Figure ((11): External arterial vascularization of the nose
Figure ((12): External arterial vascularization of the nose and nasal danger zone
Figure ((13): Differences between male and female ideals for the nasofrontal (NFr) and nasolabial (NL) angles
Figure ((14): The head of the queen Nefertiti 1365 BC (Berlin State museum, Berlin, Germany)48
Figure ((15): The same patient is photographed twice before the procedure52
Figure ((16): While HA injection results can be seen immediately, in case of botulinum injections, at least 5 days are needed
Figure ((17): Over time, the carrier gel is gradually absorbed and the calcium hydroxylapatite particles remain
Figure ((18): Local histiocytic and fibroblastic response at the site appears to result in the production of new collagen around the microspheres

Figure	(19): Vascular anatomy of the midface84
Figure	(20): Three days after the injection. Crusted papules are visible in the distribution of the angular artery92
Figure	(21): Four months after the occlusive incident.
Figure	(22): Drawings depicting injection techniques: serial puncture96
Figure	(23): Harvesting and injection cannula98
Figure	(24): Entry points and injection paths to the zones being corrected
Figure	(25): Cannulas used by the author since 1998 to perform atraumatic harvest of 2- to 3-mm fat fragments
Figure	(26): Macroscopic view of cryopreserved fat graft 2 years after harvesting105
Figure	(27): Preoperative view of a patient after a previous rhinoplasty and Postoperative view at 6 months showing nasal lipoimplantation and touch-up procedure to improve dorsal and tip contours.

Figure	(28): Transcutaneous approach: two injection points, one at each side of the medial crura110
Figure	(29): Transcutaneous approach: one injection point at the base of the columella
Figure	(30): Injection of BNT-A at the base of the collumella
Figure	(31): The m. depressor septi nasi fibers run upwards to the tip of the nose Injection of BNT-A at the middle portion of the collumella is ideal for those patients who present a long upper lip
Figure	(32): The blocking of the m. depressor septi nast promotes a slight lifting and upward rotation of the tip of the nose
Figure	(33): Treatment algorithm for development of patient care treatment plan115
Figure	(34): Restylane package 0.5 and 1cc and Emla 5 % cream
Figure	(35): Emla cream 5%, alcohol pad, needles and insulin e syringe used in fat injection124

Figure (36): Pseudohump injected with Restylane above
and below the hump pre and post pictures (our
case)131
Figure (37A,B): Mild C shape deviation of the dorsum,
right side depression over the cephalic end of
LLC, depression over the upper end of the Rt.
nasolabial fold and assymetrical nostrils (our
case)133
Figure (38A,B,C): Pollybeak deformity after many rhinoplasties,
low dorsum, depression over the left LLC and low dorsum, pre and
nost 134

List of Tables

Table no.	title	page no.
Table (1):	Nasal muscles qouted	18
Table (2):	Measurement items	44
Table (3):	The Seven Deadly Traits	53
	Clinical literature overview of use llers in nonsurgical rhinoplasty	
` ´	ASAPS member statistics year on omparisons for dermal filler produc	•
Table (6):	Patient Demographics	77
M	Clinical Improvement in Patients licrofat Injection, as Rated by Both and Patient	the Author
• •	Suggested treatment for particulat ascular compromise	
Table (9):	Sources of Hyaluronidase	87



Introduction





Objectives





Review of Literature





Patients and Methods





Results





Illustrative Cases





Discussion

