





ثبكة المعلومات الجامعية





جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها علي هذه الأفلام قد اعدت دون آية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15-20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of 15-25c and relative humidity 20-40 %



ثبكة المعلومات الجامعية







A Bacteriological Approach To Asymptomatic Bacteriuria In Pregnancy

THESIS

Submitted to The High Institute of Public Health. University of Alexandria in partial fulfillment of the requirements for the degree of Master of Public Health

(Microbiology)

Ву

Mona Mohamed El Desouky El Sayed

MBBCH, Faculty of Medicine, University of Alexandria, 1995

High Institute of Public Health University of Alexandria 2003

SUPERVISORS



Prof. Dr. Laila Ahmed El-Attar

Professor of Microbiology High Institute of Public Health University of Alexandria

Laila

Dr. Wafaa Mohamed Kamel

Lecturer of Microbiology

High Institute of Public Health

University of Alexandria

DEDICATEDTO

My loving parents, sister and brother, which husband



my beloved kids

After thanking Allah, the most gracious the most merciful, I wish to express my everless deepest appreciation to my eminent Professor Dr. Laila Ahmed El Attar, Professor of Microbiology, High Institute of Public Health, for suggesting the subject, her supervision, meticulous guidance and valuable advices throughout this work. I gratefully acknowledge her continuous support and great efforts without which this thesis would not have been possible to start or finish, her encouragement offered openheartedly helped me proceed in my study during difficult times. It was true honour working under her supervision.

I would like to extend my deep thanks and profound appreciation to Dr. Wafaa Mohammed Kamel, Lecturer of Microbiology, High Institute of Public health for her meticulous help, alert guidance, continuous encouragement, and the precious time she

offered me for preparing each part of this work, her untiring efforts, patient guidance, constructive criticism and sisterly attitude helped to bring this work to light. I do sincerely and warmly thank her for all what she has done to overcome many obstacles throughout this work.

Deep thanks to all members of the staff and personnel of Microbiology department in the High Institute of Public Health, who directly or indirectly helped to end this work.

Last but not least I owe lots of thanks and gratitude to my family for their constant support, understanding and endless help without which I would have never fulfilled this work.





CONTENTS

Introduction	1
Aim of the Work	57
Material & Methods	58
Results	70
Discussion	102
Summary & Conclusion	120
Recommendations	126
References	127
Appendices	
Protocol	
Arabic Summary	

List of Tables

Table		Page
(I)	Bacterial enzymes detected by CHROMagar Orientation and interpretation of cultures on CHROMagar Orientation	54
(II)	Identification of staphylococci and micrococci	63
(III)	Identification of lactose fermenting isolates.	65
(IV)	Confirmatory tests for identification of Ps. aeruginosa	66
(V)	Identification of <i>Proteus</i> spp.	67
(VI)	Urine isolates presumptively identified on CHROMagar Orientation according to color reaction	68
(VII)	Culture outcome of the studied urine specimens of 400 asymptomatic pregnant women	72
(VIII)	Distribution of 118 pathogens isolated from 106 urine specimens with significant bacteriuria	75
(IXa)	The relation between culture outcome and Gram stain results of 400 urine specimen	77
(IXb)	The value of using Gram stain results in the detection of positive and negative cultures	77

	(Xa)	The relation between culture outcome of the 400 studied urine specimens and pus cell count/cmm	78
	(Xb)	The value of using pyuria(≥10 leukocytes/cmm) in differentiating significant from nonsignificant bacteriuria	78
	(XI)	The relation between Gram stain results and pus cell count in the 106 specimens having significant bacteriuria.	81
ř	(XII)	Relation between the risk factors of UTI and the culture outcome of 400 asymptomatic pregnant women.	83
	(XIII)	The relation between age and culture outcome of 400 asymptomatic pregnant women	85
	(XIV)	The relation between gravidity and culture outcome of 400 asymptomatic pregnant women	87
	(XV)	The relation between duration of pregnancy and culture outcome of 400 asymptomatic pregnant women	89
	(XVI)	Distribution of the 94 monomicrobial urine isolates in relation to the used culture media.	91
·	(XVII)	Distribution of 24 isolates recovered from 12 polymicrobial cultures out of the 400 urine specimens	93

detection of different urine pathogens isolated from the 400 urine specimens	
Agreement between each pair of the media used for urine culture	98
Presumptive identification of different isolates on CHROMagar Orientation	100

List of Figures

Figure	P	age
(1)	Female urinary tract	3
(2)	Scheme of renal tubules and its vascular supply	3
(3)	Weekly 24 hour GFR in pregnant women starting before conception and measured through the 12 th gestational weeks	7
(4)	Screening methods of urine examination	35
(5)	Microscopic field (oil immersion objective) showing the accumulation of <i>E. coli</i> at the edge of the urine drop (to the left) with the Gram-stained preparation of 10 μ l of a positive urine specimen ($\geq 10^5 CFU/ml$)	37
(6)	Microscopic picture of Gram stained urine samples	37
(7)	Streaking pattern for inoculating media for semi-quantitative bacterial colony counts	44
(8a)	Blood agar plate showing monomicrobial isolate (<i>K. pneumoniae</i>)	45
(8b)	MacConkey's agar plate showing polymicrobial growth (2 types of lactose fermenting colonies)	45
(9)	Diagram of dip-strip	46