

BIOCHEMICAL STUDIES ON GARLIC PLANT USING BIOTECHNOLOGY TECHNIQUES

BY

NERMEEN MOHAMMED FOUAD SAYED ARAFA

B.Sc. Agric. Sci. (Agricultural Biochemistry), Fac. Agric., Cairo Univ., ١٩٩٩

M.Sc. Agric. Sci. (Agricultural Biochemistry), Fac. Agric., Cairo Univ., ٢٠٠٦

THESIS

**Submitted in Partial Fulfillment of the
Requirements for the Degree of**

DOCTOR OF PHILOSOPHY

In

**Agricultural Science
(Agricultural Biochemistry)**

**Department of Agricultural Biochemistry
Faculty of Agriculture
Cairo University
EGYPT**

٢٠١٢

CONTENTS

	Page
INTRODUCTION	١
REVIEW OF LITERATURE	٤
١. Proximate analysis.....	٤
٢. Pigment content	٤
٣. Vitamin content.....	٥
٤. Secondary metabolites	٦
a. Flavonoid content	٦
b. Essential oil.....	٦
c. Volatile constituents and sulfur compounds content	٨
٥. Biological and medicinal applications of garlic plants	١٣
a. Antibacterial effect	١٣
b. Antifungal and antiyeast effects	١٧
c. Antiviral effect.....	١٩
d. Antioxidant effect	٢٠
e. Antitumor effect and other medicinal applications	٢٦
٦. Tissue culture and biotechnology techniques.....	٣٢
MATERIALS AND METHODS	٦١
RESULTS AND DISCUSSION.....	٩١
١. Plant material and explant preparation of <i>Allium sativum</i> ...	٩١
٢. Activation of immature meristem-tips of garlic cultivars bulb	٩٢
٣. Plantlets formation from the activated meristem-tips of <i>Allium sativum</i> L. cultivars.....	٩٥
a. Shoots formation	٩٥
b. Roots formation.....	٩٧
٤. Callus cultures formation.....	١٠٠
a. Callus initiation.....	١٠٠
b. Callus production.....	١٠٢
c. Growth dynamics of the callus tissues of garlic cultivars.....	١٠٥
(١) Callus production of Baladi cv. cultivar	١٠٧
(٢) Callus production of Seds ٤٠ cv. cultivar.....	١٠٨
(٣) Callus production of Igaside cv. cultivar.....	١٠٨
٥. Plant regeneration of garlic cultivars through somatic embryogenesis	١١١
a. Initiation of embryogenic callus.....	١١١

b. Maintenance of embryogenic callus.....	۱۱۴
c. Germination of somatic embryos	۱۱۵
d. In vitro cloves (microbulbs) production and conversion of Embryos	۱۲۰
۶. Bulblets formation and in vitro cloves preservation	۱۲۴
a. Bulblets formation of <i>in vitro</i> cloves.....	۱۲۴
b. Preservation of in vitro cloves by encapsulation method.....	۱۲۵
۷. Shoot induction and plantlets proliferation of garlic	
Cultivars	۱۲۸
a. Shoot induction from shoot-tips of garlic cultivars.....	۱۲۹
b. Plantlets proliferation of garlic cultivars.....	۱۳۰
c. <i>In vitro</i> bulbs production	۱۳۴
۸. Effect of salts (NaCl and CaCl_۲) on the growth of shoot tips in garlic Cultivars	۱۳۷
a. Effect of NaCl and CaCl _۲ on plantlets growth of <i>Baladi cv.</i> Cultivar.....	۱۳۷
(۱) Leaves formation	۱۳۷
(۲) Roots formation.....	۱۳۹
b. Effect of NaCl and CaCl _۲ on plantlets growth of <i>Seds ۴۰ cv.</i> cultivar	۱۴۳
(۱) Leaves formation.....	۱۴۳
(۲) Roots formation.....	۱۴۶
c. Effect of NaCl and CaCl _۲ on plantlets growth of <i>Igaside cv.</i> Cultivar.....	۱۴۹
(۱) Leaves formation.....	۱۴۹
(۲) Roots formation.....	۱۵۱
۹. Transformation of garlic by <i>Agrobacterium tumefaciens</i> mediated gene transfer	۱۵۸
۱۰. Biological study	۱۶۱
a. Free radical scavenging capacity (RSC)	۱۶۱
(۱) Determination of antioxidant activity of <i>in vitro</i> cultures of Baladi cv., Seds ۴۰ cv. and Igaside cv. cultivars and their <i>in vivo</i> bulbs	۱۶۱
(۲) Determination of antioxidant activity of <i>in vitro</i> cultures of Baladi cv., Seds ۴۰ cv. and Igaside cv. cultivars and their <i>in vivo</i> bulbs at different concentrations	۱۶۳
(۳) Determination of antioxidant activity of bulblets from <i>in vitro</i> cloves of Igaside cv. cultivar cultured on MS-	

medium containing different precursors	١٦٥
b. Anti-tumor potency of the garlic cultivars extracts	١٦٧
(١) Effect of <i>in vitro</i> cultures and <i>in vivo</i> bulbs of Baladi cv., Seds ٤٠ cv. and Igaside cv. cultivars on viability of Ehrlich Ascites Carcinoma Cells (EACC).....	١٦٧
(٢) Effect of bulblet extracts of Igaside cv. <i>in vitro</i> cloves cultured on MS-medium containing different precursors on the viability of Ehrlich Ascites Carcinoma Cells (EACC)	١٦٩
c. Antimicrobial activity of garlic extracts of Baladi cv., Seds ٤٠ cv. and Igaside cv. cultivars.....	١٧٢
(١) Effect of <i>in vitro</i> cultures and <i>in vivo</i> bulb extracts of Baladi cv., Seds ٤٠ cv. and Igaside cv. cultivars on the growth of bacterial, fungal and yeast strains	١٧٢
(٢) <i>In vitro</i> activity of bulblet extracts of Igaside cv. <i>in vitro</i> cloves cultured on MS-medium containing different precursors against some of pathogenic bacteria and fungi	١٨٢
١١. Electrophoretic analysis	١٨٨
a. Characterization of protein patterns of garlic using Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS- PAGE).....	١٨٨
(١) Characterization of protein patterns of salt-free plantlets and salt tolerant plantlets of Baladi cv., Seds ٤٠ cv. and Igaside cv. cultivars	١٨٨
(٢) Characterization of protein patterns of <i>in vitro</i> plantlets of Igaside cv. cultivar cultured on MS-medium containing different concentrations of cysteine-HCl as a precursor	١٩٢
(٣) Characterization of protein patterns of <i>in vitro</i> plantlets of Igaside cv. cultivar cultured on MS-medium containing different concentrations of serine as a precursor.....	١٩٥
(٤) Characterization of protein patterns of <i>in vitro</i> plantlets of Igaside cv. cultivar cultured on MS-medium containing different concentrations of glutathione as a precursor	١٩٨
(٥) Characterization of protein patterns of bulblets of Igaside cv. <i>in vitro</i> cloves cultured on medium fortified with various precursors at different concentrations.....	٢٠٢
b. Classification of genetic variation in garlic revealed by Random Amplified Polymorphic DNA Poly Chain	

Reaction. (RAPD-PCR)	٢٠٥
(١) Genetic characterization of DNA patterns of <i>in vitro</i> plantlets and <i>in vivo</i> bulbs of Baladi cv., Seds ٤٠ cv. and Igaside cv. by RAPD	٢٠٥
(٢) Genetic characterization of DNA patterns of bulblets from <i>In vitro</i> cloves of Igaside cv. cultivar cultured on medium fortified with various precursors at different concentrations by RAPD	٢١٠
١٢. Chemical analysis of garlic cultivars	٢١٦
a. Total proline content in plantlet garlic cultivars	٢١٦
b. Chemical composition of the volatile compounds of <i>Allium sativum</i> Igaside cv. cultures obtained by steam-distillation	٢١٨
(١) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for salt-free plantlets of <i>Allium sativum</i> Igaside cv. cultivar	٢١٩
(٢) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for salt plantlets of <i>Allium sativum</i> Igaside cv. cultivar.....	٢١٩
c. Chemical composition of the volatile compounds of <i>Allium sativum</i> Igaside cv. cultures grown on potential intermediates (precursors)	٢٢٦
(١) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for <i>in vivo</i> bulbs of Igaside cv. cultivar	٢٢٧
(٢) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for bulblets of Igaside cv. <i>in vitro</i> cloves cultured on precursor-free MS-medium	٢٢٧
(٣) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for bulblets of Igaside cv. <i>in vitro</i> cloves cultured on ١٢,٨٣ mM cysteine-HCl	٢٢٩
(٤) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for bulblets of Igaside cv. <i>in vitro</i> cloves cultured on ٩,٥١ mM serine.....	٢٢٩
(٥) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for bulblets of Igaside cv. <i>in vitro</i> cloves cultured on ١,٦٢ mM glutathione	٢٣٤
(٦) Gas Chromatography-Mass spectrometry (GC-MS) of the	

volatile components for bulblets of Igaside cv. <i>in vitro</i> bulbs cultured on ١٢,٨٣ mM cysteine-HCl	٢٣٥
(٧) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for bulblets of Igaside cv. <i>in vitro</i> bulbs cultured on ٩,٥١ mM serine	٢٣٦
(٨) Gas Chromatography-Mass spectrometry (GC-MS) of the volatile components for bulblets of Igaside cv. <i>in vitro</i> bulbs cultured on ١,٦٢ mM glutathione	٢٣٧
SUMMARY	٢٤٢
REFERENCES	٢٥٣
ARABIC SUMMARY	

CONTENTS

	Page
INTRODUCT ION.....	۱
REVIEW OF LITERATURE	۴
۱. Proximate analysis.....	۴
۲. Pigment content	۴
۳. Vitamin content.....	۵
۴. Secondary metabolites.....	۶
a. Flavonoid content	۶
b. Essential oil.....	۶
c. Volatile constituents and sulfur compounds content.....	۸
۵. Biological and medicinal applications of garlic plants.....	۱۳
a. Antibacterial effect	۱۳
b. Antifungal and antiyeast effects	۱۷
c. Antiviral effect.....	۱۹
d. Antioxidant effect	۲۰
e. Antitumor effect and other medicinal applications.....	۲۶
۶. Tissue culture and biotechnology techniques.....	۳۲
a. The dormancy of shoot tips of <i>Allium sativum</i>	۳۳
b. Establishment of callus cultures of <i>Allium sativum</i>	۳۴
c. Establishment of embryogenic cell cultures of <i>Allium</i>	۳۵
d. Multiplication of <i>Allium sativum</i>	۴۴
e. <i>In vitro</i> bulbils production of <i>Allium sativum</i>	۵۲
f. Effect of salt stress on growth of <i>Allium sativum</i>	۵۳
g. Electrophoretic analysis of <i>Allium sativum</i>	۵۶
MATERIALS AND METHODS.....	۶۱
RESULTS AND DISCUSSION.....	۸۸
۱. Plant material and explant preparation of <i>Allium sativum</i>	۸۸
۲. Activation of immature meristem-tips of garlic cultivars bulb	۸۹
۳. Plantlets formation from the activated meristem-tips of <i>Allium sativum</i> L. cultivars.....	۹۲
a. Shoots formation.....	۹۲
b. Roots formation	۹۴
۴. Callus cultures formation.....	۹۷
a. Callus initiation	۹۷
b. Callus production	۹۹
c. Growth dynamics of the callus tissues of garlic cultivars	۱۰۲

9. Plant regeneration of garlic cultivars through somatic Embryogenesis	108
a. Initiation of embryogenic callus	108
b. Maintenance of embryogenic callus	111
c. Germination of somatic embryos.....	112
d. In vitro cloves (microbulbs) production and conversion of Embryos.....	117
10. Bulblets formation and in vitro cloves preservation	121
a. Bulblets formation of <i>in vitro</i> cloves.....	121
b. Preservation of in vitro cloves by encapsulation method.....	122
c. Bulblets retrieval from encapsulated in vitro cloves.....	123
11. Shoot induction and plantlets proliferation of garlic cultivars	125
a. Shoot induction from shoot-tips of garlic cultivars	126
b. Plantlets proliferation of garlic cultivars.....	127
c. <i>In vitro</i> bulbs production	131
12. Effect of salts (NaCl and CaCl ₂) on the growth of shoot tips in garlic Cultivars	134
a. Effect of NaCl and CaCl ₂ on plantlets growth of <i>Baladi cv.</i> cultivar.....	134
b. Effect of NaCl and CaCl ₂ on plantlets growth of <i>Seds 4 cv.</i> cultivar.....	140
c. Effect of NaCl and CaCl ₂ on plantlets growth of <i>Igaside cv.</i> cultivar.....	146
13. Biological study	150
a. Free radical scavenging capacity (RSC)	150
b. Anti-tumor potency of the garlic cultivars extract.....	161
c. Antimicrobial activity of garlic extracts of <i>Baladi cv.</i> , <i>Seds 4 cv.</i> and <i>Igaside cv.</i> cultivars.....	167
14. Electrophoretic analysis	183
a. Characterization of protein patterns of garlic using Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS- PAGE).....	183
b. Classification of genetic variation in garlic revealed by Random Amplified Polymorphic DNA Poly Chain Reaction. (RAPD-PCR).....	200
15. Chemical analysis of garlic cultivars	211
a. Total proline content in plantlet garlic cultivars	211

b. Chemical composition of the volatile compounds of <i>Allium sativum</i> Igaside cv. cultures obtained by steam- distillation.....	۲۱۳
CONCLUSION	۲۳۷
SUMMARY	۲۴۰
REFERENCES	۲۵۰
ARABIC SUMMARY	

LIST OF FIGURES

No.	Title	Page
١.	Schematic presentation of the preparation of encapsulated garlic <i>in vitro</i> cloves using a calcium alginate gel.....	٧٠
٢.	The activated meristem-tips of <i>Seds</i> ξ • <i>cv.</i> and <i>Igaside cv.</i> cultivars grown on MS-medium free hormone or MS-medium supplemented with ٥ mg/l BAP.....	٩١
٣.	Shoots derived from the activated merstim-tips of <i>Igaside cv.</i> (A), <i>Seds</i> ξ • <i>cv.</i> (B) and <i>Baladi cv.</i> (C) which grown on MS- medium supplemented with ٥ mg/l BAP at the end of the sixth week of culture.....	٩٣
٤.	The induced roots of plantlets of <i>Baladi cv.</i> cultivar grown on MS-basal medium (A) and MS- medium supplemented with ١ mg/l of both of NAA (B), IAA (C) and IBA (D).....	٩٦
٥.	Roots derived calli of both of <i>Baladi cv.</i> , <i>Seds</i> ξ • <i>cv.</i> and <i>Igaside cv.</i> cultivars after three months of cultivation on MS-medium supplemented with picloram (٥ mg/l).....	١٠٠
٦.	Root calli of <i>Baladi cv.</i> (A), <i>Seds</i> ξ • <i>cv.</i> (B) and <i>Igaside cv.</i> (C) after three months of culture on MS – medium supplemented with ٥ mg/l picloram+٦ mg/l thiamine-HCl.....	١٠٣
٧.	Produced callus of leaf calli of <i>Igaside cv.</i> after three months of culture medium supplemented with ٥ mg/l picloram+٦ mg/l thiamine- HCl.....	١٠٣
٨.	Somatic embryogenesis of <i>Allium sativum L. Igaside cv.</i> embryogenic callus and early stage somatic embryos (A and B) and development (C) of root calli cultured on Solid MS-medium supplemented with ٧mg/l kinetin.....	١١٠

٩. Somatic embryos of roots calli in <i>Baladi cv.</i> , (A) <i>Seds ٤٠ cv.</i> (B) and <i>Igaside cv.</i> (C) cultured on MS-medium containing ٧ mg/l Kin after sixty days of culture.....	١١٣
١٠. Mature green somatic embryos of embryogenic root calli of <i>Igaside cv.</i> cultured on MS-medium containing ٧mg/l Kin within two weeks of culture.....	١١٤
١١. Shoots derived from somatic embryos of <i>Igaside cv.</i> cultivar cultured on MS-medium supplemented with ٧ mg/l Kin after three (A) and six (B) weeks of culture.....	١١٦
١٢. Shoots derived from somatic embryos of <i>Seds ٤٠ cv.</i> cultivar culture on MS-medium supplemented with ٧ mg/l Kin after six weeks of culture.....	١١٦
١٣. <i>In vitro</i> cloves of shoots derived from somatic embryos of <i>Igaside cv.</i> cultivar cultured on MS-medium supplemented with ٥ mg/l IBA after one month of culture.....	١١٩
١٤. <i>In vitro</i> plantlets of shoots derived from somatic embryos of <i>Igaside cv.</i> cultivar cultured on MS-medium supplemented with ٣ mg/l NAA after one month of culture.....	١٢٠
١٥. Bulblets formation of <i>in vitro</i> cloves of <i>Igaside cv.</i> cultivar grown on MS-medium fortified with ٥ mg/l BAP (A) and growth regulators free MS-medium (B) after one month of culture.....	١٢٢
١٦. State of artificial garlic seeds prepared by encapsulating <i>Igaside cv.</i> cultivar <i>in vitro</i> cloves using calcium alginate gel.....	١٢٤
١٧. The recovery of <i>in vitro</i> preserved cloves of <i>Igaside cv.</i> on MS-medium containing ٥ mg/l BAP.....	١٢٤

١٨. Plantlet derived from shoot tip of <i>Igaside</i> cv. cultivar grown on growth regulator-free MS- medium.....	١٢٦
١٩. Multiple plantlets derived from shoot-tip of <i>Seds</i> ξ cv.(A), <i>Baladi</i> cv.(B) and <i>Igaside</i> cv.(C) grown on MS- medium containing ٦ mg/l BAP and <i>Igaside</i> cv.(D) grown on growth regulators-free medium after two months of culture.....	١٢٩
٢٠. Multiple plantlets derived from shoot-tip of <i>Seds</i> ξ cv. (A) and <i>Igaside</i> cv. (B) cultured on MS- medium containing ٦ mg/l BAP after two months of culture.....	١٣٠
٢١. Development stages of <i>in vitro</i> multiplication of cutting shoot-tip of <i>Igaside</i> cv. cultivar within four weeks of cultivation.....	١٣٢
٢٢. <i>In vitro</i> bulbs of shoot tip of <i>Igaside</i> cv. cultivar after one month of the third subculture on growth regulators-free MS-medium.....	١٣٣
٢٣. Leaves of shoot tips of <i>Baladi</i> cv. cultivar cultured on salts-medium (A), medium containing ٠,٤% of NaCl + ٠,٤% CaCl _٢ (B), medium containing ٠,٦% of NaCl + ٠,٦% CaCl _٢ and medium containing ٠,٨% of NaCl + ٠,٨% CaCl _٢ (D) after three weeks of culture.....	١٣٧
٢٤. Roots of plantlets of <i>Baladi</i> cv. cultivar cultured on MS – medium containing ٠,٦% NaCl.....	١٣٩
٢٥. Leaves of shoot tips of <i>Seds</i> ξ cv. cultivar cultured on salts-free medium (A), medium containing ٠,٤% of NaCl + ٠,٤% CaCl _٢ (B), medium containing ٠,٦% of NaCl + ٠,٦% CaCl _٢ (C) and medium containing ٠,٨% of NaCl + ٠,٨% CaCl _٢ (D) after three weeks of culture.....	١٤١

၃၆.	Roots of plantlets of <i>Seds</i> ၁၂ cv. cultivar cultured on MS – medium containing ၀.၁% CaCl ₂	၁၄၀
၃၇.	Leaves of shoot tips of <i>Igaside</i> cv. cultivar cultured on salts-free medium (A), medium containing ၀.၁% of NaCl + ၀.၁% CaCl ₂ (B), medium containing ၀.၂% of NaCl + ၀.၂% CaCl ₂ (C) and medium containing ၀.၄% of NaCl + ၀.၄% CaCl ₂ (D) after three weeks of culture.....	၁၄၁
၃၈.	Roots of plantlets of <i>Igaside</i> cv. cultivar cultured on MS – medium containing ၀.၂% of NaCl plus ၀.၂% CaCl ₂	၁၀၁
၃၉.	Viability of Ehrlich Ascites Carcinoma cells untreated with garlic extracts.....	၁၆၃
၄၀.	Viability of Ehrlich Ascites Carcinoma Cells treated with garlic extracts.....	၁၆၃
၄၁.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Bacillus subtilis</i>	၁၇၀
၄၂.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Staphylococcus aureus</i>	၁၇၁
၄၃.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Pseudomonas floureceans</i>	၁၇၂
၄၄.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Escheria coli</i>	၁၇၃
၄၅.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Aspergillus niger</i>	၁၇၄
၄၆.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Fusarium</i>	၁၇၅

۳۷.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Pichia</i>	۱۷۵
۳۸.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Sccharomyces cervisiae</i>	۱۷۶
۳۹.	Zones of inhibition caused by different garlic extracts on growth medium seeded with <i>Candida albicans</i>	۱۷۷
۴۰.	Zones of inhibition caused by bulblet extracts of <i>Igaside</i> cv. <i>in vitro</i> cloves cultured on MS-medium containing different precursors on growth medium seeded with <i>Staphylococcus aureus</i> (A) and <i>Escheria coli</i> (B).....	۱۸۰
۴۱.	Zones of inhibition caused by bulblet extracts of <i>Igaside</i> cv. <i>in vitro</i> cloves cultured on MS-medium containing different precursors on growth medium seeded with <i>Pseudomonas floureceans</i> (C) and <i>Aspergillus fusarium</i> (D).....	۱۸۱
۴۲.	SDS-PAGE of garlic plantlets cultured on salt-free medium and medium fortified with ۰,۶٪ of NaCl + CaCl.....	۱۸۶
۴۳.	SDS-PAGE of garlic plantlets of <i>Igaside</i> cv. cultivar cultured on MS- medium containing different concentrations of cysteine-HCl.....	۱۹۰
۴۴.	SDS-PAGE of garlic plantlets of <i>Igaside</i> cv. cultivar cultured on MS- medium containing different concentrations of serine.....	۱۹۳
۴۵.	SDS-PAGE of garlic plantlets of <i>Igaside</i> cv. cultivar cultured on MS-medium containing different concentrations of glutathione.....	۱۹۶
۴۶.	SDS-PAGE of bulblets of <i>Igaside</i> cv. cultivar cultured on MS-medium containing various precursors at different	

concentrations.....	199
47. Banding pattern of garlic cultures generated with X, Y and Z Random Amplified Polymorphic DNA (RAPD) primers and ladder of 3 kbp.....	200
48. Banding pattern of bulblets of <i>Igaside cv. in vitro</i> cloves generated with X, Y and Z Random Amplified Polymorphic DNA (RAPD) primers and ladder of 3 kbp.....	210