

# 







شبكة المعلومـــات الجامعية التوثيق الالكتروني والميكروفيا.



### جامعة عين شمس

التوثيق الالكتروني والميكروفيلم



نقسم بللله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد اعدت دون آية تغيرات



#### يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

40-20 في درجة حرارة من 15-20 منوية ورطوبة نسبية من

To be kept away from dust in dry cool place of 15 – 25c and relative humidity 20-40 %









B1.97.

# EFFECT OF PULSED ELECTROMAGNETIC FIELDS ON HEALING OF INFECTED BURN WOUND IN GUINEA PIGS

#### Ву

#### ALY OSMAN ALY SELIM

(B.Sc., in "Physical Therapy,1997")

#### A Thesis

Submitted in Partial Fulfillment for the Requirement of Master degree in Physical Therapy

Faculty of Physical Therapy
Cairo University
2003

#### **SUPERVISORS**

#### Prof. Dr. Adel Abdel El-Hamid Nossier

Chairman of The Physical Therapy Department for Surgery
Faculty of Physical Therapy
Cairo University

#### Prof. Dr. Fadel Mohamed Ali

Professor of Biophysics Faculty of Science Cairo University

#### Dr. Haytham Mohamed El-Hafez

Lecturer in Basic Science Department Faculty of Physical Therapy Cairo University

# ◄ بسم الله الرحمن الرحيم

# " قالوا سبحانك لاعلم لنا إلا ماعلمتنا إنك أنت العليم المكيم"

.صدق الله العظيم.

سورة البقرة آيـة (٣٢) TO THE SOULS OF MY MOTHER

AND FATHER,

TO MY WIFE AND MY DAUGHTER

AND TO MY FAMILY

#### **ACKNOWLEDGMENT**

First and foremost, thanks to *God*, the most gracious, the most Merciful.

I wish to express my sincere gratitude and deep appreciation to **Professor Dr. Adel Abd El- Hamid Nossier**, Professor and Chairman of Physical Therapy Department for Surgery, Faculty of Physical Therapy, Cairo University. He gave me a great deal of his valuable time and effort for this work. His comments and guidance were very helpful to me.

I am truly grateful to *Professor Dr. Fadel Mohamed Ali*, Professor of Biophysics Faculty of Science, Cairo University, for his valuable help; support, advice, guidance and effort throughout this work.

It is a great honor to me to express my most sincere and thanks to **Dr. Haytham Mohamed El-Hafez**, Lecturer in Basic Science Department Faculty of Physical Therapy, Cairo University for his valuable supervision and Kind advices throughout the whole work.

I wish to express my gratitude and special thanks also to my Professors and Colleagues in Physical Therapy Department for Surgery, Faculty of Physical Therapy, Cairo University, For their unlimited help, and support.

Alí Osman Selím 2003

#### List of Abbreviations

A : Ampere

cm :: Centimeter

cm<sup>2</sup> : Squared Centimeter

D.C. : Direct current

**ELF** : Extremely Low Frequency

**EMF** : Electromagnetic Field

Fig : Figure

G : Gauss

GP : Guinea Pigs

Hz : Hretz

HIPEF : High Intensity Pulsed Electric Field

kv : Kilo volt

mA : milliampere

MF : Magnetic Field

MG : milli gauss

MHz : Mega hertz

mm : milli meter

mV : milli volt

PS. : Pseudomonas bacteria

**QAMW** : Square Amplitude Modulated Wave

T : Tesla

TBSA : Total Body Surface Area

V : Volt

WSA : Wound Surface Area

## List of Tables

No.	Title	P
Table (4 - 1):	the mean, standard deviation, maximum, and	89
	minimum value of WSA after 24 hours post	,
:	induced burn, and after five, ten, and fifteen days of	
	post burn (Control Group A).	
Table (4– 2):	One way analysis of variance for repeated measure of	93
·	the WSA in control Group A.	•
Table (4– 3):	The mean, standard deviation, maximum, and	98
	minimum value of WSA after 24 hours post	
	induced burn and after five, ten, and fifteen days	
,	post burn (Group B).	·
Table (4 –4):	One way analysis of variance for repeated measure of	100
	the WSA in control Group D.	
Table (4 –5):	The mean, standard deviation, maximum, and	105
	minimum value of WSA after one, five, ten, and	
	fifteen days post burn and infection induction	
	(Control Group C).	
Table (4– 6):	One way analysis of variance for repeated measure of	107
	the WSA in control Group C.	
Table (4– 7):	The mean, standard deviation, maximum, and	112
	minimum value of WSA before initiation of treatment	
	application with EMF(24 hours post induced burn),	
	and after five, ten, and fifteen days of exposure to EMF	
	( Group D).	,
		·

Table (4 –8):	One way analysis of variance for repeated measure of	114
	the WSA in experimental Group D.	
Țable (4 –9):	The mean, standard deviation, maximum, and minimum value of WSA before initiation of treatment application with EMF (24 hours post induced burn),	119
	and after five, ten, and fifteen days of exposure to EMF (Experimental Group E).	
Table (4-10):	One way analysis of variance for repeated measure of the WSA in experimental Group E.	121

### **List of Figures**

No	Figure	<b>P.</b>
Fig. (2-1):	Schematic Drawing of The Skin	14
;	(Delman and Brown 1976).	
Fig. (2-2):	Papillary and Reticular Zones of Dermis Layer.	15
	(Banls 1993)	
Fig. (2-3):	Typical Hair Follicle. (Banks 1993)	18
Fig. (2- 4 - A):	Saccular Apocrine Sweat Glands. (Delman and Brown	19
	1976)	
Fig. (2-4 - B):	Tubular Apocrine, Sweat Glands (Delman and Brown,	20
	1976)	
Fig. (2-5):	Superficial Partial-Thickness Burn, (Malick and Carr,	29
	1982).	
Fig. (2-6):	Deep Partial Thickness Burn, (Malick and Carr, 1982).	30
Fig. (2-7):	Full Thickness Burn, (Malick and Carr, 1982)	31
Fig. (2-8):	Rule of Nine for Determining Percentage of Body	32
	Surface Area Burned in Adults, (O'S ullivan and Schmitz	
	1994).	
Fig. (3-1):	An Image for Exposure of One of the Experimental	75
	Animals to QAMW.	
Fig. (3-2):	An Image for the Exposure Facility to QAMW.	76
Fig. (3-3):	Wound Surface Area Measurement Tools	78
Fig. (3-4):	Handling of Guinea Pigs.	81
Fig. (3-5):	Round Aluminum Stamp Used to Induce Burn Wound.	82 .
Fig. (3-6):	Wound Surface Area Measurement Procedure	84 .
Fig.( 4 – 1):	photograph for the wound of one of the animals 24 hours	88
	post induction of the burn injury	
Fig. (4 - 2):	the mean value of WSA, after 24 hours post induced	90
	burn and after five, ten, and fifteen days of post burn	
	(Control group A)	

Fig. $(4-3)$ :	shows a photograph for the wound of one of the animal,	92
	from group A, 10 days post infection.	
Fig. (4-4):	survival rate of the animals from grou A.	95
Fig.( 4 – 5 ):	a photograph for the wound of one of the animals of	97
	group B after 10 days of wound induction.	
Fig. (4 - 6 ):	the mean value of WSA after 24 hours post induced burn	99
·	and after five, ten, and fifteen days of post burn, for	
	(Control groupB).	
Fig. $(4-7)$ :	survival rate of the animals from group B.	102
Fig. (4 – 8):	a photograph for an infected wound for one animal	104
,	from group C.	
Fig. (4 - 9):	the mean values of WSA after one, five, ten, and fifteen	106
	days post burn and infection induction (Control GroupC).	-
Fig. (4 – 10):	Survival rate of the animals of group C.	109
Fig. (4-11):	a photograph for an infected wound in one animal of	111
·	group D at day 15.	-
Fig. (4 - 12):	the mean value of WSA, before initiation of treatment	113
	application of EMF( 24 hours post induced burn) and	
	after five, ten, and fifteen days of exposure to EMF	,
	(experimental group D).	
Fig.( 4-13 ):	the survival rate of the animals from group D.	116
Fig. ( 4-14 ):	a photograph for one animal from group E at day 15	118
Fig. (4 – 15):	the mean value of WSA, before initiation of treatment	120
	application of EMF( 24 hours post induced burn) and	
	after five, ten, and fifteen days of exposure to EMF	
	(experimental group E).	
Fig. (4-16):	shows the variation of the survival rate of the animals	123
-	from group E	
<del></del> _		