

**The Use of Simvastatin Gel with Perforated
Resorbable Membranes in the Treatment of
Intrabony Defects in Chronic Periodontitis
Patients
(Clinical and Biochemical Study)**

Thesis

**Submitted to department of Oral Medicine,
Periodontology, Oral Diagnosis and Oral Radiology
Faculty of Dentistry
Ain Shams University**

**In partial Fulfillment of the Requirements of Doctor Degree
in Oral Medicine, Periodontology, Oral Diagnosis and Oral
Radiology**

By

Dalia Mohammed Rasheed Mohammed Fathy Issa

B.D.S (Mansoura University)

M.Sc (Cairo University)

**Faculty of Dentistry
Ain Shams University**

2018

Suprvisors

Prof. Dr. Ahmed Y. Gamal

Professor of Oral Medicine and Periodontology
Faculty of Dentistry
Ain Shams University

Prof. Dr. Mohamed A. Al-Shahat

Professor of Oral Medicine and Periodontology
Faculty of Oral and Dental Medicine
Delta University for Science and Technology
Former, Professor, Faculty of Oral and Dental Medicine
Mansoura University

Dr. Ahmed Abd El Aziz Hassan

Ass. Professor of Oral Medicine and Periodontology
Faculty of Dentistry
Ain Shams University

Acknowledgement

First, and for most, I feel always indebted to God, the most kind and merciful.

*I would like to express my deep gratitude and appreciation to Professor **Dr. Ahmed Y. Gamal** Professor of Oral Medicine and Periodontology, Ain Shams University Whose supervision on my thesis is a great honor to me. Without his support and his guidance this work would not have been possible. He was always accessible, willing to help me, giving generously his valuable time so, research work became smooth and rewarding for me.*

*Of course, it's necessary to express my sincere appreciation to Professor **Dr. Mohamed A. AL-Shahat** Professor of Oral Medicine and Periodontology, Faculty of Oral and Dental medicine, Mansoura University for his great assistance, academic supervision and highly cooperation in this study.*

*I am also grateful to **Ass. Professor Dr. Ahmed Abd El Aziz** Ass. Professor of Oral Medicine and Periodontology, Ain Shams University, who encouraged me and helped me throughout the work.*

I couldn't forget to express my appreciation to all the staff members and candidates of the Oral Medicine and Periodontology Department, Faculty of Dentistry, Ain Shams University for their highly cooperation.

Dedication

I'm very grateful and thankful for my father, mother, sister and brothers. Thank you for your kind support, effort, help and patience.

List of Contents

Title	Page No.
Introduction and Review of Literature.....	1
Aim of the Study.....	72
Materials and Methods.....	73
Results.....	114
Discussion.....	171
Summary.....	182
Conclusion.....	185
Recommendation.....	186
References.....	187
Appendix.....	232
Arabic Summary.....	-

LIST OF FIGURES

Figure(1)	Molecular Structure of SMV
Figure(2)	Mevalonate pathway
Figure(3)	Effect of SMV on bone
Figure(4)	The XCP film holder for posterior teeth with bite block
Figure(5)	Photostimulable phosphor plate (PSP) size (2), protective cover, and the hygiene bag.
Figure(6)	The intraoral X-ray machine was adjusted to fit into the ring of the film holder
Figure(7)	Linear measurement of bone height using Digora software
Figure(8)	Radiodensitometric measurement using Digora software
Figure(9)	SMV gel
Figure(10)	Collagen membrane (Resorba Resodont Fort)
Figure(11)	EDTA
Figure(12)	Collection of GCF with filter paper
Figure(13)	Whatman filter paper
Figure(14)	HPLC
Figure(15)	A photograph showing a 6mm pocket depth mesial

	to the lower left first molar 9 months presurgically
Figure(16)	Preoperative radiograph showing intrabony defect related to mesial surface of the lower left first molar 9 months presurgically
Figure(17)	Intrabony defect after curetteage of granulation tissues
Figure(18)	Application of SMV gel
Figure(19)	The occlusive membrane was positioned over the defect
Figure(20)	Suture using polypropylene
Figure(21)	A photograph showing a 6mm pocket depth distal to the upper left first molar 9 months presurgically
Figure(22)	Preoperative radiograph showing intrabony defect related to distal surface of the upper left first molar 9 months presurgically
Figure(23)	Intrabony defect after curetteage of granulation tissues
Figure(24)	Application of SMV gel
Figure(25)	Perforated membrane
Figure(26)	The perforated membrane was positioned over the defect
Figure(27)	Suture using polypropylene
Figure(28)	A photograph showing an 8mm pocket depth mesial

	to the upper left first molar 9 months presurgically
Figure(29)	Preoperative radiograph showing intrabony defect related to masial surface of the upper left first molar 9 months presurgically
Figure(30)	Intrabony defect after curetteage of granulation tissues
Figure(31)	Application of EDTA
Figure(32)	Application of SMV gel
Figure(33)	The occlusive membrane was positioned over the defect
Figure(34)	Suture using polypropylene
Figure(35)	A photograph showing a 6mm pocket depth mesial to the lower left first molar 9 months presurgically
Figure(36)	Preoperative radiograph showing intrabony defect related to masial surface of the lower left first molar 9 months presurgically
Figure(37)	Intrabony defect after curetteage of granulation tissues
Figure(38)	Application of EDTA
Figure(39)	Application of SMV gel
Figure(40)	The perforated membrane was positioned over the defect
Figure(41)	Suture using polypropylene

Figure(42)	Line chart representing mean PI in the four studied groups
Figure(43)	Bar chart representing mean PI in the four studied groups
Figure(44)	Line chart representing mean GI in the four studied groups
Figure(45)	Bar chart representing mean GI in the four studied groups
Figure(46)	Line chart representing mean PD in the four studied groups
Figure(47)	Bar chart representing mean PD in the four studied groups
Figure(48)	Line chart representing mean CAL in the four
Figure(49)	Bar chart representing mean CAL in the four studied groups
Figure(50)	Line chart representing mean linear measurement/ mm (DBL) in the four studied groups
Figure(51)	Bar chart representing mean linear measurement/ mm (DBL) in the four studied groups
Figure(52)	Line chart representing mean linear measurement/ mm (CBL) in the four studied groups
Figure(53)	Bar chart representing mean linear measurement/ mm (CBL) in the four studied groups

Figure(54)	Line chart representing mean radiodensitometric measurement in the four studied groups
Figure(55)	Bar chart representing mean radiodensitometric measurement in the four studied groups
Figure(56)	Line chart representing mean SMV Conc. ($\mu\text{g/ml}$) in the four studied groups
Figure(57)	Bar chart representing mean SMV Conc. ($\mu\text{g/ml}$) in the four studied groups
Figure(58)	Correlation between linear measurement / mm (DBL) with clinical attachment level/mm and radiodensitometric measurement in each group
Figure(59)	Six months postoperative photograph showing 3 mm pocket depth in the same patient.
Figure(60)	Six months postoperative radiograph
Figure(61)	Nine months postoperative photograph showing 3 mm pocket depth in the same patient.
Figure(62)	Nine months postoperative radiograph
Figure(63)	Six months postoperative photograph showing 3 mm pocket depth in the same patient.
Figure(64)	Six months postoperative radiograph
Figure(65)	Nine months postoperative photograph showing 3 mm pocket depth in the same patient.
Figure(66)	Nine months postoperative radiograph