

# **SOME ADVANCED METHODS IN PROPAGATION OF SOME FRESH WATER FISH**

A Thesis

Presented to the Graduate school

Faculty of Veterinary Medicine, Alexandria University  
In partial fulfillment of the requirements for the Degree

of

Master of Veterinary Sciences

in

Animal Breeding and Production

By

**Emad Mohamed Zidan Farag**



# بعض الطرق المستحدثة فى تفريخ بعض أسماك المياه العذبة

## رسالة علمية

مقدمة إلى الدراسات العليا بكلية الطب البيطرى – جامعة الأسكندرية إستيفاء  
للدسات المقررة للحصول على درجة الماجستير فى العلوم الطبية البيطرية

## تخصص

## التربية والإنتاج الحيوانى

## مقدمة من

ط. ب / عماد محمد زيدان فرج

(٢٠٠٩)

# لجنة الإشراف

**الأستاذ الدكتور/ محمد محمد إبراهيم شرف**

أستاذ تربية وإنتاج الدواجن

رئيس قسم رعاية الحيوان وتنمية الثروة الحيوانية

كلية الطب البيطري – جامعة الإسكندرية

**الأستاذ الدكتور/ محمد عبدالباري منـدور**

أستاذ تربية وإنتاج الدواجن

قسم رعاية الحيوان وتنمية الثروة الحيوانية

كلية الطب البيطري – جامعة الإسكندرية

## ACKNOWLEDGEMENT

First of all my deep gratitude and indeptedness to **ALLAH** who gave me the ability to start, produce and finish this work.

I'm deeply indepted to **Prof. Dr. Mohamed Mohamed Ibrahim Sharaf**, Professor of Poultry Breeding and Production, Head of Department of Animal Husbandry and Animal Wealth Development – Faculty of Veterinary Medicine – Alexandria University for his kind supervision, encouragement, support and revision of this work.

My deep thanks are to be to **Prof. Dr. Mohamed Abd El – Bary Mandour**; Professor of Poultry Breeding and Production, Department of Animal Husbandry and Animal Wealth Development - Faculty of Veterinary Medicine – Alexandria University for his supervision, encouragement, continuous help, fruitful advices throughout the course of this work as well as revision of this work.

Special thanks go to **Dr. Waleed Nader El – Hawarry**; Lecturer of Fish Breeding and Production, Department of Animal Husbandry and Animal Wealth Development - Faculty of Veterinary Medicine – Alexandria University for his great help and continuous cooperation for fulfilling the practical part as well as all items of this work.

Many Cardial thanks to all members of Animal Husbandry and Animal Wealth Development, for their continuous encouragement and providing helpful facilities during this work.

## **SUPERVISOR'S COMMITTEE**

### **Prof. Dr. Mohamed Mohamed Ibrahim Sharaf**

Professor of Poultry Breeding and Production  
Head of Department of Animal Husbandry and Animal Wealth Development  
Faculty of Veterinary Medicine  
Alexandria University

### **Prof. Dr. Mohamed Abd El – Bary Mandour**

Professor of Poultry Breeding and Production  
Department of Animal Husbandry and Animal Wealth Development  
Faculty of Veterinary Medicine  
Alexandria University

# **CONTENTS**

	<b>Pages</b>
<b>List of Tables .....</b>	<b>1</b>
<b>List of Figures.....</b>	<b>II</b>
<b>List of Abbreviations.....</b>	<b>III</b>
<b>Introduction.....</b>	<b>1</b>
<b>♣ Review of Literature</b>	
<b>1. Reproductive behavior of tilapia.....</b>	<b>2</b>
<b>2. Seed production of Tilapia hapa system.....</b>	<b>3</b>
<b>2.1. Conditioning and exchange of broodstock in hapas.....</b>	<b>3</b>
<b>2.2. Stocking sex ratio of broodfish.....</b>	<b>6</b>
<b>2.3. Frequency of eggs / fry harvesting.....</b>	<b>7</b>
<b>2.4. Fecundity.....</b>	<b>8</b>
<b>2.5. Stocking density.....</b>	<b>9</b>
<b>2.6. Effect of <i>O. niloticus</i> female exchange on serum levels of sex steroid hormones.....</b>	<b>10</b>
<b>3. Water quality.....</b>	<b>10</b>
<b>4. Nutrition of broodstock.....</b>	<b>12</b>
<b>3. Seed production of tilapia in pond system.....</b>	<b>14</b>
<b>4. Comparison between nursing of fry in hapas and earthen ponds.....</b>	<b>14</b>
<b>♣ Materials and Methods</b>	
<b>I- Conditioning of <i>O. niloticus</i> female broodstock in hapas.....</b>	<b>16</b>
<b>I.1. Location and duration of the experiment.....</b>	<b>16</b>
<b>I.2. Experimental broodstock.....</b>	<b>16</b>
<b>I.3. Spawning and conditioning hapas.....</b>	<b>16</b>

I.4. Exchange schedule and period of conditioning.....	16
2. Feeding regime.....	25
3. Measurements.....	18
4. Growth performance parameters of broodfish.....	19
5. Water quality parameters.....	20
6. Blood sampling and analysis procedure.....	21
II. Comparison between production of fry in earthen spawning ponds and Hapas.....	22
III. Comparison of fry nursing in hapas and earthen ponds.....	23
♣ Results and Discussion	
1. Parameters of reproduction performance.....	25
1.1. Total fry production.....	25
1.2. Average number of fry female <sup>-1</sup> , average number of fry female <sup>-1</sup> spawn <sup>-1</sup> .....	28
1.3. Relative fecundity.....	33
1.4. Average fry number m <sup>-2</sup> day <sup>-1</sup> .....	38
1.5. Effect of water temperature on fry production.....	41
2. Growth performance of broodfish.....	44
3. Effect of exchange of <i>O. niloticus</i> female on serum levels of sex steroid hormones and total calcium in spawned and conditioned females.....	48
3.1. Effect of exchange of <i>O. niloticus</i> female on serum levels of sex steroid Hormones in spawned and conditioned females.....	48
3.2. Effect of exchange of <i>O. niloticus</i> female on serum levels of total calcium in spawned female fish removed from spawning hapas compared to females from conditioning hapas.....	51
II. Production of fry in hapas and earthen spawning ponds.....	54
III. Fry nursing in hapas and earthen ponds.....	56
Summary and Conclusion.....	60
References.....	61
Arabic summary.	



## 1- LIST OF TABLES

<b>Table 1.</b> Hapas number and period of conditioning in each treatment.....	25
<b>Table 2.</b> Different feed ingredients of basal diet used in broodstock feeding.....	26
<b>Table 3.</b> Chemical composition of feeds used in the trial conducted in hapas for Nile tilapia ( <i>O. niloticus</i> ) broodfish.....	26
<b>Table 4.</b> Weekly physical and chemical properties of the pond's water during the experimental period.....	29
<b>Table 5.</b> Area, stocking density and sex ratio of ponds.....	31

<b>Table 6.</b> LSM $\pm$ SD of total fry number of <i>O. niloticus</i> collected according to the treatments, no conditioning (T 0), 10days conditioning (T 1), 15 days conditioning (T 2) and 20 days conditioning (T 3).....	37
<b>Table 7.</b> LSM $\pm$ SD of average number of fry produced female <sup>-1</sup> collected in no conditioning (T 0), 10days conditioning (T 1), 15 days conditioning (T 2) and 20 days conditioning (T 3).....	40
<b>Table 8.</b> Average number of fry produced female <sup>-1</sup> spawn <sup>-1</sup> within and between treatments (LSM $\pm$ SD).....	42
<b>Table 9.</b> LSM $\pm$ SD of relative fecundity of females for no conditioning (T 0), 10days conditioning (T 1), 15 days conditioning (T 2) and 20 days conditioning (T 3).....	45
<b>Table 10.</b> LSM $\pm$ SD of relative fecundity day <sup>-1</sup> for no conditioning (T 0), 10days conditioning (T 1), 15 days conditioning (T 2) and 20 days conditioning (T 3).....	47
<b>Table 11.</b> LSM $\pm$ SD of average fry number m <sup>-2</sup> day <sup>-1</sup> for no conditioning (T 0), 10days conditioning (T 1), 15 days conditioning (T 2) and 20 days conditioning (T 3).....	50
<b>Table 12.</b> LSM $\pm$ SD of broodfish growth parameters, Initial, final body weight, body weight gain (BG g fish <sup>-1</sup> ), daily weight gain (DWG g fish <sup>-1</sup> ) and specific growth rate (SGR % day <sup>-1</sup> ).....	56
<b>Table 13.</b> Effects of exchange of Nile tilapia female on serum levels of testosterone (T) and estradiol (E <sub>2</sub> ) in spawned female fish that removed from spawning hapas (SH) and female fish removed from conditioning hapas (CH) after the experimental period (LSM $\pm$ SD).....	61
<b>Table 14.</b> Effects of exchange of Nile tilapia female on serum levels of total calcium in spawned females removed from spawning hapas (SH) compared to females removed from conditioning hapas (CH) after experimental period (LSM $\pm$ SD).....	64
<b>Table 15.</b> Total fry number and fry suitable for sex reversal produced from earthen spawning ponds throughout the experimental period.....	68
<b>Table 16.</b> Total number of fry, number of fry female <sup>-1</sup> , number of fry female <sup>-1</sup> spawn <sup>-1</sup> , fry number m <sup>-2</sup> day <sup>-1</sup> produced from earthen spawning ponds through the experimental period.....	68
<b>Table 17.</b> Means $\pm$ SD for initial and final weight, survival rate of four batches of <i>O. niloticus</i> fry, as influenced by nursing period in hapas and ponds.....	70
<b>Table 18.</b> Means $\pm$ SD for average body gain (g fish <sup>-1</sup> day <sup>-1</sup> ), average daily weight gain (g fish <sup>-1</sup> day <sup>-1</sup> ) and specific growth rate (% / day) of four batches of <i>O.niloticus</i> fry. As influenced by nursing period in hapas and ponds.....	71

## II – LIST OF FIGURES

<b>Figure 1.</b> Average total fry number of <i>O. niloticus</i> of different harvests.....	38
<b>Figure 2.</b> Total fry number of different treatments during the experimental period..	38
<b>Figure 3.</b> Average number of fry produced female <sup>-1</sup> of different harvests.....	41
<b>Figure 4.</b> Total number of fry produced female <sup>-1</sup> of different treatments.....	41
<b>Figure 5.</b> Average number of fry produced female <sup>-1</sup> spawn <sup>-1</sup> of different harvests...	43
<b>Figure 6.</b> Total number of fry produced female <sup>-1</sup> spawn <sup>-1</sup> of different treatments...	43
<b>Figure 7.</b> Relative fecundity of different harvests.....	46

<b>Figure 8.</b> Relative fecundity of different treatments.....	46
<b>Figure 9.</b> Average fry number gram <sup>-1</sup> day <sup>-1</sup> within treatments.....	48
<b>Figure 10.</b> Total number of fry gram <sup>-1</sup> female day <sup>-1</sup> of different treatments.....	48
<b>Figure 11.</b> Average fry number m <sup>-2</sup> day <sup>-1</sup> of different harvests.....	51
<b>Figure 12.</b> Total number of fry m <sup>-2</sup> day <sup>-1</sup> of different treatments.....	51
<b>Figure 13.</b> Effect of temperature on fry number in T1 (10 days conditioning) throughout the experiment.....	53
<b>Figure 14.</b> Effect of temperature on fry number in T2 (15 days conditioning) throughout the experiment.....	53
<b>Figure 15.</b> Effect of temperature on fry number in T3 (20 days conditioning) throughout the experiment.....	54
<b>Figure 16.</b> Effect of temperature on fry number in T0 (control) throughout the experiment.....	54
<b>Figure 17.</b> Body weight gain of male and female broodfish of different treatments..	57
<b>Figure 18.</b> Daily weight gain of male and female broodfish.....	57
<b>Figure 19.</b> Specific growth rate of male and female broodfish of different treatments.....	58
<b>Figure 20.</b> Effects of exchange of Nile tilapia female on serum levels of testosterone in spawned female fish removed from spawning hapas (SH) and female fish removed from conditioning hapas (CH) after experimental period.....	62
<b>Figure 21.</b> Effects of exchange of Nile tilapia female on serum levels of estradiol in spawned female fish removed from spawning hapas (SH) and female fish removed from conditioning hapas (CH) after experimental period.....	62
<b>Figure 22.</b> Effects of exchange of Nile tilapia female on serum levels of total calcium in spawned female removed from spawning hapas (SH) compared to female removed from conditioning hapas (CH).....	65
<b>Figure 23.</b> Average body weight gain of fry as influenced by nursing period in hapas and ponds.....	72

<b>Figure 24.</b> Average daily body weight gain of fry as influenced by nursing period in hapas and ponds.....	72
---	----

<b>Figure 25.</b> Specific growth rate of fry as influenced by nursing period in hapas and ponds.....	73
---	----

### III – LIST OF ABBRVIATIONS

**Bw: Body weight**

**Fed : feddan**

**m<sup>-2</sup> : per square meter**

**m<sup>2</sup> : square meter**

**SH :spawning hapa**

**CH :conditioning hapa**

**M : meter**

**DWG :Daily weight gain**

**SGR: specific growth rate**

**g fish<sup>-1</sup>: gram per fish**

**g: gram**

**Batch<sup>-1</sup> : per batch**

**Day<sup>-1</sup>: per day**

**SR :survival rate**

**μl: microliter**

**T: testosterone**

**E<sub>2</sub> : 17 beta estradiol**

**ng: nanogram**

**pg: pictogram**

**ml<sup>-1</sup> : per milli**

**mg dl<sup>-1</sup>: mg per deciliter**

# 1- LIST OF TABLES

<b>Table 1.</b>	Number of hapas and period of conditioning per treatment.....	١٧
<b>Table 2.</b>	Feed ingredients of basal diet used in broodstock feeding.....	١٨
<b>Table 3.</b>	Chemical composition of feed used in the trial conducted in hapas for Nile tilapia ( <i>O. niloticus</i> ) broodfish.....	١٨
<b>Table 4.</b>	Area, stocking density and sex ratio specified for ponds.....	٢١
<b>Table 5.</b>	LSM $\pm$ SD of total fry number of <i>O. niloticus</i> collected according to the treatments, Non conditioned ( $T_0$ ), 10 days conditioning ( $T_1$ ), 15 days conditioning ( $T_2$ ) and 20 days conditioning ( $T_3$ ).....	٢٦
<b>Table 6.</b>	LSM $\pm$ SD of average number of fry produced female <sup>-1</sup> collected in non conditioned ( $T_0$ ), 10 days conditioning ( $T_1$ ), 15 days conditioning ( $T_2$ ) and 20 days conditioning ( $T_3$ ) of <i>O. niloticus</i> .....	29
<b>Table 7.</b>	Average number of fry produced female <sup>-1</sup> spawn <sup>-1</sup> within and between (LSM $\pm$ SD) of <i>O. niloticus</i> treatments	٣١
<b>Table 8.</b>	LSM $\pm$ SD of relative fecundity of females for non conditioning ( $T_0$ ), 10 days conditioning ( $T_1$ ), 15 days conditioned ( $T_2$ ) and 20 days conditioning ( $T_3$ ) of <i>O. niloticus</i> .....	٣٤
<b>Table 9.</b>	LSM $\pm$ SD of relative fecundity day <sup>-1</sup> for non conditioned ( $T_0$ ), 10 days conditioning ( $T_1$ ), 15 days conditioning ( $T_2$ ) and 20 days conditioning ( $T_3$ ) of <i>O. niloticus</i> .....	٣٦
<b>Table 10.</b>	average fry number m <sup>-2</sup> day <sup>-1</sup> for non conditioned ( $T_0$ ), LSM $\pm$ SD of 10 days conditioning ( $T_1$ ), 15 days conditioning ( $T_2$ ) and 20 days conditioning ( $T_3$ ) of <i>O. niloticus</i> .....	٣٩
<b>Table 11.</b>	Weekly physical and chemical properties of the pond's water during the experimental period.....	٤٥
<b>Table 12.</b>	LSM $\pm$ SD of broodfish growth parameters, initial, final body weight, body weight gain (BG g fish <sup>-1</sup> ), daily weight gain (DWG g fish <sup>-1</sup> ) and specific growth rate (SGR % day <sup>-1</sup> ) of <i>O. niloticus</i> .....	٤٦
<b>Table 13.</b>	Effects of exchange of <i>O. niloticus</i> female on serum levels of testosterone (T) and estradiol (E <sub>2</sub> ) in spawned female fish that	