



PROCALCITONIN IN REDUCING THE DURATION OF ANTIBIOTIC THERAPY IN SEPTIC CRITICALLY ILL PATIENTS

Thesis

Submitted for Partial Fulfillment of M.D. Degree
in Anaesthesiology

Presented By

Amr Mohamed El-Sayed Ahmed Shaaban

*M.B.B. Ch, M.Sc., Anaesthesiology
Faculty of Medicine, Ain Shams University*

Supervised By

Prof. Dr. Bahaa Eldin Ewees Hassan

*Professor of Anesthesia, Intensive Care and pain management
Faculty of Medicine - Ain Shams University*

Dr. Ahmed Mohamed Elsayed Elhenawi

*Assistant Professor of Anesthesia, Intensive Care and pain management
Faculty of Medicine - Ain Shams University*

Dr. Mai Mohsen Abdelaziz

*Lecturer of Anesthesia, Intensive Care and pain management
Faculty of Medicine - Ain Shams University*

**Faculty of Medicine
Ain Shams University
2016**



وَأَنْزَلَ اللَّهُ عَلَيْكَ
الْكِتَابَ وَالْحِكْمَةَ
وَعَلَّمَكَ مَا لَمْ تَكُن تَعْلَمُ
وَكَانَ فَضْلُ اللَّهِ عَلَيْكَ
عَظِيمًا

صِرَاقُ اللَّهِ الْعَظِيمِ

سُورَةُ النِّسَاءِ (الْآيَةُ ١١٢)



*First and above all thanks to **ALLAH**.*

I would like to express my endless gratitude and appreciation to my eminent professor,

***Prof. Dr. Bahaa Eldin Ewees Hassan**, Professor of Anesthesia, Intensive Care and pain management, Faculty of Medicine - Ain Shams University, for giving me the honor to work under her supervision and from whom I did learn a lot. She encouraged me, removed all the obstacles from my way and pushed me to achieve success.*

*My sincere thanks to **Dr. Rasha Samir Bondok**, Professor of Anesthesia, Intensive Care and pain management, Faculty of Medicine - Ain Shams University, for her continuous guidance, honest help and endurance that made this thesis come to light.*

*I would also like to express my gratefulness to **Dr. Ahmed Mohamed Elsayed Elthenawi**, Assistant Professor of Anesthesia, Intensive Care and pain management, Faculty of Medicine - Ain Shams University, for his precious advice and support throughout this whole work.*

*I thank also, **Dr. Mai Mohsen Abdelaziz**, Lecturer of Anesthesia, Intensive Care and pain management, Faculty of Medicine - Ain Shams University, for her efforts in this study.*

✎ Amr Mohamed El-Sayed Ahmed Shaaban

List of Contents

Title	Page No.
List of Abbreviations.....	ii
List of Tables	iv
List of Figures	v
Introduction	1
Aim of the work.....	3
<u>Review of Literature</u>	
• Sepsis.....	4
• Clinical Biomarkers of Sepsis.....	46
Patients and Methods	59
Results	65
Discussion.....	77
Summary	88
Conclusion.....	90
References	91
Arabic Summary	--

List of Abbreviations

ALI	Acute lung injury
ARDS	Acute respiratory distress syndrome
BBB	Blood-brain barrier
CDAD	Clostridium difficile-associated diarrhea
CRP	C-reactive protein
DIC	Disseminated intravascular coagulation
ED.....	Emergency department
ESBL	Extended-spectrum beta lactamases
FUO	Fever of unknown origin
HPV	Hypoxic pulmonary vasoconstriction
ICU	Intensive care unit
IL	interleukins
MDR	Multi-drug resistant
MODS	Multiple organ dysfunction syndrome
PCT	Procalcitonin
RTIs	Respiratory tract infections
SAPS	Simplified Acute Physiology Score
SD	Standard deviation
SIRS	Systemic inflammatory response syndrome
SOFA	Sequential organ failure assessment score

List of Abbreviations (Cont...)

SPSS	Statistical program for social science
TNF- α	Tumor necroting factor
VRE	Vancomycin-resistant Enterococcus
WBC	White Blood cell count
χ^2	Chi-square

List of Tables

Table No.	Title	Page No.
Table 1:	Sequential [Sepsis-Related] Organ Failure Assessment Score	7
Table 2:	Initial Antimicrobial Therapy for Severe Sepsis with No Obvious Source in Adults with Normal Renal Function	40
Table 3:	Use of procalcitonin levels for antimicrobial stewardship of respiratory tract infections	57
Table 4:	Comparison between groups according to the demographic data.....	65
Table 5:	Comparison between groups according SAPS II score on admission.	67
Table 6:	Comparison between groups according to SOFA score on admission.	68
Table 7:	Comparison between groups according to blood cultures on admission.....	69
Table 8:	Comparison between groups according to serial PCT levels.	70
Table 9:	Comparison between groups regarding duration of Ab treatment.....	72
Table 10:	Comparison between groups according to days alive without Ab.....	73
Table 11:	Comparison between groups according to ICU length of stay.....	74
Table 12:	Comparison between groups according to 28-day mortality.....	75
Table 13:	Comparison between groups regarding clinical cure.....	76

List of Figures

Fig. No.	Title	Page No.
Fig. (1):	The Host Response in Severe Sepsis.....	15
Fig. (2):	Organ Failure in Severe Sepsis and Dysfunction of the Vascular Endothelium and Mitochondria.....	24
Fig. (3):	Structure of Procalcitonin.....	49
Fig. (4):	Bar chart between groups according to age (years).	65
Fig. (5):	Bar chart between groups according to sex.	66
Fig. (6):	Bar chart between groups according SAPS II score.	67
Fig. (7):	Bar chart between groups according to SOFA score.	68
Fig. (8):	Bar chart between groups according blood cultures.	69
Fig. (9):	Showing chart between groups according PCT.	71
Fig. (10):	Bar chart between groups according duration of Ab. TTT.	72
Fig. (11):	Bar chart between groups according to days alive without Ab.	73
Fig. (12):	Bar chart between groups according to ICU length of stay.....	74
Fig. (13):	Bar chart between groups according outcome.	75
Fig. (14):	Bar chart between groups according clinical care.	76

INTRODUCTION

The duration of antibiotic therapy in critically ill patients with sepsis is based on empirical rules (*Calandra & Cohen, 2005*). It can result in antibiotic overuse, increasing the risk of developing multi-drug resistant (MDR) pathogens, nosocomial superinfections and excess cost (*Elizabeth, 2011*).

Recent attempts to decrease the duration of the antibiotic therapy have been successful in critically ill patients (*Nobre et al., 2008*). Patients with ventilator-associated pneumonia, were found to be equally improved when receiving an 8-day treatment compared to a 15-day course of antibiotics (*Chastre et al., 2003*). This is, however, again based on empirical rules. Guidance of the duration of the antibiotic treatment tailored for each patient based on simple biomarkers could contribute to an additional benefit (*Nobre et al., 2008*).

Newer strategies aimed at identifying patients most likely to benefit from antibiotics and those that may require shorter than standard treatments using clinical biomarkers to assess presence, severity, and progression of disease like C-reactive protein (CRP), white blood cell count (WBC), interleukins (IL) and most recently, procalcitonin (PCT) has become a target tool to improve antimicrobial stewardship (*Schuetz et al., 2009*).

Procalcitonin (PCT), the biologically active precursor of the calcium-modulating hormone calcitonin (*Meisner 2002*) has been shown in diverse studies to be closely associated with the human host response to bacterial infection (*Castelli et al., 2004*). It is elaborated by parenchymal cells throughout the body in response to endotoxin and several pro-inflammatory mediators (in particular TNF- α) and its concentration appears to be roughly linear with the degree of insult (*Brunkhorst et al., 2000*). It has tantalizing characteristics as a biomarker for bacterial infection, showing diagnostic superiority to white cell count, C-reactive protein, and a host of physiologic variables in most reports (*Wanner et al., 2000; Nobre et al., 2008*).

AIM OF THE WORK

The study aims to evaluate the efficacy of serum procalcitonin as a biomarker in reducing the duration of antibiotics treatment in septic critically ill patients.

SEPSIS

Sepsis which is derived from the Greek verb *sepo* (meaning “I rot”) is a systemic, deleterious host response to infection leading to severe sepsis (acute organ dysfunction secondary to documented or suspected infection) and septic shock (severe sepsis plus hypotension not reversed with fluid resuscitation). Severe sepsis and septic shock are major healthcare problems, Similar to polytrauma, acute myocardial infarction, or stroke, the speed and the appropriateness of therapy administered in the initial hours after severe sepsis develops are likely to influence the outcome (*Kaukonen et al., 2015*).

Definitions

Infection: defined as a pathological process caused by the invasion of a normally sterile tissue or fluid or body cavity by pathogenic or potentially pathogenic micro-organisms. It is also important to point out that, frequently, infection is strongly suspected without being microbiologically confirmed (*Dellinger et al., 2013*).

Systemic inflammatory response syndrome (SIRS): is an inflammatory reaction that produces at least two of the following four signs:

- Abnormal body temperature: hypothermia, $<96.8^{\circ}\text{F}/36^{\circ}\text{C}$; or fever, $>100.4^{\circ}\text{F}/38.3^{\circ}\text{C}$.
- Tachycardia (>90 beats/min),
- Tachypnea (>20 breaths/min or a rate sufficient to produce $\text{PaCO}_2 <32$ mm Hg),
- Abnormal white blood cell count ($>12,000/\text{mm}^3$, $<4000/\text{mm}^3$, or $>10\%$ immature forms) (*Gaieski and Goyal, 2013*).

SIRS due to abnormal white blood cell count can be triggered by an infection, but also it can arise from non-infectious sources such as trauma, haemorrhage, burns, surgery, adrenal insufficiency, pulmonary embolism, dissecting or ruptured aortic aneurysm, myocardial infarction, occult haemorrhage, cardiac tamponade, post-cardiopulmonary bypass syndrome, autoimmune disorders, pancreatitis, vasculitis, anaphylaxis, or drug overdose. SIRS can lead to organ failure, shock, and death (*Neviere, 2013*).

Severe sepsis: “sepsis syndrome, ” is present when the patient has progressed to a stage in which organs or

organ systems begin to fail. Severe sepsis is sepsis plus one of the following clinical problems: Cardiovascular system dysfunction, acute respiratory distress syndrome (ARDS), dysfunction of two or more other organs or systems (*Dellinger et al., 2013*).

Septic shock: in a patient with sepsis, is an acute circulatory failure with refractory hypotension which is unexplainable by other causes. The term *shock* describes a condition in which many tissues throughout the body become hypoxic due to poor perfusion. In shock, normal homeostatic mechanisms are either not functioning or not adequate to deliver enough oxygen to tissues. If it is not reversed it leads to organ failures and death (*Gaieski and Goyal, 2013*).

In septic shock, there is hypotension that cannot be reversed by giving adequate fluids. When the hypotension of septic shock also does not respond to vasopressors, the condition is called *refractory septic shock* (*Munford and Suffredini, 2009*).

Multiple organ dysfunction syndrome (MODS): failure of a number of organs or organ systems caused by an illness. To be considered “in failure” an organ must persist in its severe dysfunction for at least 24 hours (*Dellinger et al., 2013*).

The lack of a reliable definition of sepsis makes assessment of incidence and changes in outcomes difficult to quantify reliably leads to a new definition of sepsis (***The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3)***) which recommends the following:

Sepsis should be defined as life-threatening organ dysfunction caused by a dysregulated host response to infection. For clinical operationalization, organ dysfunction can be represented by an increase in the Sequential [Sepsis-related] Organ Failure Assessment (***SOFA***) score of 2 points or more (***see table 1***), which is associated with an in-hospital mortality greater than 10 %(***Mervyn et al., 2016***).

Table 1: Sequential [Sepsis-Related] Organ Failure Assessment Score (***Mervyn et al., 2016***).