تأثير التداخل بين بعض العناصر و المواد الدهنية على فئران التجارب

رسالة مقدمة من

نجلاء عزت ابوسمرة بكالوريوس العلوم الزراعية (صناعات غذائية)- كلية الزراعة _ جامعة القاهرة- ١٩٨٩

للحصول على

درجة الماجستير

في

العلوم الزراعية (الكيمياء الحيوية)

قسه الكيمياء الحيوية كليه كليه الزراعة جامعة القاهرة مصسر

Y . . A

تأثير التداخل بين بعض العناصر و المواد الدهنية على فئران التجارب

رسالة ماجستير في العلوم الزراعية (الكيمياء الحيوية)

مقدمة من

نجلاع عزت ابوسمرة بكالوريوس في العلوم الزراعية (الصناعات الغذائية)- كلية الزراعة - جامعة القاهرة- ١٩٨٩

لجنة الإشراف

الدكتور/ جمال سيد علي الباروطي أستاذ الكيمياء الحيوية - كلية الزراعة - جامعة القاهرة

الدكتور/ مجدى عبد العليم شعلان أستاذ الكيمياء الحيوية – كلية الزراعة - جامعة القاهرة

الدكتور/ احمد السيد بسيونى رئيس بحوث علوم وتكنولوجيا الأغذية مركز البحوث النجوث الزراعية الجيزة

تأثير التداخل بين بعض العناصر و المواد الدهنية على فئران التجارب

رسالة ماجستير في العلوم الزراعية (الكيمياء الحيوية)

مقدمة من

نجلاع عزت ابوسمرة بكالوريوس في العلوم الزراعية (الصناعات الغذائية)- كلية الزراعة - جامعة القاهرة ١٩٨٩

لجنة إجازة الرسالة: أ.د / وجيه احمد السيد المالكي رئيس بحوث الكيمياء الحيوية والتغنية – الهيئة القومية للرقابة والبحوث الدوائية

> أ.د/ عادل محمد ابو الفتوح استاذ الكيمياء الحيوية – كلية الطب البيطرى – جامعة القاهرة

أ.د/ مجدى عبد العليم شعلان استاذ الكيمياء الحيوية – كلية الزراعة – جامعة القاهرة

أ.د/ جمال سيد على الباروطى استاذ الكيمياء الحيوية - كلية الزراعة - جامعة القاهرة بتاريخ: / /

EFFECTS OF THE INTERACTION BETWEEN SOME MINERALS AND LIPIDS IN RATS

By

NAGLAA EZZAT ABOU SAMMRA

B.Sc. Agric. Sci. (Food Sci. & Tech.), Fac. Agric., Cairo University, 1989

THESIS

Submitted in Partial Fulfillment of the Requirements for the Degree of

MASTER OF SCIENCE

In

Agricultural Sciences (Agricultural Biochemistry)

Department of Biochemistry
Faculty of Agriculture
Cairo University
EGYPT

2008

APPROVAL SHEET

EFFECTS OF THE INTERACTION BETWEEN SOME MINERALS AND LIPIDS IN RATS

M. Sc. Thesis By

NAGLAA EZZAT ABOU SAMMRA

B.Sc. Agric. Sci. (Food Sci. & Tech.), Fac. Agric., Cairo Univ., 1989.

Approved by:

Dr.WAGIH AHMAD EL-MALKY

Head Researcher of Biochemistry, National Oraganization for Drug Control and Research

Dr. ADEL MOHAMED ABO EL-FETOUH

Professor of Biochemistry, Fac. Veterinary Medicine, Cairo University.

Dr. MAGDY ABDEL ALEEM SHALLAN

Professor of Agricultural Biochemistry, Fac. Agric., Cairo University.

Dr. GAMAL EL-SAYED ALY EL-BAROTY

Professor of Agricultural Biochemistry, Fac. Agric., Cairo University.

Date: / /2008

SUPERVISION SHEET

EFFECTS OF THE INTERACTION BETWEEN SOME MINERALS AND LIPIDS IN RATS

M. Sc. Thesis By

NAGLAA EZZAT ABOU SAMMRA

B.Sc. Agric. Sci. (Food Sci. & Tech.), Fac. Agric., Cairo Univ., 1989.

SUPERVISION COMMITTEE

Dr. Gamal S. El-Baroty
Professor of Biochemistry, Faculty of Agriculture, Cairo University

Dr. Magdy A. Shallan Professor of Biochemistry, Faculty of Agriculture, Cairo University

Dr. Ahmed El. S. Basuony Head Researcher of Food Sci. and Technology, Food Sci. and Technology Research Institute, Agriculture Research Center, Giza. Name of Candidate: Naglaa Ezzat Abou Sammra Degree: M.Sc.

Title of Thesis: Effects of Interaction Between Some Minerals and Lipids In

Rats.

Supervisors: Dr. Gamal S. El-Baroty, Dr. Magdy A. Shallan and Dr.

Ahmed El. S. Basuony

Department: Agricultural Biochemistry

Branch: Agricultural Biochemistry Approval: \ \2008

ABSTRACT

Interactions of dietary nutrient are important factors to consider in the study. The aim of this study was to investigate the effect of dietary interacting among iron (Fe), manganese (Mn) and two fatty types on growth parameter (body weight gain, food intake, food efficiency ratio and orgain weight), lipid characterization (as mesured by TL, TG, T-Cho, HDL and LDL) enzyme activities (CAT, POX and SOD), mineral status (as measured with Fe and Mn levels), lipid peroxidation (as measured by TBARs) and protein. Male rats (n = 6) were used in 3 x 3 x 2 factorial design and fed standard basal diet containing dietary Fe (10, 35 and 250 mg/kg), Mn (1, 45 and 250 mg/kg) and either soybean or corn oil at 15% for 8 weeks. The results of the present work can be summarized as follows:-

Body weights did not differ throughout the experiment among rats fed the differents diets. The body weight gain, food intake, food efficiency ratio and orgain weight of all rats groups were not significantly different. Thus, these parameters were not affected by the levels of dietary iron and manganese or the type of oils. Also, the Anova analysis did not show relationship between the increase of rat organs (heart, lung, kidney and spleen) and the dietry Fe, Mn and oil types. The concentration of Fe or Mn in plasma was increased through the experiment period among rat's feddifferent diets. The plasma Fe and Mn contents were gradually increased by feeding rats at higher levels of both minerals statistical analysis, showed that the plasma Fe and Mn was positively associated with Fe and Mn levels in diets.

As intake of Fe and Mn increased plasma, HDL was increased. Incontrast, concentration of plasma TL, LDLand T-Chol were decreased throughout experimental periods. In over all, the changes in TL content was affected with the time, Fe-dose, interaction between Fe and Mn and interaction among time, Fe and Mn. However, the changes in TL were altered throughout experimental periods, and these change in TL values was differ from week to another.

Rats fed high iron diets had higher concentration of lipid peroxidation products in plasma (thiobarbituric acid – reactive substances, TBRAs). TBRAs valus in plasma was increased gradually with the time. The rats fed diets contained Mn, Fe with soybean and corn oils leading to increase antioxidant enzyme activity including: SOD, CAT and POX in plasma throughout experimental period, The ANOVA analysis revealed that activities of SOD, CAT and POX were significantly affected by dietary Fe and Mn.From overall results it could be concluded that diets containing Fe at adequt level (35 mg / kg diet) or Mn could be save for supplementation of some foods to preventy my deficiency signs caused by both elementshave been found.

Key words: Iron, Manganese, Chemical Studies, Biological Studies, Rat

اسم الطالب: نجلاء عزت ابوسمرة الدرجة: الماجستير

عنوان الرسالة: تأثير التداخل بين بعض العناصر و المواد الدهنية على فئران التجارب

المشرفون: الأستاذ الدكتور: جمال سيد علي الباروطي

الأستاذ الدكتور: مجدى عبد العليم شعلان

الدكتور : الحمد سيد بسيوني

قسم: الكيمياء الحيوية فرع: الكيمياء الحيوية تاريخ منح الدرجة: / /

المستخلص العربى

تعتبر دراسة تداخل العناصر من العوامل الهامة و المؤثرة في حالة المحافظة على الصحة العامة فيلعب الغذاء المتوازن دور هام في الوقاية من امراض سوء التغذية والتي تشمل نقص أو زيادة عنصر من المواد الغذائية و التي تؤثر على صحة و حياة الفرد فأحيانا نقص احد العناصر يؤدى الى سوء التغذية بسبب نقصه في كمية الغذاء و احيانا الأفراط في تناول العناصر الغذائية يؤدى الى ظهور مشاكل صحية قد تمثل خطورة على حياة الفرد.

تهدف هذه الدراسة الى تقييم تأثير تداخل عنصرى الحديد و المنجنيز و نوعان مختلفان من الزيوت (زيت الصويا وزيت الذرة) على مستوى اللبيدات والمعادن على ذكور فئران التجاربحيث أجريت تجربة بيولوجية لدراسة مدى تأثير تداخل هذه العناصر مع اللبيدات على البلازما وكبد الفئران تم استخدم عدد 108من الفئران و قسمت الى مجموعتين مجموعة تحتوى على 37 كيزات مختلفة من الحديد و المنجنيز مع زيت الذرة بتركيز 15% و المجموعة الأخرى تحتوى على 37 كيزات مختلفة من الحديد و المنجنيز مع زيت الذرة بتركيز 15% و كانت التركيزات المستخدمة لعنصر الصويا بتركيز 10% و كانت التركيزات المستخدمة لعنصر المنجنيز هي (1-45-250ملجم/كجم) و قد استمرت التجربة لمدة 60يوما و قد تم أخذ عينات من الدم خلال فترة التجربة مع تقدير وزن الفئران لمتابعة نموها كل أسبوعين و في نهاية التجربة تم ذبح الفئران وأخذت الاعضاء المختلفة (الكبد-الطحال-القلب-الكلية) و معرفة الاوزان المختلفة لهذه الاعضاء وقد تم تقدير قياسات النمو للفئران (الزيادة في الوزن- الكفاءة الغذائية- كمية الغذاء المستهلك)وقد تم تقدير نشاط انزيمات المضادة للاكسدة (البيروكسيديز و الكتاليز والكتاليز والسوبر اوكسيد دسميوتيز) وتم تقدير محتوي المواد اللبيدية بالدم وانسجة الكبد . وقد تم تقدير الكوليستيرول واللبيدات الكلية والليبوبروتينات منخفضة وعالية الكثافة . وكذلك تقدير والـ (TBARs). وتقدير البروتين وصورة البروتينات المختلفة و من خلال التجربة كانت اهم التنتائج ما يلي:

- ملاحظة حدوث زيادة في اوزان فئران التجارب خلال التجربة مع حدوث زيادة في اوزان الفئران المغذاة على زيت فول الصويا عن الفئران المغذاة على زيت الذرة
 - تم ملاحظة زيادة نسبة عنصري الحديد والنجنيز في بلازما الدم في مجاميع الفئران المغذاة على العلائق المختلفة
 - -عدم حدوث اى تغير في نسب البروتينات في حالة المعاملة بكلا العنصرين
 - -وجد ان نتيجة المعاملة باي من العنصرين في وجود كلا من انواع الزيوت المختلفة ادى الي زيادة نشاط الانزيمات المضادة للاكسدة مثل الكتاليز ولبيروكسيبديز
 - وجد ان بزيادة تركيز كلا من العنصرين في العليقة يؤدي ذلك الي نقص في مستوي اللبيدات الكلية و الكوليسترول و الجلسريدات الكلية وعلى العكس زادة نسبة رقم الثيوبارباتيوريك
- من تلك النتائج يمكن القول بان استخدام علائق تحتوي على تركيزات متوسطة 35 ملجم/كجم عليقة من الحديد و 45 ملجم / كجم عليفة من النجنيز تعتبر امنة تماما في الاستخدام من الناحية الغذائية ومكملة في حالة نقص اي من العنصرين

الكلمات الدالة: الحديد - المنجنيز - دراسات كيميائية - دراسات بيولوجي - فنران التجارب

CONTENTS

	Page
INTRODUCTION	1
REVIEW OF LITERATURE	3
1. Iron	3
a. The risks of excessive iron storage	4
b. Catalysis of hydroxyl radical production	5
c. The initiation of lipid peroxidation	6
d. Biological function of iron	7
e. Heme proteins	10
f. Single iron containing metalloenzymes	11
g. Protein with oxygen- bridged iron	11
h. Proteins with iron-sulfur centers	12
i. Iron excess	13
j. Iron deficiency	15
k. Liabilities of iron deficiency	16
2. Effect of iron on metabolism	17
3. Effect of iron on lipid composition and metals status	20
4. Effect of iron on lipid peroxidations	27
5. Manganese	33
6. Effect of manganese on rats biochemical	35
parameters	
a. Enzyme	35
b. Mineral contents and lipid profile	39
c. Effect of manganese on lipid peroxidations	41
d. Effect of manages on lipid constitution	42
7. Iron and manganese Interaction	44
MATERIALS AND METHODS	50
1. Materials	50
a. Oils	50
b. Animals	50
c. Analytical kits	50
d. Solvents and chemicals	50
2. Nutrition experiments	51
3. Blood Sampling	53
4. Preparation of hepatic homogenate	53
5 Riochemical analysis and enzyme assays	54

a. Determination of total lipids in plasma	54
b. Determination of plasma total cholesterol	55
c. Determination of high density lipoprotein-	
cholesterol	56
d. Determination of Low density lipoprotein-cholesterol	57
e. Determination of plasma triglycerides	57
f. Determination of lipid peroxidation products	58
6. Determination of total protein in plasma and	59
liver	
7. Determination of enzymes specific activities	60
a. Antioxidant enzymes	60
1. Catalase	60
2. Peroxidase	62
3. Measurement of superoxide dismutase	63
8. Determination of minerals contents	65
9. Determination of Iron in plasma	66
10. Fractionation of protein	67
11. Isolation and extraction of fatty acid	71
a. Methylation of fatty acids	72
b. Identification of fatty acid methyl esters	72
c. Analysis of fatty acid in rat dietary diets	73
12. Statistical analysis	73
RESULTS AND DISCUSSION	74
SUMMARY	120
REFERENCES	123

7

LIST OF TABLES

No	Title	Page
1.	Types and examples of iron-containing proteins	8
2.	Composition of the basal diet	52
3.	Composition of the salt mixture contents	52
4.	Composition of vitamin mixture	52
5.	The levels of Fe and Mn in different basal diets	53
6.	Compostion of separating and stacking gels	70
7.	GLC analysis of fatty acids methylester of corn oil and soybean oils	
8.	Growth parameter of rats fed diets containing various levels of Manganese and iron with corn oil	74
9.	Growth paramater of rats fed diets containing various levels of manganese and iron with soybean oil	75
10.	Changes in weight of rats fed diet containing Fe/Mn and corn oil during the eight weeks (experimental period)	
11.	Changes in weight of rats fed diet containing Fe/Mn and soybean oil	
12.	during the eight weeks (experimental period)	78
10	soybean oil types	79
13.	Changes in Fe concentration in plasma of rats fed diet containing Fe/Mn and corn oil during the eight weeks (experimental period)	81
14.	Changes in Fe concentration in plasma of rats fed diet containing Fe/Mn and soybean oil during the	
	eight	82

	15.	Changes in Mn concentration in plasma of rats fed diet containing Fe/Mn and corn oil during the eight weeks	82
	16.	Changes in Mn concentration in plasma of rats fed diet containing Fe/Mn and soybean oil during the eight weeks	83
	17.	Changes in FE and Mn concentration in liver of rats fed diet containing Fe/Mn and soybean oil during the eight weeks	84
	18.	Changes in T.L. in plasma of rats fed diet containing Fe/Mn and corn oil during the eight weeks	85
	19.	Changes in T.L. in plasma of rats fed diet containing Fe/Mn and soybean oil during the eight weeks	86
	20.	Changes in T.L. in liver of rats fed diet containing Fe/Mn	87
	21.	Changes in T.G. in plasma of rats fed diet containing Fe/Mn and corn oil during the eight weeks	88
	22.	Changes in T.G. in plasma of rats fed diet containing Fe/Mn and soybean oil during the eight weeks	88
	23.	Changes in T.G. in liver of rats fed diet containing Fe/Mn	89
	24.	Changes in cholesterol in plasma of rats fed diet containing Fe/Mn and corn oil during the eight weeks	90
	25.	Changes in cholesterol in plasma of rats fed diet containing Fe/Mn and soybean oil during the eight weeks	91
	26.	Changes in cholesterol in liver of rats fed diet containing Fe/Mn	
	27.	Changes in HDL in plasma of rats fed diet containing Fe/Mn and corn oil during the eight weeks	94
V	28.	Changes in HDL in plasma of rats fed diet containing Fe/Mn and soybean oil during the eight	95
		IV	