

بسم الله الرحمن الرحيم

(وَقُلْ اَعْمَلُوا فَسَيَرَى اللّٰهُ عَمَلَكُمْ وَرَسُولُهُ وَالْمُؤْمِنُونَ وَسَتُرَدُّونَ اِلَىٰ عَالَمِ  
الْغَيْبِ وَالشَّهَادَةِ فَيُنَبِّئُكُمْ بِمَا كُنْتُمْ تَعْمَلُونَ)

(التوبة:105)

**Magnetic Resonance Imaging Study of**  
**Temporomandibular Joint**  
**in Yemeni Qat Chewers**

A Thesis Submitted to Faculty of Oral and Dental  
Medicine, Cairo University in partial  
Fulfillment of the Requirements  
For the Master Degree  
In Oral Radiology

By

**Abeer Abdulkareem Al-Mashraqi**  
(BDS 2004)  
Sana'a University

Faculty of Oral and Dental Medicine  
Cairo University  
2010

## **Supervisors**

**Dr. Nashwa Salah Mohamed**

Professor of Oral Radiology  
Faculty of Oral and Dental Medicine  
Cairo University

**Dr. Eman Abd Elaziz Ahmed**

Professor of Oral Radiology  
Faculty of Oral and Dental Medicine  
Cairo University

# DEDICATION

## **DEDICATION**

*To my parents who supported me,*

*To my husband for his patience and support,*

*To my beloved daughter.*

# ACKNOWLEDGEMENT

## **ACKNOWLEDGEMENT**

I would like to express my deep appreciation and gratitude to my supervisors **Prof. Dr. Nashwa Salah and Prof. Dr. Eman Abd Elaziz**, Professors of Oral Radiology, Faculty of Oral and Dental Medicine, Cairo University, for their guidance, encouragement and enlightening vision throughout the course of this study. **Dr. Nashwa Salah and Dr. Eman Abd Elaziz** not only trained me on the scientific research methods, but also exemplified the true meaning of professors. It would have been impossible to accomplish this work without their help.

It is with great pleasure that I thank the following people who have been the instrumental in the completion of this work:

**Dr. Entesar Kaheri**, Assistant Professor in Oral Medicine Department, Faculty of Oral and Dental Medicine, Ibb University, for her assistance and encouragement during sample collection.

**Prof. Dr. Aiman Ali**, Professor in Oral Medicine Department, Faculty of Oral and Dental Medicine, King Faisal University, for his help and support.

My devoted **parents** and my **husband**, for their never-ending patience and support.

All **subjects** who participate in this work and helped in bringing this research into existence.

Saudi German Hospitals Group. Sana'a, for their cooperation and help to do this work.

Last but not least, I would like to thank all the staff members of the Department of Oral Radiology, Faculty of Oral and Dental Medicine, Cairo University, for their encouragement.

## **LIST OF TABLES**

<b>Number</b>	<b>Table</b>	<b>Page</b>
1	The frequencies and percentages of clinical findings in qat chewers group.	67
2	The frequencies, percentages and results of McNemar's test for the comparison between clinical findings in the two sides of qat chewers group.	68
3	The frequencies, percentages and results of chi-square test for comparison between disc position, arthritic changes and JE in qat chewers and control groups.	69
4	The frequencies, percentages and results of McNemar's test for the comparison between MRI findings in the two sides of qat chewers group.	70
5	The frequencies, percentages and results of chi-square test for comparison between Lateral pterygoid muscle (LPM) changes in qat chewers and control groups.	71
6	The frequencies, percentages and results of McNemar's test for the comparison between LPM changes in the two sides of qat chewers group.	72
7	The frequencies, percentages and results of chi-square test for the comparison between clinical findings in the three groups.	73
8	The frequencies, percentages and results of chi-square test for the comparison between ADDR in qat chewers groups (duration) and control group.	75
9	The frequencies, percentages and results of McNemar's test for the comparison between ADD of the two sides in each group of qat chewers.	76
10	The frequencies, percentages and results of chi-square test for comparison between arthritic changes in qat chewers groups (duration) and control group.	77



11	The frequencies, percentages and results of McNemar's test for the comparison between arthritic changes of the two sides in each group of qat chewers.	78
12	The frequencies, percentages and results of chi-square test for the comparison between LPM changes in qat chewers groups (duration) and control group.	79
13	The frequencies, percentages and results of McNemar's test for the comparison between LPM changes of the two sides in each group of qat chewers.	80
14	The frequencies, percentages and results of chi-square test for the comparison between joint effusion in qat chewers groups (duration) and control group.	81
15	The frequencies, percentages and results of McNemar's test for the comparison between JE of the two sides in each group of qat chewers.	82
16	The frequencies, percentages and results of chi-square test for the comparison between clinical findings in the two groups of qat chewers.	83
17	The frequencies, percentages and results of chi-square test for the comparison between disc position in qat chewers groups (way of chewing) and control group.	84
18	The frequencies, percentages and results of McNemar's test for the comparison between ADD of the two sides in each group of qat chewers.	85
19	The frequencies, percentages and results of chi-square test for the comparison between arthritic changes in qat chewers groups (way of chewing) and control group.	86
20	The frequencies, percentages and results of McNemar's test for the comparison between arthritic changes of the two sides in each group of qat chewers.	87
21	The frequencies, percentages and results of chi-square test for the comparison between LPM changes in qat chewers groups (way of chewing) and control group.	88

22	The frequencies, percentages and results of McNemar's test for the comparison between LPM changes of the two sides in each group of qat chewers.	89
23	The frequencies, percentages and results of chi-square test for the comparison between joint effusion in qat chewers groups (way of chewing) and control group.	90
24	The frequencies, percentages and results of McNemar's test for the comparison between JE of the two sides in each group of qat chewers.	91
25	The frequencies, percentages and results of chi-square test for the association between clicking and MRI findings in qat chewers group.	92
26	The frequencies, percentages and results of chi-square test for the association between presence of TMJ pain and MRI findings in qat chewers group.	93
27	The frequencies, percentages and results of chi-square test for the association between presence of LPM pain and MRI findings in qat chewers group.	94
28	The frequencies, percentages and results of chi-square test for the association between presence of joint effusion and MRI findings in qat chewers group.	95

## **LIST OF FIGURES**

<b>Number</b>	<b>Figure</b>	<b>Page</b>
1	Qat leaves.	3
2	Qat trees in Yemen.	4
3	Qat chewing on the left side.	6
4	Drawing illustrates the anatomy of the TMJ.	16
5	Drawing illustrates the anatomy of the ligaments related to TMJ.	17
6	Normal anatomy of the Jaw (muscles and their relation to jaw).	18
7	Measurement of maximum mouth opening.	47
8	Measurement of lateral movement.	47
9	Detection of sound by stethoscope.	48
10	Palpation of the lateral aspect of the TMJ.	49
11	Palpation of the posterior aspect of the TMJ.	49
12	Palpation of origin of temporalis muscles.	50
13	Palpation of insertion of temporalis muscles.	50
14	Palpation of masseter muscle.	51
15	Palpation of lateral pterygoid muscle.	51
16	Palpation of medial pterygoid muscles.	52
17	Palpation of sternocleidomastoid muscles.	52
18	Palpation of trapezius muscles.	53
19	Shows dental status of the subject.	53
20	Normal occlusion.	54

21	Cuspid protected occlusion.	55
22	Determination of overbite.	55
23	Measurement of overbite.	56
24	Measurement of overjet.	56
25	One of the subjects in a supine position during MRI examination.	57
26	Scout view shows orientation of oblique sagittal projection.	58
27	Oblique sagittal T1W in closed mouth shows disc (solid arrow) and condyle (white arrow).	59
28	Oblique sagittal T1 FL 2D in opened mouth shows disc (solid arrow) and condyle (white arrow).	59
29	Oblique sagittal T2W in opened mouth shows disc (solid arrow) and condyle (white arrow).	59
30	Oblique sagittal T1W MRI of left TMJ at closed mouth position shows normal disc position (solid arrow) with thickening at the anterior aspect of the condyle (white arrow).	63
31	Oblique sagittal T1W MRI of left TMJ at opened mouth position shows normal disc position (solid arrow) with thickening at the anterior aspect of the condyle (white arrow).	63
32	Oblique sagittal T1W MRI of right TMJ at closed mouth position shows anterior disc displacement (white arrow) and thickening at the superior aspect of the condyle (solid arrow).	63
33	Oblique sagittal T1W MRI of right TMJ at opened mouth position shows anterior disc displacement with reduction (white arrow) and osteophyte at the anterior aspect of the condyle (solid arrow).	63
34	Oblique sagittal T1W MRI of right TMJ at closed mouth position shows anterior disc displacement (white arrow).	64
35	Oblique sagittal T1W MRI of right TMJ at opened mouth position shows anterior disc displacement without reduction (white arrow).	64

36	Oblique sagittal T2W MRI of left TMJ at closed mouth position shows joint effusion at anterior recess (solid arrow) and posterior recess (white arrow).	64
37	Oblique sagittal T2W MRI of left TMJ at opened mouth position shows joint effusion at posterior recess (black arrow).	64
38	Oblique sagittal PD MRI image of left TMJ at closed mouth position showing normal LPM structures. A high-signal fat layer can be observed between the two bellies (black arrow).	65
39	Oblique sagittal PD MRI image of left TMJ at closed mouth position shows hypertrophy of superior belly (SB) of the LPM. Disc is anteriorly displaced and deformed (black arrow). Effusion can be found at bilaminar zone (white arrow).	65
40	Oblique sagittal PD MRI image of right TMJ at closed mouth position showing shows hypertrophy of inferior belly (IB) combined with atrophy of superior belly (white arrow) of the LPM.	65
41	Oblique sagittal PD MRI image of the right TMJ at closed mouth position shows atrophy and fatty replacement in superior belly of the LPM (white arrow) with normal inferior belly (IB). Disc is anteriorly displaced (black arrow).	65
42	Shows percentage of clinical findings of qat chewers group.	67
43	Shows comparison of clinical findings between chewing side and non chewing side of qat chewers group.	68
44	Shows comparison between qat chewers and control groups in disc position, arthritic changes and JE.	69
45	Shows comparison between chewing side and non chewing side of qat chewers group in disc position, arthritic changes and JE.	70
46	Shows comparison between qat chewers group and control group in LPM changes.	71

47	Shows comparison between chewing side and non chewing side of qat chewers group in LPM changes.	72
48	Shows comparison of clinical findings between the three groups.	74
49	Shows comparison of ADD between the three groups of qat chewers and control group.	75
50	Shows comparison between chewing side and non chewing side of three groups of qat chewers in ADD.	76
51	Shows comparison of arthritic changes between the three groups of qat chewers and control group.	77
52	Shows comparison between chewing side and non chewing side of three groups of qat chewers in arthritic changes.	78
53	Shows comparison of LPM changes between the three groups of qat chewers and control group.	79
54	Shows comparison between chewing side and non chewing side of three groups of qat chewers in LPM changes.	80
55	Shows comparison of JE between the three groups of qat chewers and control group.	81
56	Shows comparison between chewing side and non chewing side of three groups of qat chewers in JE.	82
57	Shows comparison of clinical findings between the two groups of qat chewers.	83
58	Shows comparison of ADD between the two groups of qat chewers and control group.	84
59	Shows comparison between chewing side and non chewing side of two groups of qat chewers in ADD.	85
60	Shows comparison of arthritic changes between the two groups of qat chewers and control group.	86
61	Shows comparison between chewing side and non chewing side of two groups of qat chewers in arthritic changes.	87

62	Shows comparison of LPM changes between the two groups of qat chewers and control group.	88
63	Shows comparison between chewing side and non chewing side of two groups of qat chewers in LPM changes.	89
64	Shows comparison of JE between the two groups of qat chewers and control group.	90
65	Shows comparison between chewing side and non chewing side of two groups of qat chewers in JE.	91
66	Shows correlation between clicking and MRI findings.	92
67	Shows correlation between TMJ pain and MRI findings.	93
68	Shows correlation between LPM pain and MRI findings.	94
69	Shows correlation between joint effusion and MRI findings.	95
70-A	Oblique sagittal T1W MRI image of the right TMJ at closed mouth position shows anterior disk displacement (solid arrow) and thickening at anterior aspect of the condyle (white arrow).	96
70-B	Oblique sagittal T1W MRI image of the right TMJ at opened mouth position shows normal disk position (ADDR) (solid arrow).	96
70-C	Oblique sagittal T1W MRI image of the left TMJ at closed mouth position shows anterior disc displacement (solid arrow).	97
70-D	Oblique sagittal T1W MRI image of the left TMJ at opened mouth position shows normal disk position (ADDR) (solid arrow).	97
70-E	Two successive oblique sagittal T1W MRI images of right TMJ at closed mouth position show feathery appearance of high signal intensity affecting SB of LPM (solid arrows) denoting fatty changes.	98
70-F	Oblique sagittal T2W MRI image of the left TMJ at closed mouth position shows joint effusion at anterior recess (white arrow) and posterior recess (solid arrows).	98