

شبكة المعلومات الجامعية







شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار في درجة حرارة من ١٥-٥٠ مئوية ورطوبة نسبية من ٢٠-٠٠% To be Kept away from Dust in Dry Cool place of 15-25- c and relative humidity 20-40%



بعض الوثائـــق الإصليــة تالفــة



بالرسالة صفحات لم ترد بالإصل

EFFECT OF SOME REPLACEMENT FLUIDS ON BLOOD COAGULATION AND PLATELET FUNCTION

Thesis submitted for partial fulfillment of M.D. degree in Anaethesiology

2 TN

BY

Ayman Zaki Ahmed Hassan M.B.B.,Ch., M. Sc., Cairo University

Supervised by

Prof. MOHAMED SALAH EL KHATEEB

Professor of Anaethesiology Faculty of Medicine- Cairo University

Prof. ABDEL RAHMAN M. FATHALLAH

Professor of Anaethesiology
Faculty of Medicine -Cairo University

Prof. HODA ALY SADEK

Professor of Clinical Pathology Faculty of Medicine -Cairo university

Ass. Prof. MOHAMED ESAWY

Assistant professor of Anesthesiology Faculty of Medicine - Cairo University

> Faculty of Medicine Cairo Univeristy 2002



Acknowledgments

I would like to express my deepest thanks to Prof. Dr. .

Mohamed Salah El Khateeb for his guidance.

I wish to express my sincere gratitude and thanks to Prof. Dr. Abdel Rahman Fathallah for his guidance.

I would like to express my deepest thanks and appreciation to Ass. Prof. Mohamed Esawy for his continuous and kind help.


I would like also to offer my sincere thanks to Prof. Dr Hoda Sadek for her kind help.

I am grateful for the valuable advice of Prof. Dr. Mohamed Abdel Latif .



List of Figures

| Fig (1) | Overview of coagulation cascade |
|----------|--|
| Fig (2) | Protein C Pathway |
| Fig (3) | The fibrinolytic pathway |
| Fig (4) | The effect of HES (6%) on bleeding time |
| Fig (5) | The effect of HES (6%) on platelet count |
| Fig (6) | The effect of HES (6%) on platelet aggregation |
| Fig (7) | The effect of HES (6%) on v W.F |
| Fig (8) | The Effect of HES (6%) on fibrinogen concentration |
| Fig (9) | The effect of Gel (4%) on bleeding time |
| Fig (10) | The effect of Gel (4%) on platelet count |
| Fig (11) | The effect of Gel (4%) on platelet aggregation |
| Fig (12) | The effect of Gel (4%) on v W. F |
| Fig (13) | The effect of Gel (4%) on fibrinogen concentration |
| Fig (14) | The effect of Dextran 70 on bleeding time |
| Fig (15) | The Effect of Dextran 70 on platelet count |
| Fig (16) | The effect of Dextran 70 on platelet aggregation |
| Fig (17) | The effect of Dextran 70 on v W. F |
| Fig (18) | The effect of Dextran 70 on fibrinogen concentration |
| Fig (19) | The effect of HS (2.7%) on bleeding time |
| Fig (20) | The effect of HS (2.7%) on platelet count |
| | |



ABSTRACT

The main reason for using colloids as volume replacement is to maintain the circulating blood volume by stabilizing plasma oncotic pressure in the preoperative period. The effects of colloids on haemostasis potentially cause an alteration ogf the haemostatic mechanisms such as interference with platelet functions. Thus the said effects of colloids may be relevant clinically.

The aim of this study is to clarify the clinical effects of colloids on blood coagulation and platelet functions.

INTRODUCTION AND AIM OF THE WORK

This work will be concerned with the study of some replacement fluids (crystalloids and colloids) and their effect on blood coagulation and platelet functions in surgical patients so that coagulopathies related to intravenous fluid replacement could be anticipated, diagnosed and treated.