THE EFFECT OF MODE OF DELIVERY AND OTHER MATERNAL AND INFANT FACTORS ON ACQUISITION OF SOME ORAL MICROFLORA IN A GROUP OF EGYPTIAN INFANTS

Thesis

Submitted in partial fulfillment of the Masters degree in

Pediatric Dentistry

Faculty of Dentistry

Ain Shams University

By

Nora Sherif Salman

B.D.Sc Ain Shams University, 2008

Faculty of Dentistry

Ain Shams University

2016

Supervisors:

Dr. Nadia Ezz El Din Metwalli

Professor of Pediatric Dentistry and Dental Public Health

Faculty of Dentistry

Ain Shams University

Dr. Mariem Osama Wassel

Lecturer of Pediatric Dentistry and Dental Public Health

Faculty of Dentistry

Ain Shams University

ACKNOWLEDGEMENT

First of all, I would like to thank **ALLAH** the Kindest and the most Merciful for assisting me and directing me to the right way.

I would like to express my deepest gratitude and admiration to Dr. Nadia Ezz El Din Metwalli, Professor in Pediatric Dentistry and Dental Public Health Department, Faculty of Dentistry, Ain Shams University, the person who taught me a lot and owe everything I achieved so far. She was my idol since I was a student in my fourth-year at dental school. She invested her time and effort with me and all her students. She was always there for me with her continuous guidance, encouragement, and positive attitude. Words are not enough to express how grateful and thankful I am.

My most sincere appreciation and gratitude to, **Dr Mariem**Osama Wassel, Lecturer in Pediatric Dentistry and Dental Public
Health Department, Faculty of Dentistry, Ain Shams University, for
her guidance, time, advice, and support during the years of my study.
It was an honor having her as my supervisor. Her insightful criticism
and valuable comments and her expertise are reflected in this work.
Her kind words of encouragement, support and valuable advice were
a push every time I was pulled backwards.

I would like to express my deepest respect to chairman, my professors, colleagues and staff members of Pediatric Dentistry and Dental Public Health department for their support and concern.

Finally endless thanks are extended for all the participants in this study. Without volunteers, this thesis work would not have been possible.

DEDICATION

I dedicate this thesis to all my family and friends for their endless love and support

My Father, for all the encouragement and motivation to fight and face all the obstacles and to keep pushing. You have been the best symbol of a great father, I Hope I always make you proud

My Mother, the symbol of love and giving, she always leads me through valleys of darkness with light and hope. Her unconditional and pure love, moral support and prayers day and night are the reasons I am who I am today

To the soul of my beloved grandmother, I wish you were there to share this success with me. You passed away in the middle of my journey but I am sure your pure soul is always there for me and guiding me from above, God bless your soul

Last but not least to my lovely brother and Amazing best friends Noha and Noreen for being there for me through ups and downs

List of contents

| Title | Page No. |
|----------------------------------|----------|
| List of Figures | I |
| List of Tables | V |
| List of Abbreviations | VII |
| Introduction | 1 |
| Review of Literature | 3 |
| Aim of the Study | 29 |
| Subjects, ,Materials and Methods | 30 |
| Results | 38 |
| Discussion | 78 |
| Summary | 92 |
| Conclusions | 95 |
| Recommendations | 96 |
| References | 97 |
| Appendix | - |
| Arabic Summary | - |

List of figures

| Figure | Title | Page No. |
|--------|--|------------|
| No. | | |
| | | |
| 1 | Data collection via questionnaire | 32 |
| | interview | |
| 2 | Salivary sample collection from mother | 35 |
| _ | - infant pair | |
| | The state of the s | |
| 3 | S.mutans and S. mitis colonies on Mitis | 37 |
| | Salivarius Agar | |
| | | |
| 4 | Bar chart representing acquisition of S. | 42 |
| | mutans in infants at 3 and 6 months in | |
| | relation to mode of child delivery | 40 |
| 5 | Bar chart representing acquisition of S. | 42 |
| | mitis in infants at 3 and 6 months in | |
| | relation to mode of child delivery | 4.4 |
| 6 | Bar chart representing counts of S. | 44 |
| | mutans in infants at 3 and 6 months in | |
| | relation to mode of child delivery | 4.4 |
| 7 | Bar chart representing counts of S. mitis | 44 |
| | in infants at 3 and 6 months in relation | |
| | to mode of child delivery | 4.5 |
| 8 | Bar chart representing acquisition of S. | 46 |
| | mutans in infants at 3 and 6 months in | |
| | relation to maternal S. mutans counts | • • |
| 9 | Bar chart representing acquisition of S. | 46 |
| | mitis in infants at 3 and 6 months in | |
| 10 | relation to maternal S. mutans counts | 40 |
| 10 | Bar chart representing acquisition of S. | 48 |
| | mutans in infants at 3 and 6 months in | |
| 11 | relation to maternal S. mitis | 40 |
| 11 | Bar chart representing acquisition of S. | 48 |
| | mitis in infants at 3 and 6 months in | |
| 10 | relation to maternal S. mitis | 5 0 |
| 12 | Bar chart representing acquisition of S. | 50 |
| | mutans in infants at 3 and 6 months in | |
| | relation to maternal DMFT | |

| 13 | Bar chart representing Acquisition of S. mitis in infants at 3 and 6 months in relation to maternal DMFT | 50 |
|----|--|----|
| 14 | Correlation between infants' S. mutans mean counts and maternal DMFT scores at three months | 52 |
| 15 | Correlation between infants' S. mutans mean counts and maternal DMFT scores at six months | 52 |
| 16 | Correlation between infants' S. mitis mean counts and maternal DMFT scores at three months | 53 |
| 17 | Correlation between infants' S. mitis mean counts and maternal DMFT scores at six months | 53 |
| 18 | Correlation between mean maternal S. mutans counts and mean S. mutans counts in infants at three months | 54 |
| 19 | Correlation between mean maternal S. mutans counts and mean S. mutans counts in infants at six months | 54 |
| 20 | Correlation between mean maternal S. mutans counts and mean S. mitis counts in infants at three months | 55 |
| 21 | Correlation between mean maternal S. mutans counts and mean S. mitis counts in infants at six months | 55 |
| 22 | Correlation between mean maternal S. mitis counts and mean S. mutans counts in infants at three months | 56 |
| 23 | Correlation between mean maternal S. mitis counts and mean S. mutans counts in infants at six months | 56 |
| 24 | Correlation between mean maternal S. mitis counts and mean S. mitis counts in infants at three months | 57 |
| 25 | Correlation between mean maternal S. mitis counts and mean S. mitis counts in infants at six months | 57 |

| <u> </u> | <u> </u> | |
|----------|---|------------|
| 26 | Bar chart representing acquisition of S. mutans in infants at 3 and 6 months in | 59 |
| 25 | relation to infants' gender | 5 0 |
| 27 | Bar chart representing acquisition of S. | 59 |
| | mitis in infants at 3 and 6 months in | |
| • | relation to infants' gender | |
| 28 | Bar chart representing counts of S. | 61 |
| | mutans in infants at 3 and 6 months in | |
| • | relation to infants' gender | |
| 29 | Bar chart representing counts of S. mitis | 61 |
| | in infants at 3 and 6 months in relation | |
| | to infants' gender | |
| 30 | Bar chart representing acquisition of S. | 63 |
| | mutans in infants at 3 and 6 months in | |
| | relation to presence of siblings | |
| 31 | Bar chart representing acquisition of S. | 63 |
| | mitis in infants at 3 and 6 months in | |
| | relation to presence of siblings | |
| 32 | Bar chart representing counts of S. | 65 |
| | mutans in infants at 3 and 6 months in | |
| | relation to presence of siblings | |
| 33 | Bar chart representing counts of S. mitis | 65 |
| | in infants at 3 and 6 months in relation | |
| | to presence of siblings | |
| 34 | Bar chart representing acquisition of S. | 67 |
| | mutans in infants at 3 and 6 months in | |
| | relation to the introduction of | |
| | complementary food and beverage | |
| 35 | Bar chart representing acquisition of S. | 67 |
| | mitis in infants at 3 and 6 months in | |
| | relation to the introduction of | |
| | complementary food and beverage | |
| 36 | Bar chart representing counts of S. | 69 |
| | mutans in infants at 3 and 6 months in | |
| | relation to the introduction of | |
| | complementary food and beverage | |
| 37 | Bar chart representing counts of S. mitis | 69 |
| | in infants at 3 and 6 months in relation | |
| | to the introduction of complementary | |
| | food and beverage | |

| 38 | Bar chart representing acquisition of S. | 71 |
|----|---|----|
| | mutans in infants at 3 and 6 months in | |
| | relation to the use of sweetened pacifier | |
| 39 | Bar chart representing acquisition of S. | 71 |
| | mitis in infants at 3 and 6 months in | |
| | relation to the use of sweetened pacifier | |
| 40 | Bar chart representing counts of S. | 73 |
| | mutans in infants at 3 and 6 months in | |
| | relation to the use of sweetened pacifier | |
| 41 | Bar chart representing counts of S. mitis | 73 |
| | in infants at 3 and 6 months in relation | |
| | to the use of sweetened pacifier | |
| 42 | Bar chart representing acquisition of S. | 75 |
| | mutans in infants at 3 and 6 months in | |
| | relation to the nursing pattern | |
| 43 | Bar chart representing acquisition of S. | 75 |
| | mitis in infants at 3 and 6 months in | |
| | relation to the nursing pattern | |
| 44 | Bar chart representing counts of S. | 77 |
| | mutans in infants at 3 and 6 months in | |
| | relation to the nursing pattern | |
| 45 | Bar chart representing counts of S. mitis | 77 |
| | in infants at 3 and 6 months in relation | |
| | to the nursing pattern | |

List of tables

| Table No. | Title | Page No. |
|-----------|---|----------|
| 1 | Mean age of mothers | 40 |
| 2 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to mode of child delivery | 41 |
| 3 | Counts of S. mutans and S. mitis in infants at 3 and 6 months in relation to mode of child delivery | 43 |
| 4 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to maternal S.mutans counts | 45 |
| 5 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to maternal S.mitis counts | 47 |
| 6 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to maternal DMFT | 49 |
| 7 | Correlation between S. mutans and S. mitis counts in infants at 3 and 6 months and maternal DMFT, materanal S. mutans and S. mitis counts | 51 |
| 8 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to infants' gender | 58 |
| 9 | Counts of S. mutans and S. mitis in infants at 3 and 6 months in relation to infants' gender | 60 |

| 10 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to presence of siblings | 62 |
|----|---|----|
| 11 | Counts of S. mutans and S. mitis in infants at 3 and 6 months in relation to presence of siblings | 64 |
| 12 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to the introduction of complementary food and beverage | 66 |
| 13 | Counts of S. mutans and S. mitis in infants at 3 and 6 months in relation to the introduction of complementary food and beverage | 68 |
| 14 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to the use of sweetened pacifier | 70 |
| 15 | Counts of S. mutans and S. mitis in infants at 3 and 6 months in relation to the use of sweetened pacifier | 72 |
| 16 | Acquisition of S. mutans and S. mitis in infants at 3 and 6 months in relation to the nursing pattern | 74 |
| 17 | Counts of S. mutans and S. mitis in infants at 3 and 6 months in relation to the nursing pattern | 76 |

List of Abbreviations

| Abb. | Full term |
|---------------|---|
| AAP | The American Academy of Pediatrics |
| AAPD | American Academy of Pediatric Dentistry |
| AWHONN | The Association of Women's Health, Obstetric and Neonatal Nurses |
| CDCP | The Center for Disease Control and Prevention |
| C-section | Cesarean-section |
| DMFT/dmft | D/d = decayed M/m = missing F/f = filled T/t = tooth |
| EAPD | European Academy of Pediatric Dentistry |
| ECC | Early childhood caries |
| HS | High significant difference |
| MDSA | Maternally Derived Streptococcus mutans Disease |
| MS | Mutans Streptococci |
| NS | No significant difference |
| S | Significant difference |
| S.mitis | Streptococcus mitis |
| S.mutans | Streptococcus mutans |
| S.oralis | Streptococcus oralis |
| S.salivarious | Streptococcus salivarius |

| S.sobrinus | Streptococcus sobrinus |
|------------|---|
| SD | Standard deviation |
| S-ECC | Severe Early Childhood Caries |
| SIgA | secretary immunoglobulinA |
| USDHHS | The United State Department for Health Education and Human Service |
| WHO | World Health Organization |

Introduction

Dental caries affects humans of all ages throughout the world and remains the major dental public health problem among school children globally. (1)

Dental caries can be defined as a bacterial disease of the calcified tissues of the teeth. It is characterized by demineralization of the inorganic substrate and destruction of the organic substrate of the tooth. ⁽²⁾ It is a major cause of tooth loss, pain and discomfort worldwide; especially amongst children. ⁽³⁾

Dental caries has a multifactorial etiology, and although it is a preventable disease, its prevention is complicated by the complex interaction of cultural, social, behavioral, nutritional, and biological risk factors that are associated with its initiation and progression. (4)

Early childhood caries (ECC) is a virulent type of dental caries. It is characterized by early onset and rapid progression. ECC is a major public health problem, being the most common chronic infectious childhood disease, which is hard to control. Although it's not life-threatening, its impact on individuals and communities is extensive, resulting in pain, impairment of function, harmful effects on the child's growth rate, body weight, and ability to thrive, thus reducing quality of life. ⁽⁵⁾

ECC is etiologically associated with bacterial colonization in deciduous dentition. Mutans streptococci (MS), especially Streptococcus mutans (S.mutans) and Streptococcus sobrinus (S.sobrinus), have been related to dental caries based on well-defined cariogenicity. (6-9)

A body of evidence suggests that children acquire S. mutans primarily from their mothers, (10-13) and possibly other sources. (14,15)

Some studies suggested that mode of child delivery whether cesarean-section (C-section) or vaginal can lead to differences in the initial foundation of microbiota in the gut and, consequently, contribute to variations in child health. Whether the mode of child delivery can alter the infant's oral microbiota and the development of ECC in the future still remains unclear. (16,17)

Therefore, the main objective of the present study was to investigate the relation between the mode of child delivery (vaginal versus C-section), and other maternal and infant factors to the colonization of oral microflora in infants.