# PHYSIOLOGICAL STUDIES ON BREAKING BUD DORMANCY OF GRAPES (VITIS VINIFERA L.)

By

### AMIN MOHAMED GALAL EL-DIN AMIN SHADDAD

B.Sc. Agric. Sc. (Horticulture), Ain Shams University, 1994M.Sc. Agric. Sc. (Pomology), Zagazig University, 1999

A thesis submitted in partial fulfillment

of

the requirements for the degree of

# DOCTOR OF PHILOSOPHY in

Agricultural Science (Pomology)

Department of Horticulture
Faculty of Agriculture
Ain Shams University

2010

### **Approval Sheet**

## PHYSIOLOGICAL STUDIES ON BREAKING BUD DORMANCY OF GRAPES (VITIS VINIFERA L.)

By

## AMIN MOHAMED GALAL EL-DIN AMIN SHADDAD

B.Sc. Agric. Sc. (Horticulture), Ain Shams University, 1994M.Sc. Agric. Sc. (Pomology), Zagazig University, 1999

This thesis for Ph. D. degree has been approved by:
Prof. Dr. Ahmed Salem Tawfeek
Prof. of Pomology, Faculty of Agriculture, Cairo, University.
Prof. Dr. Mohamed Abou-Rawash Ali Badr
Prof. Emeritus of Pomology, Faculty of Agriculture, Ain Shams
University.
Prof. Dr. Nazmy Abd El-Hamed Abd El-Ghaney
Prof. of Pomology, Faculty of Agriculture, Ain Shams University
(Supervisor).
Prof. Dr. Abd El-Azim Mohamed El- Hammady
Prof. Emeritus of Pomology, Faculty of Agriculture, Ain Shams
University (Principal Supervisor).

**Date of Examination**: 15/11/2009

## PHYSIOLOGICAL STUDIES ON BREAKING BUD DORMANCY OF GRAPES (VITIS VINIFERA L.)

#### By

### AMIN MOHAMED GALAL EL-DIN AMIN SHADDAD

B.Sc. Agric. Sc. (Horticulture), Ain Shams University, 1994M.Sc. Agric. Sc. (Pomology), Zagazig University, 1999

#### **Under the supervision of:**

#### Prof. Dr. Abd El-Azim Mohamed El- Hammady

Prof. Emeritus of Pomology, Department of Horticulture, Faculty of Agriculture, Ain Shams University (Principal Supervisor)

#### Prof. Dr. Nazmy Abd El-Hamed Abd El-Ghaney

Prof. of Pomology, Department of Horticulture, Faculty of Agriculture, Ain Shams University

#### Dr . Ahmed Abd El-Hamid Awad

Associate Prof. of Pomology, Department of Horticulture, Faculty of Agriculture, Ain Shams University

#### **ABSTRACT**

Amin Mohamed Galal El-Dine Amin Shaddad: Physiological Studies on Breaking Bud Dormancy of Grapes (*Vitis Vinifera L.*). Unpublished Ph.D., Department of Horticulture, Faculty of Agriculture, Ain Shams University, 2010.

Garlic and onion oils both at 5& 10 and 15% and GA<sub>3</sub> at 150&200and 250 ppm in comparison with dormex 5% (commercial application) were evaluated as breaking dormancy agents on "Superior" grapevines and cuttings during 2006 and 2007 seasons.

As for Superior grapevines, these materials were sprayed on three dates, December 27<sup>th</sup>, January 5<sup>th</sup> and January 13<sup>th</sup> to study the effect on bud burst%, bud fertility %, vegetative growth parameters and yield as well as cluster and berries characteristics. The vines sprayed with 15% garlic oil and 5% onion oil on January 5<sup>th</sup> had significantly higher bud burst % and bud fertility % as compared with other treatments and dates. Vegetative growth significantly increased on vines treated with garlic at 15 %, onion at 5% and dormex in the second application dates. Garlic applications at 15 & 10 % and onion at 15% on January13<sup>th</sup> markedly increased both number of clusters and yield per vine.

There was a positive relationship between fruit quality characteristics and garlic or onion application. Berry length and diameter; berry adherence; brush length; TSS; TSS/acid ratio and total sugar increased with the vines treated with garlic at 15% and onion at 5% on January 5<sup>th</sup>. Whereas, total acidity was reduced .On the other hand ,GA3 at 150 ppm was not favorable treatment where it gives similar effect as the control .

As for "Superior" cuttings, an experiment was conducted to examine the interaction between chilling exposure (100,150 and 200 hours at  $7 \pm 1^{\circ}$ c) and breaking dormancy treatments on cuttings burst % and total bud burst %. The maximum percentage of cuttings burst and

total bud burst were observed with cuttings exposed to 150 hours when treated with onion at 15% and garlic at 15%.

## **Key Words**:

Superior , grapevines , cuttings , Breaking , bud dormancy Dormex , Garlic , Onion  $\,$  ,  $GA_3$  , bud burst , bud fertility , yield ,fruit quality .

#### **ACKNOWLEDGEMENT**

I wish to offer, my sincere gratitude to **Prof. Dr. Abd El-Azim El-Hammady** professor emeritus of pomolgy, Horticulture Department, Faculty of Agriculture, Ain Shams University, for his supervision guidance, encouragement, valuable suggestions, endless advice and constructive criticism during the whole work.

Great thanks are offered to **Prof. Dr. Nazmy Abd El-Hamed Abd El-Ghaney** professor of pomolgy, Horticulture Department, Faculty of Agriculture, Ain Shams University, for his supervision, valuable suggestions, guidance and the preparation of this manuscript.

Thankful to **Dr. Ahmed Abd El-Hamid Awad**, associate prof of pomolgy, Horticulture Department, Faculty of Agriculture, Ain Shams University, for his supervision and helping in the preparation of this manuscript.

I am grateful to staff members of Department of Horticulture, Faculty of Agriculture, Ain Shams University for spiritual help and assistance.

Deepest thanks are offered to my parents, my wife and my sons for their continuous outstanding and great helping during this period.

### **CONTENTS**

	Page
LIST OF TABLES	iv
1. INTRODUCTION	1
2. REVIEW OF LITERATURE	3
3. MATERIALS AND METHODS	31
4. RESULTS AND DISCUSSION	37
4-1-Regulation and breaking of bud dormancy of Superior	37
grapevines	37
4-1-1- Budburst and Bud fertility %.	37
4- 1-1-1- Budburst%	37
4-1-1-2-Bud fertility %.	43
4-1-2- Vegetative growth parameters.	48
4-1-2-1-Average shoot length (cm)	48
4-1-2-2-Average number of leaves per shoot	50
4-1-3- Yield and cluster characteristics	53
4-1-3-1- Total yield/vine (kg).	53
4-1-3-2- Average number of clusters.	62
4-1-3-3-Cluster weight (g).	64
4-1-3-4Cluster volume (cm³)	66
4-1-3-5Cluster width (cm)	68
4-1-3-6-Cluster length (cm)	70
4-1-4-Physical characteristics of berries	72
4-1-4-1- Average berry length ( cm )	72
4-1-4-2-Weight of 100 berries (g)	74
4-1-4-3-Average berry diameter (cm)	76
4-1-4-Berry firmness (g / cm²)	76
4-1-4-5-Berry adherence strength (g/ cm²)	79
4-1-4-6-Berry brush length (cm)	81
4-1-4-7-Berry torus thickness ( cm )	83
4-1-4-8-Berry juice %	85
4-1-5-Chemical characteristics of herries	87

4-1-5-1-Total soluble solids (TSS %.)	87
4-1-5-2 Total titratable acidity	89
4-1-5-3-TSS /acid ratio	91
4-1-5-4-Total sugar %	96
4-1-5-5-Reducing sugars %	99
4-1-5-6- Non reducing sugars %	101
4-1-6- Chemical analysis.	103
4-1-6-1- Total carbohydrates.%	103
4-1-6-2-Total nitrogen %.	105
4-1-6-3-C/N ratio.	107
4-2-Breaking bud dormancy of Superior grapevines cuttings.	114
4-2-1- Cuttings burst percentage	114
4-2-2-Total bud burst percentage	122
4-2-3-Buds endogenous hormone analysis	129
5. SUMMERY AND CONCLUSION	132
6. REFRENCES	146
7 ADARIC SIIMMADV	

## LIST OF TABLES

		Page
Table(A)	Garlic analysis	32
Table(B)	Onion analysis	33
Table (1)	Effect of some breaking dormancy treatments	
	and different spraying dates on bud burst	
	percentage of "Superior" grapevines during	
	2006 and 2007 seasons.	38
Table (2)	Effect of some breaking dormancy treatments	
	and different spraying dates on bud fertility	
	percentage of "Superior" grapevines during	
	2006 and 2007 seasons.	44
Table (3)	Effect of some breaking dormancy treatments	
	and different spraying dates on average shoot	
	length (cm) of "Superior" grapevines during	
	2006 and 2007 seasons.	49
Table (4)	Effect of some breaking dormancy treatments	
	and different spraying dates on average number	
	of leaves per shoot of "Superior" grapevines	
	during 2006 and 2007 seasons.	52
Table (5)	Effect of some breaking dormancy treatments	
	and different spraying dates on vine yield in(	
	Kg) of "Superior" grapevines during 2006 and	
	2007 seasons.	55
Table (6)	Effect of some breaking dormancy treatments	
	and different spraying dates on average number	
	of clusters of "Superior" grape vines during	
	2006 and 2007 seasons.	63
Table (7)	Effect of some breaking dormancy treatments	
	and different spraying dates on cluster weight	
	(g) of "Superior" grapevines during 2006 and	
	2007 seasons.	65

Table (8)	Effect of some breaking dormancy treatments	
	and different spraying dates on cluster	
	volume(cm³)of "Superior" grapevines during	
	2006 and 2007 seasons.	67
Table (9)	Effect of some breaking dormancy treatments	
	and different spraying dates on cluster width	
	(cm) of "Superior" grapevines during 2006 and	
	2007 seasons	69
Table (10)	Effect of some breaking dormancy treatments	
	and different spraying dates on cluster length	
	(cm) of "Superior" grape vines during 2006 and	
	2007 seasons.	71
Table (11)	Effect of some breaking dormancy treatments	
	and different spraying dates on average berry	
	length (cm) of "Superior" grapevines during	
	2006 and 2007 seasons	73
Table (12)	Effect of some breaking dormancy treatments	
	and different spraying dates on weight of 100	
	berries (g) of "Superior" grapevines during 2006	
	and 2007 seasons.	75
Table (13)	Effect of some breaking dormancy treatments	
	and different spraying dates on average berry	
	diameter (cm²) of Superior grapevines during	
	2006 and 2007 seasons.	77
Table (14)	Effect of some breaking dormancy treatments	
	and different spraying dates on berry firmness	
	(g/cm²) of "Superior" grapevines during 2006	
	and 2007 seasons	78
Table (15)	Effect of some breaking dormancy treatments	
	and different spraying dates on berry adherence	
	strength (g/cm²) of "Superior" grapevines 2006	
	and 2007 seasons.	80

Table (16)	Effect of some breaking dormancy treatments	
	and different spraying dates on brush length	
	(cm) of "Superior" grapevines during 2006 and	
	2007 seasons.	82
Table (17)	Effect of some breaking dormancy treatments	
	and different spraying dates on torus thickness	
	(cm) of "Superior" grape vines during 2006 and	
	2007 seasons.	84
Table (18)	Effect of some breaking dormancy treatments	
	and different spraying dates on berry juice %	
	/100berries of "Superior" grapevines during	
	2006 and 2007 seasons.	86
Table (19)	Effect of some breaking dormancy treatments	
	and different spraying dates on total soluble	
	solids(TSS %) of "Superior" grapevines during	
	2006 and 2007 seasons.	88
Table (20)	Effect of some breaking dormancy treatments	
	and different spraying dates on total titratable	
	acidity % of "Superior" grapevines during 2006	
	and 2007 seasons.	90
Table (21)	Effect of some breaking dormancy treatments	
	and different spraying dates on TSS/acid ratio of	
	"Superior" grapevines during 2006 and 2007	
	seasons.	92
Table (22)	Effect of some breaking dormancy treatments	
	and different spraying dates on total sugar% of	
	"Superior" grapevines during 2006 and 2007	
	seasons.	98
Table (23)	Effect of some breaking dormancy treatments	
	and different spraying dates on reduced sugars%	
	of "Superior" grapevines during 2006 and 2007	100

seasons.

Table (24)	Effect of some breaking dormancy treatments	
	and different spraying dates on Non reduced	
	sugars% of "Superior" grape vines during 2006	
	and 2007 seasons	102
Table (25)	Effect of some breaking dormancy treatments	
	and different spraying dates on total	
	carbohydrates of "Superior" grapevines during	
	2006 and 2007 seasons.	104
Table (26)	Effect of some breaking dormancy treatments	
	and different spraying dates on total nitrogen of	
	"Superior" grapevines during 2006 and 2007	
	seasons.	106
Table (27)	Effect of some breaking dormancy treatments	
	and different spraying dates on C/N ratio of	
	"Superior" grapevines during 2006 and 2007	
	seasons.	108
Table (28)	Effect of some breaking dormancy treatments	
	and different chilling hours on cutting burst	
	percentage of Superior grape cuttings during	
	2006 season.	115
Table (29)	Effect of some breaking dormancy treatments	
	and different chilling hours cutting burst	
	percentage of Superior grape cuttings during	
	2007 season	117
Table (30)	Effect of some breaking dormancy treatments	
	and different chilling hours on total buds burst	
	percentage of Superior grape cuttings during	
	2006season.	125
Table (31)	Effect of some breaking dormancy treatments	
	and different chilling hours on total buds burst	
	percentage of Superior grape cuttings during	127

	2007 season	
Table (32a)	Effect of some breaking dormancy treatments	
	and different chilling hours on buds endogenous	
	hormones (GA <sub>3</sub> &IAA) of "Superior" grape	
	cuttings, during 2006 season	130
Table (32b)	Effect of some breaking dormancy treatments	
	and different chilling hours on buds endogenous	
	hormones (ABA&Zeatin R) of "Superior" grape	
	cuttings ,during 2006 season.	130
Table (32c)	Effect of some breaking dormancy treatments	
	and different chilling hours on buds endogenous	
	hormones (Zeatin&Benzyladenin) of "Superior"	
	grape cuttings, during 2006 season	131

## LIST OF FIGURES

		Page
Fig. (1)	Effect of garlic concentrations as breaking	
	dormancy treatment and different spraying dates	
	on bud burst percentage of "Superior" grapevines	
	during 2006 and 2007 seasons.	39
Fig. (2)	Effect of onion concentrations as breaking	
	dormancy treatment and different spraying dates	
	on bud burst percentage of "Superior" grapevines	
	during 2006 and 2007 seasons	40
Fig. (3)	Effect of.GA <sub>3</sub> concentrations as breaking	
	dormancy treatment and different spraying dates	
	on bud burst percentage of "Superior" grapevines	
	during 2006 and 2007 seasons.	41
Fig. (4)	Effect of garlic concentrations as breaking	
	dormancy treatment and different spraying dates	
	on bud fertility percentage of "Superior"	
	grapevines during 2006 and 2007 seasons.	45
Fig. (5)	Effect of onion concentrations as breaking	
	dormancy treatment and different spraying dates	
	on bud fertility percentage of "Superior"	
	grapevines during 2006 and 2007 seasons	46
Fig. (6)	Effect of GA <sub>3</sub> concentrations as breaking	
	dormancy treatment and different spraying dates	
	on bud fertility percentage of "Superior"	
	grapevines during 2006 and 2007 seasons.	47
Fig. (7)	Effect of garlic concentrations as breaking	
	dormancy treatment and different spraying dates	
	on total yield of "Superior" grapevines during	
	2006 and 2007 seasons	56

	Effect of onion concentrations as breaking	
	dormancy treatment and different spraying dates	
Fig. (8)	on total yield (Kg) of "Superior" grapevine during	
	2006 and 2007 seasons.	57
Fig. (9)	Effect of GA <sub>3</sub> concentrations as breaking	
	dormancy treatment and different spraying dates	
	on total yield(Kg) of "Superior" grapevines during	
	2006 and 2007 seasons	58
Fig. (10)	Effect of garlic concentrations as breaking	
	dormancy treatment and different spraying dates	
	on TSS/acid ratio of "Superior" grapevines during	
	2006 and 2007 seasons.	93
Fig. (11)	Effect of onion concentrations as breaking	,,,
	dormancy treatment and different spraying dates	
	on TSS/acid ratio of "Superior" grapevines during	
	2006 and 2007 seasons.	94
Fig. (12)	Effect of GA <sub>3</sub> concentrations as breaking	74
	dormancy treatment and different spraying dates	
	on TSS/acid ratio of "Superior" grapevines during	
	2006 and 2007 seasons.	95
Fig. (13)	Effect of garlic concentrations as breaking	)3
<i>6</i> · ( - )	dormancy treatment and different spraying dates	
	on C/N ratio of "Superior" grapevines during 2006 and 2007 seasons.	109
Fig. (14)		109
1 181 (1 1)	6	
	dormancy treatment and different spraying dates	
	on C/N ratio of "Superior" grapevines during 2006	110
Fig. (15)	and 2007 seasons.	110
115. (13)	Effect of GA <sub>3</sub> concentrations as breaking	
	dormancy treatment and different spraying dates	
	on C/N ratio of "Superior" grapevines during	111
	2006 and 2007 seasons	111