

**BIOREMEDIATION OF OLIVE MILL
WASTEWATER FOR USING IN
BIOFERTILIZATION**

By

ROKIA YEHIA ALI AWADALLAH

B.Sc. Agric. Sci. (Agric. Microbiology), Fac. of Agric., Ain Shams Univ., 2010

**A Thesis Submitted in Partial Fulfillment
Of
The Requirements for the Degree of**

**MASTER OF SCIENCE
in
Agricultural Sciences
(Agriculture Microbiology)**

**Department of Agriculture Microbiology
Faculty of Agriculture
Ain Shams University**

2017

Approval Sheet

BIOREMEDIATION OF OLIVE MILL WASTEWATER FOR USING IN BIOFERTILIZATION

By

ROKIA YEHIA ALI AWADALLAH

B.Sc. Agric. Sci. (Agric. Microbiology), Fac. of Agric., Ain Shams Univ., 2010

This thesis for M. Sc. degree has been approved by:

Dr. Mervat Ahmed Taher Emara
Researcher prof. of Microbiology, Desert Research Center

Dr. Sohair Ahmed Ebrahim Naser
Prof. of Microbiology, Faculty of Agriculture, Ain Shams University

Dr. Ahmed Abdel-Wahab Abdel-Hafez
Prof. of Microbiology, Faculty of Agriculture, Ain Shams University

Dr. Elshahat Mohamed Ramadan
Prof. Emeritus of Microbiology, Faculty of Agriculture, Ain Shams
University

Date of Examination: 8 / 11/ 2017

BIOREMEDIATION OF OLIVE MILL WASTEWATER FOR USING IN BIOFERTILIZATION

By

ROKIA YEHIA ALI AWADALLAH

B.Sc. Agric. Sci. (Agric. Microbiology), Fac. of Agric., Ain Shams Univ., 2010

Under the supervision of:

Dr. Elshahat Mohamed Ramadan

Prof. Emeritus of Microbiology, Department of Microbiology,
Faculty of Agriculture, Ain Shams University, (Principal Supervisor)

Dr. Ahmed Abdelwahab Abdelhafez

Prof. of Microbiology, Department of Microbiology, Faculty of
Agriculture, Ain Shams University

Dr. Azza Ahmed Mohamed

Head Research of Microbiology, Department of Microbiology
Research, Institute of Soil, Water and Environment, ARC

ACKNOWLEDGEMENT

All Praise and thanks to ALLAH, the most merciful for directing me to the right way and provides me all I have.

I would like to express my sincere gratitude to **Prof. Dr. ElShahat Mohamed Ramadan**, Prof. of Agric. Microbiology, Faculty of Agriculture, Ain Shams University for his kind supervision, guidance, valuable help, continuous support and encouragement, precious advices, and useful discussion during the study.

I deeply thank to **Dr. Ahmed Abdelwahab Abdelhafez**, Prof. of Agric. Microbiology, Faculty of Agriculture, Ain Shams University for his motivation, enthusiasm, useful discussion during the study and continuous help during the study and during writing up the manuscript.

Special thanks to **Prof. Dr. Azza Ahmed Mohamed**, Prof. of Microbiology research Dept., Institute of Soil, Water and Environment, ARC for her help, advice, her kind supervision. I would gratefully acknowledge her valuable support. valuable technical advice, introducing all lab facilities and continuous help during the study.

I owe my sincere thanks to **Prof. Dr. Hassan Ismail**, Prof. of Microbiology research Dept, Institute of Soil, Water and Environment, ARC, for his genius guidance, valuable help and his encouragement during this study.

CONTENTS

	Page
LIST OF TABLES	i
LIST OF FIGURES	ii
LIST OF ABBREVIATIONS	iii
1. INTRODUCTION	1
2. REVIEW OF LITERATURE	4
2.1. Olive oil extraction	4
2.1.1. Pressure extraction system	5
2.1.2. Centrifugation system	7
2.1.2.1. Three-phase system	7
2.1.2.2. Two phase system	8
2.2. General characteristics of olive mill wastewater	8
2.3. Remediation of Olive Mill Wastewater	8
2.3.1. Physical treatments	8
2.3.1.1. Dilution	9
2.3.1.2. Sedimentation	9
2.3.1.3. Filtration	9
2.3.1.4. Flotation	9
2.3.1.5 Centrifugation	9
2.3.1.6 Membrane Processes	9
2.3.2. Physico-thermal processes	10
2.3.2.1. Evaporation/distillation	10
2.3.2.2. Dring	10
2.3.2.3 Pyrolysis	10
2.3.2.4 lagooning	10
2.3.3. Physico-Chemical Treatments	11
2.3.3.1. precipitation	11
2.3.3.2. coagulation and flocculation	11
2.3.3.3. Chemical oxidation processes	11
2.3.4. Biological treatments	11
2.4. 2.4. Effect of OMWW on the Environment	13

II

	Page
2.4.1. Effect on soil	14
2.4.2. Effect on water	14
2.4.3. Effect on Atmosphere	15
2.5. Utilization of OMWW	15
2.5.1. Use OMWW as medium for growth algae and edible mushroom	15
2.5.2. OMWW as Fertilizer	16
2.5.3. . Use OMWW in composting	19
2.5.4. Recovery of Antioxidants from OMWW	19
2.5.5. Production of Biogas and Biopoymers	19
2.5.6. Recovery of Enzymes	20
2.5.6. Use as Pesticide	20
3. MATERIALS AND METHODS	21
3.1. MATERIALS	21
3.1.1. Sampling	21
3.1.2. Selected Microorganisms	21
3.1.3. Microbial media	21
3.1.4. Plant seeds	25
3.2. Experiments	25
3.2.1. Bioremediation of OMWW by Fungi	25
3.2.1.1. Isolation of fungi from OMWW	25
3.2.1.2. Screening for isolates efficient in phenolic compounds degradation and COD reduction	26
3.2.1.3. Comparison of the most potent isolate with <i>Pleurotus columbinus</i> as a reference	26
3.2.2. phycoremediation of OMWW	26
3.2.3. Utilization of treated and untreated OMWW in irrigation	26
3.2.3.1. Germination test	26
3.2.3.2. Pot experiment	27
3.3. Methods of determination and analysis	28
3.3.1. Determination of total phenol	28

	Page
3.3.2. Determination of total carbohydrates	29
3.3.3 Determination of COD	29
3.3.4. Determination of pH and EC	29
3.3.5. Determination of microelements	29
3.3.6. Determination of total nitrogen	30
3.3.7. Determination of chlorophyll pigment	30
3.3.8. Decolorization Assay	30
3.3.9. Determination of dehydrogenase activity	30
3.3.10. Determination of Nitrogenase activity	30
3.3.11. Determination of CO ₂ evolution	30
3.3.12. Determination of Indole Acetic Acid (IAA)	31
3.3.13. Determination of Gibberellins	31
3.4. Statistical analysis	31
4. RESULTS AND DISCUSSION	32
4.1. Fungal treatment of OMWW	32
4.1.1. Chemical Analysis of the collected OMWW samples	32
4.1.2. Isolation and Screening the efficiency of fungal isolates.	32
4.1.3. Comparison of the most potent fungal isolate with <i>Pleurotus columbinus</i> as a reference	34
4.1.3.1. Phenolic compounds degradation by fungal isolate 5 and <i>Pleurotus columbinus</i>	34
4.1.3.2. COD reduction by fungal isolate 5 and <i>Pleurotus columbinus</i>	38
4.1.3.3. Decolorization of OMWW by fungal isolate 5 and <i>Pleurotus columbinus</i>	41
4.1.3.4. Growth of isolate 5 and <i>Pleurotus columbinus</i> in OMWW	44
4.1.3.5. pH and EC change after treatment by isolate 5 and <i>Pleurotus columbinus</i>	46
4.1.3.6. Determination of total carbohydrates in fungal treated OMWW	46

	Page
4.1.3.7.Determination of Indol Acetic Acid and Gibberellins contents in OMWW treated with <i>P. columbinus</i> .	48
4.2.Phycoremediation of OMWW by algae	51
4.2.1. Chemichal Analysis of OMWW	51
4.2.2. Degradation of phenolic compounds by algae	51
4.2.3 Reduction of COD value in OMWW by algal	55
4.2.4. Effect of algal strains and species on carbohydrate content of OMWW.	58
4.2.5. Decolorization of OMWW treated with algal strains and species	61
4.2.6. Growth of algal strain in OMWW	64
4.2.7. pH value of OMWW after treated with algae	66
4.2.8 Indol acetic acid (IAA) and gibberellins (GA)of OMWW after treated with algae treated with algae	68
4.3. Evaluation of treated OMWW as fertilizer	73
4.3.1. Plant height, shoot and roots weight	74
4.3.2. NPK of Shoot and root determination	78
4.3.4. Dehydrogenase and Nitrogenase activity in rhizosphere of pea	81
4.3.5. Determination of Chlorophyll, carotene and CO ₂ in peas plants	83
5. SUMMARY	85
6. REFERENCES	92
ARABIC SUMMARY	

LIST OF TABLES

Table No.		Page
3.1	Physical and chemical properties of the experimental soil	28
4.1	Chemical composition of OMWW from Agricultural Research Center	32
4.2	Effect of fungal isolates on phenol content and COD of 25% diluted OMWW after 2 weeks incubation at 28°C.	33
4.3	Degradation of phenolic compounds in OMWW at different concentrations by fungal isolate 5, compared to <i>P. columbinus</i> as reference.	36
4.4	Determination of COD (g/L), in OMWW at different concentration by fungal isolate 5 compared to <i>P. columbinus</i> as reference.	39
4.5	Decolorization of OMWW by isolate 5 compared to <i>P. columbinus</i> as reference	42
4.6	Growth of fungal isolate 5 and <i>P. columbinus</i> (dry weight g/L) in OMWW at different concentrations, incubated for 4 weeks at 28°C	45
4.7	Determination of carbohydrate content (g/L) released from <i>P. columbinus</i> in OMWW	47
4.8	Determination of Indole Acetic Acid and gibberellins in 10% and 20% OMWW, treated with <i>P. columbinus</i>	49
4.9	Chemical composition of OMWW collected from Wadi Food industry	51
4.10	Degradation of phenolic compounds in 20% OMWW by algae incubated at 30°C/3000 lux for 30days.	53

Table No.		Page
4.11	Degradation of phenolic compounds in 10% OMWW by algae incubated at 30°C/3000 lux for 30days	54
4.12	Reduction of COD in 20% OMWW treated by algae incubated at 30°C/3000 lux for 30days	56
4.13	Reduction of COD value in 10% OMWW by algae incubated at 30°C/3000 lux for 30days.	57
4.14	Carbohydrate content (g/L) in 10 and 20% OMWW treated with algal strains and species incubated at 30°C/3000 lux for 30days	59
4.15	Decolorization of OMWW of 10 and 20% concentrations treated with algal strains and species.	62
4.16	Growth of algal strains and species in OMWW at 20% and 10% concentrations	65
4.17	Change in pH of 20 and 10% OMWW treated with algae.	67
4.18	Indole acetic acid and gibberellins content at (20%) OMWW concentrations treated with <i>Spirulina platensis</i> and <i>Wollea</i> sp.	70
4.19	Growth parameters of Peas plants after 45days of plantation, irrigated with 10% OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp. (diluted to 50% with tap water).	75
4.20	Shoot and root NPK of Peas plants after 45 days of plantation, irrigated with 10% OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp then diluted to 50% with tap water	79

Table No.		Page
4.21	Dehydrogenase and Nitrogenase activities in peas plants rhizosphere after 45days of plantation, irrigated with OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp and diluted to 50%.	82
4.22	Determination of Chlorophyll, carotene and CO ₂ in peas plants after 45days of plantation, irrigated with OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp and diluted to 50%.	84

LIST OF FIGURES

Fig. No.		Page
2.1	Diagram of the three olive oil extraction methods.	5
2.2	Pressure extraction system	6
2.3	Olive oil extraction by centrifugation (continuous) system	7
4.1	Effect of fungal isolates on phenolic compounds degradation and COD reduction of 25% diluted OMWW after 2 weeks incubation at 28°C	33
4.2a	Degradation of phenolic compounds in OMWW at different concentrations by fungal isolate 5	37
4.2b	Degradation of phenolic compounds in OMWW at different concentrations by <i>P. columbinus</i>	37
4.3a	Determination of COD in OMWW at different concentrations as treated with isolate 5	40
4.3b	Determination of COD in OMWW at different concentration as treated with <i>P. columbinus</i>	40
4.4	Decolorization of 10% OMWW by <i>P. columbinus</i> after 4 weeks	43
4.5a	Decolorization (O.D at 395nm) of OMWW at 40, 30, 20 and 10% by isolate 5	43
4.5b	Decolorization (O.D at 395nm) of OMWW at 40, 30, 20 and 10% by <i>P. columbinus</i>	44
4.6a	Growth of isolate 5 (dry weight/L) in OMWW at 40, 30, 20 and 10% incubated for 4 weeks at 28°C	45
4.6b	Growth of <i>P. columbinus</i> (dry weight,g/L) in OMWW at 40, 30, 20 and 10% incubated for 4 weeks at 28°C	46
4.7	Determination of total carbohydrate in 10, 20, 30 and 40% OMWW treated with <i>p. columbinus</i> for 4 weeks.	48

Fig. No.		Page
4.8a	Determination (IAA) in 10 and 20% OMWW treated with <i>P. columbinus</i> .	50
4.8b	Determination (GA) in 10 and 20% OMWW treated with <i>P. columbinus</i> .	50
4.9	Degradation of phenolic compounds in 20% OMWW by algae incubated at 30°C/3000 lux for 30days.	53
4.10	Degradation of phenolic compounds in 10% OMWW by algae incubated at 30°C/3000 lux for 30days.	54
4.11	Reduction of COD in 20% OMWW by algae incubated at 30°C/3000 lux for 30 days	57
4.12	Reduction of COD value in 10% OMWW by algae incubated at 30°C/3000 lux for 30 days	58
4.13a	Carbohydrate content in 20% OMWW by algal strains and species incubated at 30°C/3000 lux for 30 days.	60
4.13b	Carbohydrate content in 10% OMWW by algal strains and species incubated at 30°C/3000 lux for 30 days.	60
4.14a	Decolorization of 10% OMWW treated with algae	63
4.14b	Decolorization of 20% OMWW treated with algae	63
4.15a	Growth of algal strains and species (dry weight g/L) in 20% OMWW	65
4.15b	Growth of algal strains and species (dry weight g/L) in 10% OMWW	66
4.16 a	Change in pH of 20% OMWW treated with algae.	67
4.16 b	Change in pH of 10% OMWW treated with algae.	68
4.17 a	IAA content at 20 and 10% OMWW treated with <i>Spirulina platensis</i>	71

Fig. No.		Page
4.17 b	IAA content at 20 and 10% OMWW treated with <i>Wollea</i> sp	71
4.18 a	GA content at 20 and 10% OMWW treated with <i>Spirulina platensis</i>	72
4.18 b	GA content at 20 and 10% OMWW treated with <i>Wollea</i> sp	72
4.19 a	Plant height of Peas plants after 45days of plantation, irrigated with 10% diluted OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp. then diluted to 50% with tap water	76
4.19 b	Shoot and root dry weight of Peas plants after 45days of plantation, irrigated with 10% OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp. then diluted to 50% with tap water	77
4.20 a	Shoot NPK of Peas plants after 45 days of plantation, irrigated with OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp then diluted to 50% with tap water	80
4.20 b	Root NPK of Peas plants after 45 days of plantation, irrigated with 10% OMWW treated with <i>P. columbinus</i> , <i>Spirulina platensis</i> and <i>Wollea</i> sp then diluted to 50% with tap water.	81