

**RESISTANCE TO REINFECTION IN MICE INFECTED  
WITH *SCHISTOSOMA MANSONI* AFTER USING  
COMBINED ANTIHELMINTHIC AND ANTIFIBROTIC  
AGENTS**

By

**EMAN ALI MOHAMED MOSTAFA**

B. Science, Ain Shams University, 1980,  
Master in Environmental Sci., Ain Shams University, 1996

**A Thesis Submitted In Partial Fulfillment  
For The Ph.D. Degree**

In

Environmental Sciences  
Department of Biological and Natural Sciences

Under Supervision of

**Prof. Dr. Ameen Abd El-Baqy Ashour**

Prof. of Parasitology,  
Vice Dean of Community and Environmental Development-  
Faculty of Science, Ain Shams University


**Prof. Dr. Mohamed Abd El-Kader El-Khafif**

Prof. of Biochemistry,  
Dean, Institute of Environmental Studies and Research-  
Ain Shams University

**Prof. Dr. Soad Ismail Hassan**

Prof. of Parasitology,  
Head of parasitology Departement,  
Theodor Bilharz Research Institute

**2005**



"وَقُلْ رَبِّ زِدْنِي عِلْمًا"

سورة طه الآية: 114

الْعَظِيمُ

## Acknowledgments

*I am very thankful to **Prof. Dr. Ameen A. Ashour**, Professor of Parasitology, Vice Dean for Community and Environmental Development, Faculty of Science, Ain Shams University, for his outstanding devotion, great help and guidance that have been of utmost help in performing this work.*

*My cordial thanks are due to **Prof. Dr. Mohamed A. El-Khafif**, Professor of Biochemistry, Dean, Institute of Environmental Studies and Research, Ain Shams University, for his encouragement, instructive guidance, valuable assistance and meticulous revision of all details of this thesis.*

*I wish to express my tender thanks and deepest gratitude towards **Prof. Dr. Soad I. Hassan**, Professor of Parasitology, and Head of Parasitology Department, Theodor Bilharz Research Institute, for her endless help and support, she guided me throughout the preparation of this thesis, and patiently supervised every bit of the details.*

*I am really grateful to **Ass. Prof. Dr. Nevein N. Giurgius**, Assistant Professor of Parasitology, Theodor Bilharz Research Institute, for her brilliant scientific ideas, tremendous concern and unfailing support which helped to bring this work to light.*

*I feel deeply thankful for **Ass. Prof. Dr. Neimat Mousa** Assistant Professor of Parasitology, Theodor Bilharz Research Institute, for her cooperative supervision, generous help and sincere guidance during this study.*

*I wish to express my thanks to **Ass. Prof. Dr. Mona Noseir**, Assistant Professor of Pathology, Theodor Bilharz Research Institute, for her help in performing the histopathological study in this work.*

*I am very thankful to Ass. Prof. Dr. Amira H.M. Hassan, Assistant Professor of Electron Microscope, Theodor Bilharz Research Institute, for her kind help in preparing this work.*

*Words would not be sufficient to express my deepest, heartfelt thanks towards my colleagues in the Parasitology Department and to all my colleagues in Schistosome Biological Supply Program unit (SBSP) at Theodor Bilharz Research Institute, who provided all the animals and S. mansoni cercariae used in this study. Their care and encouragement gave me the power to complete this work.*

*Eman A.M. Mostafa*  
*2005*

## LIST OF ABBREVIATIONS

Abbreviation	Definition
<b>AdDP</b>	Adamantylmide dipeptide
<b>ALT</b>	Alanine aminotransferase
<b>AST</b>	Aspartate amino-transferase
<b>B/6</b>	A strain of mice
<b>BAPN</b>	$\beta$ -aminopropionitrile
<b>C3H</b>	A strain of mice
<b>C<sub>57</sub>BL/6</b>	A strain of mice
<b>CBA</b>	A strain of mice
<b>CDI</b>	A strain of mice
<b>Chall.</b>	Challenged
<b>ESP</b>	Eosinophil stimulating promotor
<b>IFN</b>	Interferon
<b>Ig</b>	Immunoglobulin
<b>IL-5</b>	Interleukin-5
<b>Inf</b>	Infected
<b>IWB</b>	Immature worm burden
<b>LO</b>	Lysyl oxidase enzyme
<b>M</b>	Macrophages
<b>PZQ</b>	Praziquantel
<b>R0-11-0761</b>	[3, (3,5 dinitro-2-thienyl) thiozolidine]
<b>SEA</b>	Soluble egg antigen

Abbreviation	Definition
--------------	------------

## *List of Abbreviations*

---

<b>TEM</b>	Transmission Electron Microscopic
<b>Th</b>	T helper cells
<b>TWB</b>	Total worm burden
<b>U/L</b>	Unit per liter

# CONTENTS

	<i>Page</i>
<b>Acknowledgement</b>	<b>i</b>
<b>List of Abbreviations</b>	<b>iii</b>
<b>Contents</b>	<b>v</b>
<b>List of Tables</b>	<b>x</b>
<b>List of Figures</b>	<b>xiii</b>
<b>Abstract</b>	<b>xxiv</b>
<b>Introduction</b>	<b>1-3</b>
<b>Aim of the Work</b>	<b>4</b>
<b>Review of the Literature</b>	<b>5-35</b>
- Schistosome species	5
- Susceptibility and resistance	6
- Resistance to reinfection in schistosomiasis <i>mansoni</i>	7
- Chemotherapy of schistosomes	13
-Praziquantel	
- Resistance to reinfection with <i>Schistosoma mansoni</i> after specific chemotherapy	19
-β-aminopropionitrile	
- Effect of schistosomiasis on liver functions	25
- Pathology of schistosomiasis	27
- Schistosomal liver granulomatous reaction	28
- Effect of granulomas on the resistance	32
- Transmission electron microscopy	33
- Electron microscopic picture of hepatic schistosomiasis	34
<b>MATERIAL AND METHODS</b>	<b>36-45</b>
I. Experimental animals	
II. Cercarial count	
III. Route of infection and challenge	
IV. Test drugs	
V. Performed investigations	

1- Parasitological investing	
a. Hepatic perfusion	
b. Mesenteric perfusion	
c. Egg count in liver & intestine tissue	
d. Oogram studies	
2- Count of schistosome eggs in stools	
3- Enzyme assay investigation	
4- Histopathological investigations	
A. Preservation and processing	
B. Sectioning	
C. Histopathological study	
D. Measurement of granuloma size	
5- Electron microscopic preparation	
6- Statistics	
VI. Experimental design	46-54
- Experiment (1)	
- Experiment (2)	
Parameters of the study:	
1) Parasitological parameters	
2) Biochemical studies	
3) Histopathological parameters	
4) Electron microscopic study	
5) Statistical methods	
RESULTS	57-173
Effect of treatment with $\beta$ -aminopropionitril on worm burden and distribution in <i>Schistosoma mansoni</i> -infected mice	57
1. Infected untreated control	
2. Infected-challenged control	
a- Total mature worm burden	
b- Mature worm distribution	
3. Infected-PZQ treated-challenged group	
4. Infected-BAPN treated-challenged group	
5. Infected-PZQ-BAPN treated-challenged group	
	<i>Page</i>
The percentage of resistance to reinfection in <i>S. mansoni</i>	



infected mice challenged with 240 cercariae and treated with praziquantel and $\beta$ -aminopropionitril	
a. Total mature worm burden	
b. Total immature worms	
c. Percentage of resistance	
d. Percentage of reduction in worm burden	
Oogram pattern in <i>S. mansoni</i> infected mice challenged with 240 cercariae and treated with praziquantel and B-aminopropionitril	66
a. The total immature stage	
b. The total mature stage	
c. The dead ova	
Tissue egg load in <i>S. mansoni</i> infected mice challenged with 240 cercariae and treated with praziquantel and $\beta$ -aminopropionitril	69
a. The hepatic egg load	
b. The intestinal egg load	
c. The percentage of reduction	
Effect of treatment with $\beta$ -aminopropionitril on egg excretion in stool in <i>S. mansoni</i> infected mice	73
Serum enzymes in <i>S. mansoni</i> infected mice challenged with 240 cercariae and treated with praziquantel and $\beta$ -aminopropionitril	76
1. Serum alanine aminotransferase (ALT)	
2. Serum asparate aminotransferase (AST)	
Effect of treatment with $\beta$ -aminopropionitril on collagen content in <i>S. mansoni</i> infected mice	79
Hepatic granuloma diameter in <i>S. mansoni</i> infected mice challenged with 240 cercariae and treated with praziquantel and $\beta$ -aminopropionitril	81
	<i>Page</i>
Liver pathology in <i>S. mansoni</i> infected mice challenged with 240 cercariae and treated with praziquantel and $\beta$ -	84

**aminopropionitril**

<b>Electron microscopic examination in <i>S. mansoni</i>-infected mice challenged with 240 cercariae and treated with praziquantil and <math>\beta</math>-aminopropionitrile</b>	<b>100</b>
--	------------

1. Animals sacrificed 10 weeks after infection
2. Animals sacrificed 12 weeks after treatment with a single therapeutic dose of praziquantel (PZQ)
3. Animals sacrificed 12 weeks after treatment with a single therapeutic dose of B-aminopropionitrile (BAPN)
4. Animals treated with praziquantel then B-aminopropionitrile therapeutic dose and sacrificed after 12 weeks

<b>Effect of chemotherapy and/or BAPN (<math>\beta</math>-aminopropionitril) on resistance to reinfection in <i>Schistosoma mansoni</i>-infected mice</b>	<b>115</b>
---	------------

**The mean number of worm burden and worm distribution**

<b>Evaluation of resistance to reinfection</b>	<b>120</b>
--	------------

<b>Percent egg developmental stages (oogram)</b>	<b>124</b>
--	------------

<b>The mean tissue egg load</b>	<b>127</b>
---------------------------------	------------

<b>Effect of treatment with BAPN on egg excretion in stool in <i>Schistosoma mansoni</i>-infected mice</b>	<b>131</b>
--	------------

<b>Effect of treatment with BAPN on the liver function as assessed by the levels of some serum enzymes in <i>Schistosoma mansoni</i>-infected mice</b>	<b>134</b>
--	------------

1. Serum alanine aminotransferase (ALT)
2. Serum aspartate aminotransferase (AST)

<b>The mean value of the hepatic tissue collagen</b>	<b>138</b>
--	------------

*Page*

<b>Hepatic granuloma diameter</b>	<b>140</b>
-----------------------------------	------------

<b>Histological study of liver</b>	<b>144</b>
------------------------------------	------------

<b>Electron microscopic examination in <i>S. mansoni</i>-infected mice challenged with 120 cercariae and treated with praziquantil and <math>\beta</math>-aminopropionitrile</b>	<b>160</b>
<b>DISCUSSION</b>	<b>174-202</b>
<b>SUMMARY</b>	<b>203-209</b>
<b>REFERENCES</b>	<b>210-255</b>
<b>ARABIC SUMMARY</b>	<b>1-5</b>

## LIST OF TABLES

No.	Table	Page
1-	Mean worm burden and distribution in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	60
2-	Percentage of resistance and worm reduction in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ (500mg/kg for 2 days) 2 weeks before challenged and/or treated with BAPN (daily) for 14 days.	64
3-	The percentage egg developmental stages in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	67
4-	Mean number of ova/gm tissue in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	71
5-	Mean number of egg excretion in infected mice primarily with (100) <i>Schistosoma mansoni</i> cercariae and treated with PZQ and/or BAPN.	74
6-	Serum enzymes in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ (500mg/kg for 2 days) 2 weeks before challenged and/or treated with BAPN (daily) for 14 days before challenge infection.	77

No.	Table	Page
-----	-------	------

7-	Mean hepatic collagen content in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ (500mg/kg for 2 days) 2 weeks before challenge and/or treated with BAPN (daily) for 14 days before challenge infection.	80
8-	Mean hepatic granuloma diameter in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ (500mg/kg for 2 days) before challenge and/or treated with BAPN versus infected untreated challenged animals..	82
9-	Mean worm burden and distribution in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	118
10-	Percentage resistance to reinfection in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ (500mg/kg for 2 days) 2 weeks before challenged and/or treated with BAPN (daily) for 14 days.	122
11-	Percentage egg developmental stages in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ (500 mg/kg for two days) before challenged and/or treated with BAPN (5mg in 0.5 ml saline) for 14 successive days.	125
12-	Mean number of ova/gm tissue in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	129

- |     |   |     |
|-----|---|-----|
| 13- | Mean number of egg excretion in infected mice primarily with (40) <i>Schistosoma mansoni</i> cercariae and treated with PZQ and/or BAPN.  | 132 |
| 14- | Serum alanine aminotransferase (ALT, units/L), aspartate aminotransferase (AST, units/L) in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ (500mg/kg for 2 days) 2 weeks before challenged and/or treated with BAPN (daily) for 14 days before challenge infection. | 136 |
| 15- | Mean hepatic collagen content in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ (500mg/kg for 2 days) 2 weeks before challenged and/or treated with BAPN versus control group and chronic infection group.  | 139 |
| 16- | Mean hepatic granuloma diameter in <i>Schistosoma mansoni</i> infected mice (40 cercariae) and treated with PZQ before challenged and/or treated with BAPN versus infected untreated challenged animals.  | 142 |

## LIST OF FIGURES

No.	Figure	Page
<b>Material and Methods:</b>		
<b>Diagrams:-</b>		
1-	Representation of the work studying the resistance to reinfection in murine schistosomiasis <i>mansoni</i> .	47
2-	Experimental design of group (A) infection of mice with 100 <i>Schistosoma mansoni</i> cercariae and challenged with 240 cercariae.	48
3-	Representation of groups of the experimental animal model for study the effect of PZQ and/or BAPN on schistosomiasis.	50
4-	Experimental design of group (B) infection of mice with 100 <i>Schistosoma mansoni</i> cercariae and challenge infection.	51
5-	Representation of groups of experimental model for study of the effect of administration of $\beta$ -aminopropionitrile with or without Praziquantel on resistance to reinfection.	53
<b>Results:</b>		
<b>Figures:-</b>		
1-	The mean worm burden in <i>Schistosoma mansoni</i> -infected mice (100 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	61
2-	The mean number of worms and the percentage of resistance to reinfection in <i>Schistosoma mansoni</i> infected mice (100 cercariae) and treated with PZQ before challenged and/or treated with BAPN (daily) for 14 days.	65
No.	Figure	Page