



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
على هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

Contrast Sensitivity before and after Laser in Situ Keratomileusis

Thesis

Submitted for the partial fulfillment of medical doctorate degree in
ophthalmology

by

Amre Abd-Elbari Hamed Hassan
M.B., B. Ch. , M .Sc.
Supervised by

Prof. Dr. Shaker Ahmed Khedr
Professor of ophthalmology
Ain Shams University

Prof. Dr. Mervat Salah
Professor of ophthalmology
Ain Shams University


Prof. Dr. Sherif Mahmoud Zohni
Professor of ophthalmology
Research Institute of Ophthalmology

Prof. Dr. Bahaa Abd-Allah
Assistant Professor of ophthalmology
Ain Shams University

Ain Shams University
Cairo
2001

B

9221




بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قَالُوا سُبْحَانَكَ لَا عِلْمَ لَنَا إِلَّا مَا عَلَّمْتَنَا
إِنَّكَ أَنْتَ الْعَلِيمُ الْحَكِيمُ

صدق الله العظيم

قرآن كريم

الآية ٣٢ من سورة البقرة



توقيع لجنة المناقشة علي رسالة الدكتوراة المقدمة من

الطبيب

عمرو عبد الباري

وتتكون لجنة المناقشة من السادة :-

(برناتس)

أ.د. بشر قناوي

(ميسر)

أ.د. حسين شاكر


(نازك)

أ.د. شاكر خضر

~~~~~

~~~~~

~~~~~



| <i>Figure</i>                                                                                                                                                                                                                                                                                           | <i>Page</i> |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Fig (34): Treatment screen of Nidek EC-5000 excimer laser system showing planned spherical correction, cylindrical correction, optical zone, transitional zone, operation time, ablation depth, and scan number for treatment of - 2D spherical error, and -1D cylindrical error with axis 15 °.</b> |             |
| <b>Fig (35): Treatment screen of Nidek EC-5000 excimer laser system showing Planned spherical correction, cylindrical correction,optical zone, Transitional zone, operation time, ablation depth, and scan number for treatment of- 10D spherical error.</b>                                            | <b>94</b>   |
| <b>Fig (36): Treatment screen of Nidek EC-5000 excimer laser system showing Planned spherical correction, cylindrical correction, optical zone, Transitional zone, operation time, ablation depth, and scan number for treatment of- 6.25D spherical error.</b>                                         | <b>95</b>   |
| <b>Fig (37): Bilateral post LASIK corneal topography showing right central islands</b>                                                                                                                                                                                                                  | <b>96</b>   |
| <b>Fig (38): post LASIK bilateral corneal topography showing eccentric ablation of the right eye</b>                                                                                                                                                                                                    | <b>97</b>   |
| <b>Fig (39): Differential topographic map showing pre and post LASIK corneal changes</b>                                                                                                                                                                                                                | <b>98</b>   |
| <b>Fig (40): Postoperative corneal topography showing adjustable post LASIK map with flattening of the central cornea.</b>                                                                                                                                                                              | <b>99</b>   |

| <i>Figure</i>                                                                                              | <i>Page</i> |
|------------------------------------------------------------------------------------------------------------|-------------|
| <b>Fig (41): Bilateral post LASIK corneal topography showing well centered homogeneous ablation zones.</b> |             |
| <b>Fig (42): Number of eyes included in the study.</b>                                                     | <b>103</b>  |
| <b>Fig (43): Number of females and males in the study.</b>                                                 | <b>104</b>  |
| <b>Fig.(44):Mean pre and postoperative visual acuities of group A.</b>                                     | <b>111</b>  |
| <b>Fig (45): Mean contrast sensitivity values for group (A) 2 weeks postoperatively.</b>                   | <b>112</b>  |
| <b>Fig (46): Mean contrast sensitivity values for group (A) 4 weeks postoperatively.</b>                   | <b>113</b>  |
| <b>Fig (47): Mean contrast sensitivity values for group (A) 12 weeks postoperatively.</b>                  | <b>114</b>  |
| <b>Fig.(48):Mean pre and postoperative visual acuities of group B.</b>                                     | <b>122</b>  |
| <b>Fig (49):Mean contrast sensitivity values for group (B) 2 weeks postoperatively.</b>                    | <b>123</b>  |
| <b>Fig (50):Mean contrast sensitivity values for group (B) 4 weeks postoperatively.</b>                    | <b>124</b>  |

## *L*ist of figures

---

| <i>Figure</i>                                                                                                | <i>Page</i> |
|--------------------------------------------------------------------------------------------------------------|-------------|
| Fig (11) :Epithelial ingrowth.                                                                               |             |
| Fig (12): Post LASIK corneal topography showing Homogeneous uniform ablation pattern.                        | 62          |
| Fig (13): Post LASIK corneal topography showingirregular ablation pattern                                    | 63          |
| Fig(14) : Post LASIK corneal topography showing toric ablation pattern                                       | 64          |
| Fig(15): Post LASIK corneal topography showing central island pattern.                                       | 65          |
| Fig (16): Cambridge low contrast Gratings book                                                               | 75          |
| Fig (17): Cambridge low contrast Gratings with grated and blank pages hanged to the wall                     | 76          |
| Fig (18): Contrast sensitivity testing chart                                                                 | 77          |
| Fig (19): Preoperative examination unit in refractive clinic (Research institute of ophthalmology).          | 78          |
| Fig (20): TMS 1 system corneal topography                                                                    | 79          |
| Fig (21): Absolute map in the TMS 1 corneal topography screen Showing sever steepening of the central cornea | 80          |
| Fig (22): Rabinowitz, klyce indices for topographic keratoconus detection                                    | 81          |

| <i>Figure</i>                                                                                                                      | <i>Page</i> |
|------------------------------------------------------------------------------------------------------------------------------------|-------------|
| Fig (23): preoperative Rabinowitz , Klyce indices for Screening of topographic keratoconus in one of the studied eyes              |             |
| Fig (24): ultrasonic pachymetry (Sonoscan model 4000AP)                                                                            | 83          |
| Fig (25): Nidek EC-5000 Excimer laser system                                                                                       | 84          |
| Fig (26): Scanning slit system of Nidek EC-5000 ( Lavery, 1996 )                                                                   | 85          |
| Fig (27): Alignment system of Nidek EC-5000 ( Lavery, 1996 )                                                                       | 86          |
| Fig (28): Hansatome microkeratome.                                                                                                 | 87          |
| Fig (29): Hansatome microkeratome parts showing the motor, suction rings, and blades for 160 $\mu$ m and 180 $\mu$ m lamellar cuts | 88          |
| Fig (30): suction ring (8.5 mm) diameter and blade for 160 $\mu$ m lamellar cut in position.                                       | 89          |
| Fig (31): Hansatome microkeratome blade on its handle for fixation.                                                                | 90          |
| Fig (32): Suction ring of the Hansatome microkeratome applied to the eye.                                                          | 91          |
| Fig (33): The microkeratome head moving along its pass.                                                                            | 92          |