

شبكة المعلومات الجامعية







شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



شبكة المعلومات الجامعية

جامعة عين شمس

التوثيق الالكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها على هذه الأفلام قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيدا عن الغبار في درجة حرارة من ١٥-٥٠ مئوية ورطوبة نسبية من ٢٠-٠٠%. To be Kept away from Dust in Dry Cool place of 15-25- c and relative humidity 20-40%



بعض الوثائـــق الإصليــة تالفــة



بالرسالة صفحات لم ترد بالإصل

LAPAROSCOPIC MANAGEMENT OF GASTRO-OESOPHAGEAL REFLUX DISEASE PROCEDURES



Essay

Submitted for Partial Fulfilement of Master Degree in General Surgery

By

Ahmed Mohamed El Shaer

M.B.B.Ch of Medicine, Ain Shams University

Supervised By

Prof. Dr. Nabil Shedid

Professor of General Surgery
Faculty of Medicine
Benha University

Prof. Dr. Essam Sadek

Professor of General Surgery Faculty of Medicine Benha University

Dr. Mostafa Saleh

Lecturer of General Surgery Faculty of Medicine Benha University

Faculty of Medicine
Benha University
2006



Acknowledgement

First I thank **GOD** the most helpful and the most merciful.

No.

S. San

SCOOLS.

X X X

i i

SSESS

XXXXX

MANAGE

RANANA

NAMAN

MANANA

WWWW

WWW.

I would like to express my inner feeling of thankfulness and sincerity for **Prof. Dr Nabil Shedid**, for his help and advises.

Also I hope to success in thanking and showing may appreciation for **Prof. Dr Essam Sadek**, for spending much time in supervision assistance patience an guidance in the review.

Finally I thank **Dr Mostafa Saleh** for his cooperation in this work.

List of Contents

Title	Page No.
Introduction and aim of the work	1-3
Anatomical consideration	4-25
Physiological considerations	26-31
Pathology of GERD	32-40
Diagnosis of GERD	41-62
Investigations	63-89
Treatment	90-169
Summary	170-172
References	173-194

List of Figures

Fig. No.	age No.
Fig. (1): Topography and constriction of the oesophagus	5
Fig. (2): Relation of the abdominal part of the oesophagus	
Fig. (3): Arterial supply of the esophagus	
Fig. (4): Venous supply of the esophagus	
Fig. (5): Lymphatic drainage of the oesophagus	
Fig. (6): Anatomy of the diaphragm (Thoracic part)	
Fig. (7): Anatomy of the diaphragm (Abdominal part)	18
Fig. (8): Gastroesophageal junction	24
Fig. (9): Evaluating symptoms in oesophageal disorders	
Fig. (10): Normal manometry	
Fig. (11): Position of the reflux operations	
Fig. (12): Transabdominal open nissen fundoplication	
Fig. (13): A transthoracic mobilization of the stomach and esophagus	
Fig. (14): Transthoracic open Nissen fundoplication	
Fig. (15): Floppy Nissen fundoplication	
Fig. (16): Belsy Mark IV procedure	
Fig. (17): Collis gastroplasty	
Fig. (18): Hill posterior gastropexy	
Fig. (19): Toupet fundoplication	
Fig. (20): Operating room layout for laparoscopic antireflux	
Fig. (21): Port locations for laparoscopic antireflux surgery	
Fig. (22): Laparoscopic Nissen fundoplication	
Fig. (23): Technique of laparoscopic antireflux surgery	

List of Tables

Tab. No. Pag	ge No.
Table (1): Results of laparoscopic antireflux surgery LARS	150
Table (2): Reasons for failure of the primary antireflux repair	156



Introduction and Aim of the Work

INTRODUCTION

Gastro-oesophageal reflux disease (GERD) is the medical term for chronic heartburn. Many people have heartburn at times it's caused by irritation from stomach acid that backs up into the oesophagus. The oesophagus is the tube that carries food from your mouth to your stomach sometimes stomach acid backs all the way up to the throat or into the mouth. This acid has a sour taste (Fennerty, 2000).

The simplest approach is to define the disease by its symptoms. However, symptoms thought to be indicative of GERD, such as heart burn, acid regurgitation, are very common in general population and many individuals consider them to be normal and do not seek medical advice (Kahrijas, 2001).

GERD can usually be diagnosed based on the clinical presentation alone. In some patients, however, the diagnosis may require endoscopy and, ambulatory pH monitoring. Management includes lifestyle modifications and pharmacologic therapy, refractory disease requires surgery the therapeutic goals are to control symptoms, heal esophagitis and maintain remission so that morbidity is decreased and quality of life is improved (Velanbvich, 2001).

Patients who did not respond well to lifestyle changes and drug therapy, especially those with grade three and four oesophagitis, and the young age group, should be treated surgically. Reduction of the hiatus hernia if present, repair of the oesophageal hiatus and fundoplication is a surgical procedure that has stood the test of time. However, the traditional open procedure requires a large incision and along, painful recovery (Spechler, 2001).

Laparoscopy has dramatically changed operations for gastroesophageal reflux disease (GERD), making them increasingly common in the last decade. As experience with laparoscopic antireflux procedures ahs increased, operating times, conversion rates, post perioperative complications, and hospital length of stay have all decreased. Through long term outcomes data are not yet available, it is clear that the laparoscopic approach to this procedure is equally effective compared with the open approach. In addition, the benefit to the patient in terms of postoperative recovery is indisputable (Flipi and Lehman, 2001).

AIM OF THE WORK

The aim of this study is review the literature and highlight update in diagnosis and treatment of gastro-oesophageal reflux disease (GERD regarding medical and surgical treatment with special stress on laparoscopic techniques. Evaluation of different ways for this technique will be discussed and correlated.