



شبكة المعلومات الجامعية

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ





شبكة المعلومات الجامعية



شبكة المعلومات الجامعية

التوثيق الالكتروني والميكرو فيلم

جامعة عين شمس

التوثيق الالكتروني والميكرو فيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأفلام قد اعدت دون أية تغيرات



يجب أن

تحفظ هذه الأفلام بعيداً عن الغبار

في درجة حرارة من 15 – 20 مئوية ورطوبة نسبية من 20-40 %

To be kept away from dust in dry cool place of
15 – 25c and relative humidity 20-40 %



شبكة المعلومات الجامعية



بعض الوثائق الأصلية تالفة



شبكة المعلومات الجامعية



بالرسالة صفحات
لم ترد بالأصل

High-Frequency Ultrasonography as a Tool in the Diagnosis of Functional and Organic Colopathies

Thesis

Submitted for Partial Fulfillment of
Master Degree in Tropical Medicine

By

Naglaa Adel Shawky El Hosary

M.B., B.Ch.

Faculty of Medicine
Ain Shams University

Supervised by

Dr. Hassan Salah El-Din Hamdy

Assist. Prof. of Tropical Medicine
Faculty of Medicine
Ain Shams University

Prof. Dr. Ahmed Abass El Katib

Prof. of Tropical Medicine
Faculty of Medicine
Ain Shams University

Prof. Dr. Waheed Hussein Tantawy

Prof. of Radiodiagnosis
Faculty of Medicine
Ain Shams University

**Faculty of Medicine
Ain Shams University**

2001

B 7702

List of Contents

	Page
Introduction and aim of the work	1
Review of Literature:	3
• Chapter 1: Anatomical, Histological and Physiological Consideration of the Colon	
• Chapter 2: Colonic Diseases	20
Subjects and Methods	97
Results	103
Discussion	128
Conclusion	135
Recommendations	136
Summary	137
References	139
Arabic Summary	175



List of Tables

Table in Review

- Differentiation between diverticular disease of the colon and colonic carcinoma by means of radiological features

Tables in Results

- Table (1): Socio-demographic data of the studied groups
- Table (2): Symptomatology of the studied groups
- Table (3): General symptoms in the studied groups
- Table (4): General examination of the studied groups
- Table (5): Abdominal examination of the studied groups
- Table (6): Results of laboratory investigation in the studied groups

I- Stool analysis in the studied groups

II- CBC and ESR in the studied groups

- Table (7): Colonoscopic findings in the studied groups
- Table (8): Sonographic findings in the studied groups
- Table (9): Results of sonography in comparison to results of colonoscopy in the studied groups
- Table (10): Sensitivity and specificity of ultrasonography in different studied groups
- Table (11): Descriptive data of results of colonoscopy and sonography in total cases



List of Figures

- **Figure (1):** High-frequency bowel sonography shows normal wall thickness of the colon in a healthy control.
- **Figure (2):** Bowel sonography shows thickened hypoechoic oedematous rectal wall in a patient with ulcerative colitis.
- **Figure (3):** High-frequency bowel sonography shows marked thickening of mucosa and submucosa of rectosigmoid colon in a patient with ulcerative colitis.
- **Figure (4):** High-frequency bowel sonography shows thickened descending colon in a patient with ulcerative colitis.
- **Figure (5):** High-frequency bowel sonography shows thickened hypoechoic descending colon in a patient with ulcerative colitis reaching 1cm with multiple surrounding paracolic lymph nodes the largest is about 6 mm in diameter.
- **Figure (6):** High-frequency bowel sonography shows thickened and oedematous transverse colon in a patient with ulcerative colitis.
- **Figure (7):** High-frequency bowel sonography shows thickened ascending colon and caecum and dilated, distended ileum in a patient with Crohn's disease together with large pericolic lymph node.
- **Figure (8):** High-frequency bowel sonography shows irregular soft tissue mass at the rectosigmoid colon with small pericolic lymph node.



List of Figures

- **Figure (9):** High-frequency bowel sonography shows large circumferential hypoechoic soft tissue mass at the sigmoid colon with small pericolic lymph node.
- **Figure (10):** High-frequency bowel sonography shows irregular soft tissue mass infiltrating the splenic flexure with gas seen inside it (splenic flexure mass).
- **Figure (11):** High-frequency bowel sonography shows irregular hypoechoic soft tissue mass infiltrating the hepatic flexure with gas seen in its middle (hepatic flexure mass).
- **Figure (12):** High-frequency bowel sonography shows left paracolic abscess in a patient with diverticular disease of the colon

