

بسم الله الرحمن الرحيم





شبكة المعلومات الجامعية

التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغيرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بعض الوثائق الأصلية تالفة





بالرسالة صفحات لم ترد بالأصل



Role of Salivary Progesterone In Prediction of Preterm Labour

Thesis

Submitted for Partial Fulfillment of
Master Degree in Obstetrics and Gynecology



By

Walaa Adel El-Sayed El-Bahat

M.B., B.Ch

Cairo University (2003)

Resident of obstetrics and Gynecology

Shubra General Hospital

Supervisors

Dr. Magd El Din M. Mohammed

Professor of Obstetrics and Gynecology

Faculty of medicine

Ain Shams University

Dr. Mohammed Hussein Mostafa


Lecturer of Obstetrics and Gynecology

Faculty of medicine

Ain Shams University

Faculty of medicine
Ain Shams University

2011





Acknowledgment

*First of all, I wish to express my special thanks to **Dr. SABRY SAYED HASSAN**, Professor of Obstetrics and Gynecology, Ain Shams University for his early supervision and guidance.*

*I would like to express my sincere appreciation and my deep gratitude to **Dr. MAGD EL DIN M. MOHAMMED**, Professor of Obstetrics and Gynecology, Ain Shams University for his faithful supervision and guidance.*

*I would like to express my great thanks to **Dr. MOHAMMED HUSSEIN MOSTAFA**, Lecturer of Obstetrics and Gynecology, Ain Shams University for the tremendous effort he has done in the meticulous revision of this work.*

Lastly, I am indebted to my family, my husband and thanks to every body supported me.

Walaa Adel El Sayed

Contents

Subjects	Page
• List of Abbreviations.....	I
• List of Tables.....	III
• List of Figures	IV
• Introduction	1
• Aim of the work.....	5
• Review of literature:	
• Chapter (1): Epidemiology of preterm labour	6
• Chapter (2): Prediction of preterm labour.....	26
• Chapter (3): Progesterone.....	49
• Patients and Methods	66
• Results	73
• Discussion	83
• Summary	88
• Conclusion	90
• Recommendations	91
• References	92
• Arabic Summary	

List Abbreviations

- **AF:** amniotic fluid
- **ALP:** alkaline phosphatase
- **AUC:** area under the curve
- **BMI:** body mass index
- **CI:** confidence interval
- **CL:** cervical length
- **CRH:** corticotrophin-releasing hormone
- **DHEA-S:** dehydroepiandrosterone sulphate
- **ELISA:** enzyme-linked immunosorbant assay
- **FFN:** fetal fibronectin
- **HCG:** human chorionic gonadotropin
- **HMD:** hyaline membrane disease
- **HUAM:** home uterine activity monitoring
- **IL-1:** interleukine-1
- **IL-6:** interleukine-6
- **ILGFBP-1:** insulin-like growth factor binding protein-1
- **LDL:** low density lipoprotein
- **MFMU:** maternal-fetal medicine units
- **MIAC:** microbial invasion of the amniotic cavity
- **NBV:** negative predictive value

 List of Abbreviations

- **NICHHD:** National Institute of Child Health and Human Development
- **PIGFBP-1:** phosphorylated insulin-like growth factor binding protein1
- **PPROM:** preterm premature rupture of membranes
- **PPV:** positive predictive value
- **PROM:** premature rupture of membranes
- **RDS:** respiratory distress syndrome
- **ROC:** receiver operating characteristic
- **ROP:** retinopathy of prematurity
- **SPSS:** statistical package for the social sciences
- **TNF:** tumor necrosis factor
- **TVU:** trans vaginal ultrasound

List of Tables

Table No.	Title	Page No.
1a	Progesterone pharmacological types and protocols	64
1	Demographic characteristics of the study groups.	73
2	Salivary progesterone levels in preterm group	74
3	Salivary progesterone levels in full term group	75
4	Comparison between preterm and full term groups according to the mean salivary progesterone levels	76
5	Comparison between preterm and full term groups according to the mean maternal age, salivary progesterone level two weeks before delivery and cervical length (26 weeks).	78
6	Comparison between preterm and full term groups according to salivary progesterone levels.	81
7	Correlation between cervical length (at 26 weeks) and salivary progesterone level two weeks before delivery.	82

