

بِسْمِ اللّٰهِ الرَّحْمٰنِ الرَّحِیْمِ





شبكة المعلومات الجامعية التوثيق الالكتروني والميكروفيلم



جامعة عين شمس

التوثيق الإلكتروني والميكروفيلم

قسم

نقسم بالله العظيم أن المادة التي تم توثيقها وتسجيلها
علي هذه الأقراص المدمجة قد أعدت دون أية تغييرات



يجب أن

تحفظ هذه الأقراص المدمجة بعيدا عن الغبار





بالرسالة صفحات لم ترد بالأصل





بعض الوثائق الأصلية تالفة



C 52

The role of Hyaluronidase in periocular anaesthesia

Thesis submitted in partial fulfillment of master degree in ophthalmology

By

Mohamed Abd El-Hameed Abd EL-Fattah Kabil
M.B.,B.Ch.



Supervised By

65703

Prof. Dr. Shaker Ahmed Khedr

Professor of Ophthalmology
Ain Shams University

Prof. Dr. Samir Mohamed Kasim

Assist. Professor of Ophthalmology
Ain Shams University


Faculty of Medicine
Ain Shams University

2000

Cairo

617.78
M.A.





بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ


Acknowledgment

I wish to express my deep gratitude and my profound indebtedness to prof. Dr. Shaker Khedr, Professor of ophthalmology, Ain Shams University, for his kind supervision, unfailing advice and constructive guidance. It was through his continuous help and moral support that this work was completed.

I am also deeply grateful to Prof. Dr. Sam~~er~~er Kas~~im~~im, ^{xx}
Ass. Professor of Ophthalmology, Ain Shams University, who devoted much of his time, effort and experience in the supervision of this work. His precious advice contributed a lot in the preparation of this study.

I would also like to express my sincere gratitude and thanks to Assist. Lecturer Dr. Hatem Ayman for his generous contribution to the arrangement of many of the figures included in this study.

Special thanks are also due to my family for their unlimited support and continuous help to assist this work to be completed successfully.



Contents (protocol)

Introduction.....	1
Aim of the work.....	3
Review of literature	
• Anatomy.....	4
• Pharmacology.....	41
• Complications of ophthalmic regional anaesthesia.....	60
Subjects and methods.....	65
Results.....	71
Discussion.....	87
Conclusion and recommendations.....	94
Summary.....	95
References.....	97
Arabic summary	

List of Figures

Chapter 3

- Fig. 3-1 An artist drawing of the sensory areas of the eye and orbit on the right and the sensory nerves on the left.....4
- Fig. 3-2 Patient with high brow, lax lids, wide palpebral fissure, and forward set globe.....7
- Fig. 3-3 Patient with short, tight lids, and deep set globe.....8
- Fig. 3-4 Schematic drawing for the geometric relationship of the walls of the orbit9
- Fig. 3-5 A computer generated drawing through the inferior orbital rim to show the orbital path length at the plane of the inferotemporal limbus... 11
- Fig. 3-6 Atkinson up and in position of the eye..... 12
- Fig. 3-7 Frontal view of figure 6..... 13
- Fig. 3-8 Incision of the conjunctiva of the superior fornix reveals connective tissue strands between frontal periorbit and Tenon's capsule.....15
- Fig. 3-9 Connective tissue septa between the lacrimal bone, periorbit and Tenon's capsule.....16
- Fig. 3-10 Needles inserted for medial block at 4:00, for a superotemporal block at 10:45 and for inferotemporal block at 7:00 and 8:00.....18
- Fig. 3-11 Coronal section parallel to the orbital axis through the hind surface of the globe.....19
- Fig. 3-12 Coronal section perpendicular to the optic nerve 5 mm. Posterior to the hindsurface of the globe.....20
- Fig. 3-13 Coronal section parallel to the orbital axis 4.6 mm. Anterior to the hindsurface of the globe.....22
- Fig. 3-14 Transverse mid-axial section through the optic nerve and the medial and lateral rectus muscles.....24
- Fig. 3-15 Malignant and non-malignant Graves' ophthalmopathy.....26
- Fig. 3-16 Frontal section of the right orbit posterior to the globe demonstrating intermuscular septum, which ensheathes the rectus muscles.....27
- Fig. 3-17 Histological section shows no intermuscular septum ensheathing the rectus muscles and connecting them to each other28
- Fig. 3-18 Drawing of the connective tissue system of the medial rectus muscle... 29
- Fig. 3-19 An artist's depiction of the septa in the posterior orbit viewed from behind looking into the mid-orbit.....31
- Fig. 3-20 Drawing of the connective tissue system of the extraocular muscles ... 33

Chapter 4

- Fig. 4-1 Structural formula of Procaine..... 42
- Fig. 4-2 Structural formula of Bupivacaine.....43
- Fig. 4-3 Typical anatomy of peripheral nerves.....46
- Fig. 4-4 Diagram showing a myelinated and non-myelinated nerves.....47

- Fig. 4-5 Myelinated nerves with thin axons are more readily blocked than large thick axons.....48
- Fig. 4-6 Local anaesthetic injected extraneurally to avoid nerve trauma.....49
- Fig. 4-7 The anaesthetic front has traversed the nerve's mantle region but not the core region.....49
- Fig. 4-8 The anaesthetic front although weakened reaches the core.....50
- Fig. 4-9 Ultimately, the intra- and extraneural concentrations equalise.....50
- Fig. 4-10 With the extraneural drug reservoir depleted, the diffusion gradient reverses.....51
- Fig. 4-11 Mantle bundles lose local anaesthetic faster than core bundles.....51

Chapter 6

- Fig. 6-1 The first periocular injection site.
- Fig. 6-2 The second periocular injection site at the medial canthal angle nasal to the caruncle.
- Fig. 6-3 The upper eyelid becomes ptotic at the conclusion of the second injection.
- Fig. 6-4 Facial block was given using Obrien's technique whenever needed.