

**Impact of Extracoronary Calcification and Pericardial Fat Volume
on Extent of Coronary Artery Disease in Patients with Type 2
Diabetes Mellitus**

Thesis submitted for partial fulfillment of MD degree in cardiology

By

Samir Mohamed Hassan Sorour, M.Sc.

Assistant Researcher of Cardiology

National Research Center

Under supervision of

Azza Abdel Moniem Farrag, MD

Professor of Cardiology

Cairo University

Hala Mohamed Reslan, M.D

Professor of Internal Medicine

National Research Center

Sameh Wadie Bakhoun, M.D

Assistant Professor of Cardiology

Cairo University

Mohamad Ali Salem, MD

Lecturer of Radiology

Cairo University

Cairo University

2015

ACKNOWLEDGEMENT

First and foremost, I thank **God** for helping and guiding me in accomplishing this work.

I would like to express my gratitude and deepest thanks to ***Prof. Dr. Azza Farrag***, Prof. of Cardiology, Cairo University, for her supervision, valuable remarks, advices and continuous encouragement throughout this work.

I am greatly indebted to ***Prof. Dr. Hala Mohamed Raslan***, Prof. of Medicine, National Research Center, for her meticulous supervision, superfluous advices and sincere help.

I must extend my warmest gratitude to ***Dr. Sameh Bakhoun***, Ass. Professor of Cardiology, Cairo University, for his continuous support, meticulous supervision, great valuable remarks, encouragement and assistance until this work was fulfilled.

I would like to express my deepest thanks to ***Dr. Mohamad Ali Salem***, Lecturer of Radiology, Cairo University, for his sincere help, valuable remarks and advices throughout this work.

My gratitude goes also for ***Dr. Ashraf Elsherbiny***, Ass. Professor of Medicine, National Research Center, for his suggestions, advices and encouragement.

Finally, I would like to express my special thanks and gratitude to my family for their continuous support and encouragement.

Abstract

Background: Type 2 diabetes mellitus (T2DM) is now considered a coronary artery disease (CAD) equivalent. We hypothesized that coronary artery calcium score (CACS), extracoronary calcium score (ECCS) and pericardial fat volume (PFV) may represent useful additional markers for risk assessment in diabetic patients.

Methods: This prospective study was conducted on 1032 patients. They included 402 patients with T2DM. Using multidetector computed tomography (MDCT), we assessed PFV, CACS and ECCS. The severity of CAD was quantified using the Gensini Score. Correlation between CACS, ECCS and PFV and severity of CAD was studied.

Results: Diabetic patients with ECCS > 0 in comparison with diabetic patients with ECCS=0 had higher CACS (163 ± 239 vs. 72 ± 176 , $p < 0.000$), higher Gensini score (22.3 ± 31.6 vs. 9.7 ± 16.3 , $p < 0.000$) and higher prevalence of patients with Gensini score > 0 ($p < 0.000$). There was a significant linear correlation between ECCS and Gensini score ($r=0.328$, $p < 0.001$). Comparing diabetic patients with $PFV \geq 100 \text{ cm}^3$ with diabetic patients with $PFV < 100$ revealed that prevalence of patients with Gensini score > 0 was not significantly higher in patients with $PFV \geq 100 \text{ cm}^3$ ($p = 0.205$). Gensini score also was not significantly different between the 2 groups (13.9 ± 25.1 vs. 12.8 ± 21.3 , $p = 0.370$). Logistic regression analysis was applied to detect possible association between age, male gender, hypertension, diabetes mellitus duration, CACS, ECCS, PFV and Gensini score > 0. Male gender (OR = 3.021, $p = 0.007$, 95% CI: 1.352-6.750), and CACS (OR=1.021, $P < 0.000$, 95% CI: 1.011-1.031) were the only independent predictors of Gensini score > 0.

Conclusion: In T2DM, neither ECCS > 0 nor $PFV \geq 100 \text{ cm}^3$ were predictors of Gensini score > 0. Male gender and CACS were independent predictors of Gensini > 0.

List of contents

	Page
• Abbreviations.....	I
• List of tables	IV
• List of figures	V
• Introduction.....	1
• Aim of work.....	3
• Literature review	
Chapter 1: Coronary artery calcification.....	4
Chapter 2: Extracoronary calcification	16
Chapter 3: Pericardial fat volume	37
• Methods.....	55
• Results.....	62
• Discussion.....	75
• Summary.....	87
• References.....	90
• Appendix 1: Master table.....	121
• Appendix 2: coding for master table.....	139
• Arabic summary	

Abbreviations

AAC	Ascending aorta calcium
AC	Aortic calcification
ACE	Angiotensin converting enzyme
AD	Aortic distensibility
ADVANCE	Action in Diabetes and Vascular Disease: Preterax and Diamicron Modified Release Controlled Evaluation
ARC	Aortic root calcification
ATP	Adult treatment panel
AU	Agatston units
AVC	Aortic valve calcium
AVF	Abdominal visceral fat
AVS	Aortic valve sclerosis
AWT	Aortic wall thickness
BMI	Body mass index
BMP	Bone morphogenic proteins
CABG	Coronary artery bypass grafting
CAC	Coronary artery calcification
CACS	Coronary artery calcium score
CAD	Coronary artery disease
CAG	Conventional coronary angiography
CAP	Calcified atherosclerotic plaque
CHD	Coronary heart disease
CKD	Chronic kidney disease
CMR	Cardiovascular magnetic resonance
CRP	C reactive protein
CT	Computed tomography
CTCA	Computed tomographic coronary angiography
CVD	Cardiovascular disease
CVE	Cardiovascular events
DAC	Descending aorta calcium
DHS	Diabetes Heart Study
DM	Diabetes mellitus

EAT	Epicardial adipose tissue
EBCT	Electron beam computed tomography
ECCS	Extracoronary calcium score
ESRD	End-stage renal disease
EVASCAN	Medical Evaluation of Scanner in Coronary Syndrome
FFAs	Free fatty acids
FPG	Fasting plasma glucose
FRS	Framingham Risk Score
HDL	High density lipoprotein
HOMA-IR	Homeostasis model assessment of insulin resistance
HU	Hounsfield units
HVC	Heart valve calcium
ICAM	Intercellular adhesion molecule
IDF	International diabetes federation
IFG	Impaired fasting glucose
IL-6	Interleukin-6
LDCT	Low-dose ungated chest CT
LDL	Low density lipoprotein
MAC	Mitral annulus calcification
MACE	Major adverse cardiovascular events
MCAP	Mixed coronary atherosclerotic plaque
MCP-1	Monocyte chemoattractant protein-1
MDCT	Multidetector computed tomography
MESA	Multiethnic Study of Atherosclerosis
MetS	Metabolic syndrome
MI	Myocardial infarction
MPS	Myocardial perfusion scintigraphy
NDR	National Diabetes Register
NCAP	Noncalcified coronary atherosclerotic plaque
NFG	Normal fasting glucose
PAI-1	Plasminogen activator inhibitor-1
PDHS	The Penn Diabetes Heart Study
PET	Positron emission tomography
PF	Pericardial fat
PFT	Pericardial fat thickness
PFV	Pericardial fat volume

PREDICT	Prospective Evaluation of Diabetic Ischaemic Disease by Computed Tomography
ROS	Reactive oxygen species
RV	Right ventricle
SEAS	Simvastatin and Ezetimibe in Aortic Stenosis
SIS	Segment involvement score
SPECT	Single-photon emission computed tomography
sPLA2-IIA	Secretory type II phospholipase A2
SS	Shear stress
SSS	Segment stenosis score
T2DM	Type 2 diabetes mellitus
TAC	Thoracic aortic calcification
TFV	Intrathoracic fat volume
TNF- α	Tumor necrosis factor alpha
UKPDS	United Kingdom Prospective Diabetes Study
VADT	Veterans Affairs Diabetes Trial
VAT	Visceral abdominal fat
VC	Valve calcification
VPF	Pericardial fat volume indexed by body surface area
VSMCs	Vascular smooth muscle cells
WC	Waist circumference
WHR	Waist to Hip ratio
WHtR	Waist to Height ratio

List of tables

	Page
Table 1 Risk factors for coronary calcification.....	6
Table 2 Associations between novel cardiovascular risk factors and CACS.....	7
Table 3 Multiple regression analysis identifying the independent risk factors associated with mean PF thickness.....	46
Table 4 Demographic and clinical characteristics of diabetic and non diabetic patients.....	63
Table 5 Multidetector computed tomography data of the studied population.....	64
Table 6 Demographic and clinical characteristics within the diabetic group.....	65
Table 7 MDCT findings comparison between diabetic patients with ECCS > 0 and diabetic patients with ECCS = 0.....	66
Table 8 Correlation between DM duration and different parameters.....	66
Table 9 Comparison of demographic data and clinical characteristics of diabetic patients with $PFV \geq 100 \text{ cm}^3$ and patients with $PFV < 100 \text{ cm}^3$	70
Table 10 Comparison of MDCT findings in diabetic patients with $PFV \geq 100$ and patients with $PFV < 100$	71
Table 11 linear correlation of CACS, ECCS and PFV with Gensini score.....	72
Table 12 Regression analysis for variables assessment in diabetic patients to predict Gensini score >0.....	74

List of figures

	Page
Figure 1 The prevalence and extent of calcified plaque in coronary arteries in asymptomatic patients < 40 years of age according to absence and presence of diabetes.....	11
Figure 2 The relationship between the extent of coronary calcification and the prevalence/severity of myocardial perfusion abnormality	13
Figure 3 Cardiovascular mortality with higher CAC scores in the DHS cohort compared with CAC score <10 in a full model.....	14
Figure 4 Hazard ratios for effect of glycemc treatment (intensive vs. standard) in multivariable-adjusted models.....	15
Figure 5 The odds ratio for the presence of mitral annular calcification based on CAC Agatston score category.....	18
Figure 6 Combined end point (events), coronary events, and all-cause mortality in patients with and without valve calcification.....	20
Figure 7 Cumulative incidence of incident CVD, CVD death, and all-cause death in up to 15 years of follow-up in patients with MAC.....	21
Figure 8 Regional and local forces affect endothelial alignment in aortic valve...	23
Figure 9 Prevalence of AVC by diabetes mellitus, Metabolic syndrome, or neither condition, by gender.....	26
Figure 10 Cardiovascular and coronary events in patients with and without aortic valve calcifications.....	27
Figure 11 Cumulative survival for those with or without AVC.....	28
Figure 12 Multi slice computed tomography showing calcification of aorta.....	32
Figure 13 Kaplan–Meyer survival curves for all CHD events with TAC.....	34

Figure 14 Cumulative survival for subjects with or without thoracic aortic calcification	35
Figure 15 Pericardial fat and intrathoracic fat in an axial image.....	38
Figure 16 Hypothetical mechanisms whereby pericardial adipokines might play a role in coronary atherogenesis	41
Figure 17 Pericardial fat in normal fasting glucose and impaired fasting glucose subjects.	44
Figure 18 Association between PF thickness and severity of coronary artery disease.....	47
Figure 19 Comparison of PF thickness according to the presence of the metabolic syndrome (MS) and coronary artery disease.....	47
Figure 20 Major adverse cardiovascular event (MACE) prediction using Framingham Risk Score and coronary calcium score and pericardial fat volume.....	49
Figure 21 Relationship between the stenosis score and the indexed volume of pericardial adipose tissue	51
Figure 22 Mean values of PFT in normal (control) and CAD groups.....	52
Figure 23 Sensitivity and specificity for coronary artery calcium and pericardial fat as predictors of ischemia.....	54
Figure 24 Schematic drawing of the Gensini score	59
Figure 25 Measurement of pericardial fat volume by computed tomography	60
Figure 26 Correlation between DM duration and ECCS, CACS and Gensini score.....	67
Figure 27 Correlation between PFV and BMI.....	68
Figure 28 Linear correlation between CACS and Gensini score.....	72
Figure 29 Linear correlation between ECCS and Gensini score.....	73

Introduction